

Arithmetic Sequences and Series

A-level Maths Topic Summaries - Series

Subject & topics Maths Algebra Series		
Status Not started	Stage & difficulty A Level Practice 2	

Fill in the blanks below to complete the notes on arithmetic sequences and series.

Part A
Arithmetic sequences

An example of an arithmetic sequence is

3 5 7 9 11 13 ...

1st term: $u_1 = a$

2nd term: $u_2=u_1+d=a+d$


 $\text{3rd term: } u_3=u_2+d=a+2d$

•

The nth term in the sequence is given by the formula

$$u_n = a + iggl[iggl] d$$

Items:



Part B

Arithmetic series

An example of an arithmetic series is

$$3+5+7+9+11+13+...$$

An arithmetic series is a sum of terms in an arithmetic sequence. The sum of the first n terms, $S_{
m n}$, is given by

$$S_n=u_1+u_2+u_3+...+u_{
m n}$$

If the value of the last term in the series is l, this formula can also be written

$$S_n=rac{n}{2}(a+igcap)$$

Items:



Created for isaacphysics.org by Jonathan Waugh



Geometric Sequences and Series

A-level Maths Topic Summaries - Series

Subject & topics Maths Algebra Series		
Status Not started	Stage & difficulty A Level Practice 2	
Not started	A Level Flactice 2	

Fill in the blanks below to complete the notes on geometric sequences and series.

Part A	
Geometric sequences	
An example of a geometric sequence is	
$3 6 12 24 48 96 \dots$	
n a geometric sequence there is a constant ratio between successive terms. If is the first term in	n an
geometric sequence, and is the common ratio,	
1st term: $u_1=a$	
2nd term: $u_2=u_1 imes r=ar$	
3rd term: $u_3=u_2 imes r=ar^2$	
···	
The n th term in the sequence is given by the formula	
$u_n=a$	
tems:	
$egin{array}{cccccccccccccccccccccccccccccccccccc$	

Part B

Geometric series

An example of a geometric series is

$$3+6+12+24+48+96+...$$

A geometric series is a sum of terms in a geometric sequence. The sum of the first n terms, S_n , is given by

$$S_n=u_1+u_2+u_3+...+u_{\mathsf{n}}$$

or, equivalently,

$$S_n = a rac{igg(1 - r igg)}{1 - r}$$

If $\widehat{}$, the sum of the terms tends to a finite limit as n tends to infinity. The sum to infinity is given by

$$S_{\infty}= iggl($$

Items:

Created for isaacphysics.org by Jonathan Waugh

Question deck:



Sequences and Series 1i

pics		
	Stage & difficulty	
ed	A Level Practice 2	
ea	A Level Fractice 2	

A sequence of terms $u_1,\,u_2,\,u_3,\,...$ is defined by

$$u_1 = 2 \text{ and } u_{n+1} = 1 - u_n$$

for
$$n \geqslant 1$$

Part A Values			
Give the values of $u_2,\ u_3$ and $u_4.$			
$u_2 = \bigcirc$			
$u_3 = \bigcirc$			
$u_4=$			

Pa	art	В
	4	

Behaviour

Describe the behaviour of the sequence. $\qquad \qquad \text{The sequence is periodic, with a period of four. The first two values that repeat are 2 and -1.}$

It is a geometric sequence, with first term 2 and constant ratio $-\frac{1}{2}$.

The sequence is periodic, with a period of three. It cycles through values of 2, -1 and 1.

The sequence is periodic, with a period of two. It alternates between values of 2 and -1.

Part C **Sum**

Find $\sum\limits_{n=1}^{100}u_n.$


Adapted with permission from UCLES A-level Maths papers, 2003-2017.

Question deck:



Arithmetic Progression 1

Pre-Uni Maths for Sciences C2.2



In an arithmetic progression, the fifth term is 32 and the tenth term is 57.

Part A		
First term		
Find the first term, a .		
Part B		
Common difference		
Common difference		
Find the common difference, d .		


Part C Sum of f	first 70 terms			
Hence, finc	d the sum of the first 70 term	S.		

Created for isaacphysics.org by Julia Riley.

Question deck:



Arithmetic Series 2ii



The tenth term of an arithmetic progression is equal to twice the fourth term. The twentieth term of the progression is 44.

Part A First Term	
Find the first term.	
Part B Common Difference	
Find the common difference.	


Part C Sum of	the Series		
Find the s	sum of the first 50 terms.		

Used with permission from UCLES A-level Maths papers, 2003-2017.

Question deck:



Geometric Series 1ii



Records are kept of the number of copies of a certain book that are sold each week. In the first week after publication, 3000 copies were sold, and in the second week 2400 copies were sold. The publisher forecasts future sales by assuming that the number of copies sold each week will form a geometric progression with first two terms 3000 and 2400. Calculate (to the nearest number of whole books) the publisher's forecasts for:

Part A $20^{\rm th}$ Week The number of copies that will be sold in the $20^{\rm th}$ week after publication.

Part B Total copies sold in 20 weeks The total number of copies sold during the first 20 weeks after publication.

Part C Total sold copies
The total number of copies that will ever be sold.

Used with permission from UCLES A-level Maths papers, 2003-2017.

Question deck:



Geometric Series 2ii

Subject & topics Maths		
Status Not started	Stage & difficulty A Level Practice 2	

Part A $ \textbf{Geometric progression 1 - } r $
In a geometric progression, the sum to infinity is four times the first term.
Find the common ratio.

Part B Geometric progression 1 - a Given that the third term is 9, find the first term.




Part D Geometric progression 2 In another geometric progression, the first term is 6 and the sum to infinity is 10. Find the common ratio.

Adapted with permission from UCLES A-level Maths papers, 2003-2017.

Question deck:



Geometric Series 4ii



In a geometric progression, the first term is 5 and the second term is 4.8.

Part A Sum to Infinity Find the sum to infinity.

Part B Value of n

The sum of the first n terms is greater than 124. By showing that

 $0.96^n < 0.008$

and using logarithms, calculate the smallest possible value of n.

Used with permission from UCLES A-level Maths papers, 2003-2017.


Question deck:



Part B

STEM SMART Single Maths 25 - Sequences & Series

Arithmetic Series 1i



In an arithmetic progression the first term is 5 and the common difference is 3. The $n^{\rm th}$ term of the progression is denoted by u_n .

Part A ${f Value\ of\ }u_{20}$ Find the value of $u_{20}.$ The following symbols may be useful: u_20

Find the value of $\sum_{n=10}^{20} u_n$.

Part C

 ${\bf Value\ of}\ N$

Find the value of N such that $\displaystyle\sum_{n=N}^{2N}u_n=2750.$

The following symbols may be useful: N

Used with permission from UCLES A-level Maths papers, 2003-2017.