



Physics. *You work it out.*

[Home](#)

[Gameboard](#)

[Maths](#)

[Calculus: Improper Integration 1i](#)

# Calculus: Improper Integration 1i

Further A



Evaluate  $\int_0^{\infty} 2xe^{-x} dx$ .

Note that  $xe^{-x} \rightarrow 0$  as  $x \rightarrow \infty$ .

---

Adapted with permission from UCLES, A Level, Sample Paper 2017, Paper Y541, Question 5.

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



## Calculus: Improper Integration 2i

Further A



A large cuboid block is covered with a tarpaulin.

The tarpaulin is placed symmetrically over the block.

It is in contact with the block where the block is horizontal, and the shape of the tarpaulin where it hangs over the ends of the block can be modelled by the function:

$$f(x) = \frac{1}{x^2}$$




Figure 1: Diagram of the tarpaulin.

If the block is 2.00 m in length and 1.00 m in height, find the area of the cross-section of this system, which is shown in **Figure 1**.

Give your answer to 3 significant figures.

Created for isaacphysics.org by Sally A Waugh

Gameboard:

**STEM SMART Double Maths 30 - Inverse Trigonometric Calculus & Polar Coordinates**



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Integration](#) [Mean values 1](#)

# Mean values 1

A Level Further A  
C C C P P P

Find the mean values of the following.

**Part A**  $\frac{x}{\sqrt{4-x^2}}$  between 0 and 2

Find the mean value of

$$\frac{x}{\sqrt{4-x^2}}$$

between  $x = 0$  and  $x = 2$ .

**Part B**  $\frac{\sin(2\theta)}{(1-\cos^2\theta)^3}$  between  $\frac{\pi}{6}$  and  $\frac{\pi}{2}$

Find the mean value of

$$\frac{\sin(2\theta)}{(1-\cos^2\theta)^3}$$

between  $\theta = \frac{\pi}{6}$  and  $\theta = \frac{\pi}{2}$ , giving your answer in exact form.

The following symbols may be useful:  $\pi$

---

**Part C**     $\frac{1}{1+e^{-2t}}$  **between 0 and 3**

Find the mean value of

$$\frac{1}{1 + e^{-2t}}$$

between  $t = 0$  and  $t = 3$ , giving your answer in exact form.

The following symbols may be useful:  $e$ ,  $\ln()$ ,  $\log()$

---

---

**Part D**     $4b \tan \phi$  **between 0 and  $b$**

The mean value of the function  $4b \tan \phi$  between  $\phi = 0$  and  $\phi = b$  is equal to  $2 \ln 2$ , where  $b < \frac{\pi}{2}$ .  
Deduce the value of  $b$ .

The following symbols may be useful:  $b$ ,  $\pi$

---

Created for isaacphysics.org by Julia Riley

Gameboard:

**STEM SMART Double Maths 30 - Inverse Trigonometric Calculus & Polar Coordinates**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Integration](#) [Mean values 2](#)

# Mean values 2

A Level Further A  
C C C P P P

Find the mean values of the following.

**Part A**  $\sin\left(\frac{\pi x}{a}\right)$  between 0 and  $a$

Find the mean value of  $\sin\left(\frac{\pi x}{a}\right)$  between  $x = 0$  and  $x = a$ .

The following symbols may be useful:  $a$ ,  $\pi$

**Part B**  $\sin\left(\frac{\pi x}{a}\right)$  between  $-a$  and  $a$

Find the mean value of  $\sin\left(\frac{\pi x}{a}\right)$  between  $x = -a$  and  $x = a$ .

The following symbols may be useful:  $a$ ,  $\pi$

**Part C**  $\sin^2\left(\frac{\pi x}{a}\right)$  between 0 and  $a$

Find the mean value of  $\sin^2\left(\frac{\pi x}{a}\right)$  between  $x = 0$  and  $x = a$ .

The following symbols may be useful:  $a$ ,  $\pi$

---

**Part D**     $\sin^2\left(\frac{\pi x}{a}\right)$  **between  $-a$  and  $a$**

Find the mean value of  $\sin^2\left(\frac{\pi x}{a}\right)$  between  $x = -a$  and  $x = a$ .

The following symbols may be useful: a, pi

---

---

**Part E**     $x \sin^2\left(\frac{\pi x}{2a}\right)$  **between 0 and  $2a$**

Find the mean value of  $x \sin^2\left(\frac{\pi x}{2a}\right)$  between  $x = 0$  and  $x = 2a$ .

The following symbols may be useful: a, pi

---

---

**Part F**     $x \sin^2\left(\frac{\pi x}{2a}\right)$  **between  $-a$  and  $a$**

Find the mean value of  $x \sin^2\left(\frac{\pi x}{2a}\right)$  between  $x = -a$  and  $x = a$ .

The following symbols may be useful: a, pi

---

Created by for isaacphysics.org by Julia Riley

Gameboard:

**STEM SMART Double Maths 30 - Inverse Trigonometric Calculus & Polar Coordinates**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus: Inverse Trigonometry 2i](#)

# Calculus: Inverse Trigonometry 2i

Further A



## Part A Derivative of $\arcsin x$

Find the derivative of  $\arcsin x$

The following symbols may be useful:  $x$

---

## Part B Implicit differentiation

Given that

$$\arcsin 2x + \arcsin y = \frac{1}{2}\pi$$

find the exact value of  $\frac{dy}{dx}$  when  $x = \frac{1}{4}$ .

---

Adapted with permission from UCLES, A Level, January 2009, Paper 4726, Question 3.

Gameboard:

**STEM SMART Double Maths 30 - Inverse Trigonometric Calculus & Polar Coordinates**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Calculus](#) [Integration](#) [Integration using inverse trig 1](#)

# Integration using inverse trig 1

A Level Further A  
C C C P P P

Find the following integrals.

**Part A**  $\frac{1}{\sqrt{1-x^2}}$

Find the indefinite integral

$$\int \frac{dx}{\sqrt{1-x^2}}$$

using a suitable trigonometric substitution.

The following symbols may be useful: C, arccos(), arcsin(), arctan(), c, k, x

**Part B**  $\frac{5}{\sqrt{9-x^2}}$

Find the integral

$$\int_{\frac{3}{\sqrt{2}}}^{\frac{3\sqrt{3}}{2}} \frac{5dx}{\sqrt{9-x^2}}$$

giving your answer in exact form.

The following symbols may be useful: pi



---

Part C

$$\frac{2}{\sqrt{1-2x^2}}$$

Find the integral

$$\int_0^{\frac{1}{2}} \frac{2dx}{\sqrt{1-2x^2}}$$

giving your answer in exact form.

The following symbols may be useful:  $\pi$

---

Created for isaacphysics.org by Julia Riley

Gameboard:

**STEM SMART Double Maths 30 - Inverse Trigonometric Calculus & Polar Coordinates**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#)[Gameboard](#)[Maths](#)[Calculus](#)[Integration](#)[Integration using inverse trig 2](#)

# Integration using inverse trig 2

A Level Further A  
C C C P P P

Find the following integrals.

**Part A**  $\frac{1}{1+x^2}$

Find the indefinite integral

$$\int \frac{dx}{1+x^2}$$

using a suitable trigonometric substitution.

The following symbols may be useful: C, arccos(), arccosec(), arccot(), arcsec(), arcsin(), arctan(), c, k, x

**Part B**  $\frac{4}{4x^2+9}$

Find the integral

$$\int_0^\infty \frac{4dx}{4x^2+9}$$

giving your answer in exact form.

The following symbols may be useful: pi

---

Part C  $\frac{3}{x\sqrt{4x^2-1}}$

Find the integral

$$\int_1^\infty \frac{3dx}{x\sqrt{4x^2-1}}$$

giving your answer in exact form.

The following symbols may be useful: pi

---

Created for isaacphysics.org by Julia Riley

Gameboard:

**STEM SMART Double Maths 30 - Inverse Trigonometric Calculus & Polar Coordinates**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



Physics. *You work it out.*

[Home](#)   [Gameboard](#)   [Maths](#)   Polar Coordinates: General 1ii

# Polar Coordinates: General 1ii

Further A



A curve  $C$  has the cartesian equation  $x^3 + y^3 = axy$ , where  $x \geq 0$ ,  $y \geq 0$  and  $a > 0$ .

## Part A   Polar equation of $C$

Express the polar equation of  $C$  in the form  $r = f(\theta)$ .

The following symbols may be useful: `a`, `cos()`, `r`, `sin()`, `tan()`, `theta`

---

## Part B   Range of $\theta$

$$0 \leq \theta \leq \beta.$$

Find  $\beta$ .

The following symbols may be useful: `pi`

---

---

### Part C    Line of symmetry

The line  $\theta = \alpha$  is a line of symmetry of  $C$ .

Find and simplify an expression for  $r = f\left(\frac{1}{2}\pi - \theta\right)$ .

The following symbols may be useful: `a`, `cos()`, `r`, `sin()`, `tan()`, `theta`

---

Hence find the value of  $\alpha$ .

The following symbols may be useful: `pi`

---

---

### Part D    $r$ at line of symmetry

Find the value of  $r$  when  $\theta = \frac{1}{4}\pi$ .

The following symbols may be useful: `a`

---

Part E    Sketch of  $C$

Sketch the curve  $C$ .

Which curve in **Figure 1** is closest to your sketch?




Figure 1: Four curves.

- ☐ Curve P
- ☐ Curve Q
- ☐ Curve R
- ☐ Curve S

Adapted with permission from UCLES, A Level, June 11, Paper 4726, Question 4.



Physics. *You work it out.*

[Home](#) [Gameboard](#) [Maths](#) [Polar Coordinates: General 2i](#)

# Polar Coordinates: General 2i

Further A



A curve has polar equation  $r = \cos \theta \sin 2\theta$ , for  $0 \leq \theta \leq \frac{1}{2}\pi$ .

## Part A Maximum value of $r$

Find the maximum value of  $r$ .

## Part B Cartesian equation of the curve

Find a cartesian equation of the curve.

Give your answer in the form  $(x^2 + y^2)^2 = f(x, y)$

The following symbols may be useful:  $x$ ,  $y$

Adapted with permission from UCLES, A Level, June 2012, Paper 4726, Question 2.

Gameboard:

**STEM SMART Double Maths 30 - Inverse Trigonometric Calculus & Polar Coordinates**

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.



# Polar Coordinates: Area 2i

Further A



The equation of a curve, in polar coordinates, is

$$r = 2 \cos 2\theta \quad (-\pi < \theta \leq \pi).$$

## Part A Tangents at the poles

Find the values of  $\theta$  which give the directions of the tangents at the pole.

Give your answers in order of lowest to highest (most negative to most positive).

Find the lowest value,  $\theta_1$ .

The following symbols may be useful:  $\pi$

Find the second-lowest value,  $\theta_2$ .

The following symbols may be useful:  $\pi$

Find the third-lowest value,  $\theta_3$ .

The following symbols may be useful:  $\pi$

Find the highest value,  $\theta_4$ .


The following symbols may be useful:  $\pi$



---

**Part B**    **Area enclosed by one loop**

A loop of the curve is shown in the diagram.



---

**Figure 1:** One loop of  $r = 2 \cos 2\theta$ .

Find the exact value of the area of the region enclosed by the loop.

The following symbols may be useful:  $\pi$

---

Adapted with permission from UCLES, A Level, Specimen Paper, Paper 4726, Question 4.

All materials on this site are licensed under the **Creative Commons license**, unless stated otherwise.