

Section 1

Exercise 1.1.i

- (i) Let $f : x \rightarrow y$ be a morphism and $g, g' : y \rightarrow x$ be inverses. Then $fg = \text{id}_y$ and $gf = \text{id}_x$ (resp g'). Hence $gf = g'f$ so

$$\begin{aligned} gfg &= g'fg \\ g &= g' \end{aligned}$$

Therefore any morphism has at most one inverse. ///

- (ii) Suppose there are morphisms $g, h : y \rightrightarrows x$ such that $fg = \text{id}_y$ and $hf = \text{id}_x$. Then

$$\begin{aligned} hfg &= h \\ g &= h \end{aligned}$$

Hence $fg = \text{id}_y$ and $gf = \text{id}_x$ so f is an isomorphism. ///

Exercise 1.1.ii Consider a category C and the subset of isomorphisms it contains. The subset contains the identity morphisms for each object because the identity is an isomorphism. The subset inherits associativity from C . Now pick two isomorphisms $f : x \rightarrow y$ and $g : y \rightarrow z$ and consider gf . We have $(f^{-1}g^{-1})gf = f^{-1}g^{-1}gf = f^{-1}f = \text{id}_x$ and $gf(f^{-1}g^{-1}) = \text{id}_y$ so gf is also an isomorphism. Hence C contains a maximal groupoid. ///

Exercise 1.1.iii

- (i) Observe that for any object $f : c \rightarrow x$ in c/C , the morphism $\text{id}_x : x \rightarrow x$ is the identity morphism for f . Now pick two morphisms in c/C , $h : x \rightarrow y$ and $k : y \rightarrow z$ and consider kh .

$$\begin{array}{ccccc} & & c & & \\ & f \swarrow & \downarrow g & \searrow b & \\ x & \xrightarrow{h} & y & \xrightarrow{k} & z \end{array}$$

Because $hf = g$ and $kg = b$ we have $khf = b$ so kh is a morphism in c/C which is enough to show it is a category. ///

- (ii) A similar argument as above shows that C/c is a category. ///

Section 2

Exercise 1.2.ii

- (i) (\implies) Suppose $f : x \rightarrow y$ is a split epimorphism. Now pick a morphism h in $C(c, y)$. We may write

Exercise 1.2.iii Let $f : x \hookrightarrow y$ and $g : y \hookrightarrow z$ be monomorphisms and $h, k : w \rightarrow x$ be morphisms such that $gfh = gfk$. Because g is monic, it follows that $fh = fk$. Similarly, because f is monic we have $h = k$. Hence gf is monic.

Now suppose f and g are simply morphisms and that gf is monic. BWOC suppose that f is not monic. Then there exist morphisms h, k such that $fh = fk$ but $h \neq k$. Hence we have $gfh = gfk$ this is a contradiction because gf being monic implies that $h = k$. Hence f is monic.

Now observe that the monomorphisms in a category form a subcategory. Every object has an identity morphism because the identity is monic. Associativity is inherited from the larger category and composition is preserved because the composition of two monomorphisms is also a monomorphisms as shown above.

By duality the epimorphisms also form a subcategory. ///

Exercise 1.2.v Consider the inclusion map $i : \mathbb{Z} \hookrightarrow \mathbb{Q}$. Now pick two morphisms in Ring, $f, g : \mathbb{R} \rightrightarrows \mathbb{Z}$ such that $if = ig$.