



Simulation of the phenological development of wheat and maize at the global scale

L. G. J. van Bussel^{1,2*}, E. Stehfest², S. Siebert³, C. Müller⁴ and F. Ewert³

¹Plant Production Systems Group, Wageningen University, PO Box 430, NL-6700 AK, Wageningen, The Netherlands, ²Netherlands Environmental Assessment Agency, PO Box 303, NL-3720 AH, Bilthoven, The Netherlands, ³Institute of Crop Science and Resource Conservation, University of Bonn, Katzenburgweg 5, D-53115 Bonn, Germany, ⁴Earth System Analysis, Potsdam Institute for Climate Impact Research (PIK), PO Box 60 12 03, D-14412, Potsdam, Germany

ABSTRACT

Aim To derive location-specific parameters that reflect the geographic differences among cultivars in vernalization requirements, sensitivity to day length (photo-period) and temperature, which can be used to simulate the phenological development of wheat and maize at the global scale.

Location Global.

Methods Based on crop calendar observations and literature describing the large-scale patterns of phenological characteristics of cultivars, we developed algorithms to compute location-specific parameters to represent this large-scale pattern. Vernalization requirements were related to the duration and coldness of winter, sensitivity to day length was assumed to be represented by the minimum and maximum day lengths occurring at a location, and sensitivity to temperature was related to temperature conditions during the vegetative development phase of the crop.

Results Application of the derived location-specific parameters resulted in high agreement between simulated and observed lengths of the cropping period. Agreement was especially high for wheat, with mean absolute errors of less than 3 weeks. In the main maize cropping regions, cropping periods were over- and underestimated by 0.5–1.5 months. We also found that interannual variability in simulated wheat harvest dates was more realistic when accounting for photoperiod effects.

Main conclusions The methodology presented here provides a good basis for modelling the phenological characteristics of cultivars at the global scale. We show that current global patterns of growing season length as described in cropping calendars can be largely reproduced by phenology models if location-specific parameters are derived from temperature and day length indicators. Growing seasons can be modelled more accurately for wheat than for maize, especially in warm regions. Our method for computing parameters for phenology models from temperature and day length offers opportunities to improve the simulation of crop productivity by crop simulation models developed for large spatial areas and for long-term climate impact projections that account for adaptation in the selection of varieties.

Keywords

Agricultural management, crop calendars, cultivar/variety characteristics, global crop modelling, global harvest dates, phenology.

*Correspondence: L. G. J. van Bussel, Plant Production Systems Group, Wageningen University, PO Box 430, NL-6700 AK, Wageningen, The Netherlands.
E-mail: lenny.vanbussel@wur.nl

INTRODUCTION

The phenological development of crops determines the duration and timing of essential periods for crop growth, and thus to

a large extent the quality and quantity of crop yields (Porter & Semenov, 2005). In the scientific literature it is well established that temperature (directly and in the case of winter cultivars also indirectly via vernalization requirements) and day length (also

referred to as photoperiod) are the main components determining crop development. The development rate shows a positive linear correlation with temperature (Slafer & Rawson, 1994). Vernalization is the influence of cold temperatures on the timing of flowering (Raven *et al.*, 2005): development is delayed until the plant has experienced sufficient days at vernalizing temperatures (Miralles & Slafer, 1999). Finally, photoperiodism is the response to a change in the proportions of light and darkness in a 24-h cycle. For long-day plants (e.g. wheat) development accelerates when the photoperiod increases, for short-day plants (e.g. maize) development accelerates when the photoperiod decreases (Raven *et al.*, 2005). Moreover, crop development will be synchronized between years due to photoperiodism, buffering the impact of interannual variability in weather conditions (Hay & Kirby, 1991; Gouesnard *et al.*, 2002; Craufurd & Wheeler, 2009). Crops like wheat and maize are cultivated in a wide range of environments (Gouesnard *et al.*, 2002; Trethowan *et al.*, 2006). This is possible due to a variety of cultivars that respond differently to temperature and photoperiod; farmers can therefore choose a cultivar that is well adapted to local temperature and photoperiod conditions (Olesen *et al.*, 2012).

Observed climatic warming has resulted in changes in crop phenology, which has been described for Europe (Menzel *et al.*, 2006), China (He *et al.*, 2015) and the USA (Twine & Kucharik, 2009), with associated consequences for crop productivity. Crop simulation models are suitable tools for assessing the impact of climate change on crop productivity. The accuracy of simulated crop yield is determined to a large extent by the accuracy of the phenological module of the model (Wheeler *et al.*, 2000). Moreover, potential adaptations of farmers to climate change through the selection of cultivars and sowing dates should be considered when assessing the impacts of climate change (Olesen *et al.*, 2012). Accordingly, static crop calendars, for example based on observations, should not be used as input for studies on adaptation or climate impact.

Given the widely reported sensitivity of crops to photoperiod, its role as a key determinant of the adaptation of a crop to local conditions, and its synchronizing function, it seems important to consider its effects in the phenological modules of crop models (Craufurd & Wheeler, 2009). Vernalization is often, but not always, considered in field-scale (Asseng *et al.*, 2013) and global-scale (Rosenzweig *et al.*, 2014) wheat models, while photoperiodism is covered by many field-scale models for wheat (Asseng *et al.*, 2013) and maize (Bassu *et al.*, 2014), but typically not in global-scale models (Rosenzweig *et al.*, 2014). One reason is certainly the lack of information required to parameterize photoperiodism and vernalization responses of crops in phenology models applied at global scale. The consequences of omitting photoperiodism and vernalization relations for phenology simulations, for example on the interannual variability of the length of the simulated cropping periods, have not been evaluated at the global scale to date.

As stated above, farmers choose cultivars adapted to local conditions, but also as an adaptation strategy to changing cli-

matic conditions. Most global crop growth models use climate data to define sowing dates (see, e.g., Waha *et al.*, 2012), thus accounting for possible adaptation strategies. However, to date, equivalent methods to define the model parameters necessary to account for cultivar selection as an adaptation strategy are lacking for the global scale. This limits simulation of phenological development and consequently hinders simulation of accurate crop yields and variety-related adaptation strategies, especially within climate impact assessments.

The objectives of this study were therefore: (1) to compare phenology models that differ with regard to consideration of vernalization and/or photoperiod responses; (2) to develop algorithms to define location-specific crop variety parameters for these phenology models; and (3) to investigate the interannual variability in simulated growing period lengths based on the different phenology models. Wheat and maize are used as example crops as they account for approximately 30% of the total harvested global crop area (FAO, 2012). We thus hypothesize that farmers' selection of crop varieties is mainly aimed at optimizing the growing period as defined by local climatic conditions and that the parameters for phenology models can be derived from simple climatic indicators.

The outcomes of this study can help to improve simulations of crop phenology in crop growth models applied over large spatial areas. This will make an important contribution to studies of the impact of climate on global crop productivity.

METHODS

The phenological model AFRCWHEAT2

To simulate the phenological development of wheat and maize we used the well-established concept of heat units as implemented in the model AFRCWHEAT2 (Weir *et al.*, 1984; Porter, 1993; Ewert *et al.*, 1996). Daily temperature (T_i , °C) is accumulated above a base temperature (T_b , °C), resulting in a heat unit sum (HU_{sum} , °C day) until the required heat units from emergence to physiological maturity (HU_{req} , °C day) are reached. The increment in heat units is modified by the effects of photoperiod (photoperiod factor, P_i), and in case of winter wheat by the effects of vernalization (vernalization factor, V_i). The crop is harvested when the heat unit sum accumulated for the period since sowing, HU_{sum} , becomes equal to or larger than HU_{req} :

$$HU_{sum} = \sum_{i=1}^N (T_i - T_b) \times P_i \times V_i \quad \text{if } HU_{sum} \leq HU_{req} \quad (1)$$

with N the simulated length of the cropping period in days. The simulation of maize was in addition stopped when temperatures dropped continuously below the base temperature. Further details on the phenology model and the required parameters are provided in Appendix S1 in the Supporting Information.

Data

Datasets of observed cropping periods

The global dataset MIRCA2000

To derive regression parameters for estimating heat unit requirements and to test how well application of the obtained regression equations at the global scale could reproduce spatial patterns of observed phenology, we used the global dataset MIRCA2000 (Portmann *et al.*, 2010). MIRCA2000 reports monthly growing areas of 24 irrigated and rainfed crops at a spatial resolution of $5' \times 5'$ for the period around the year 2000. The data were aggregated to the resolution of the weather data used in this study ($30' \times 30'$).

Up to five possible cropping periods per grid cell are indicated in MIRCA2000, reflecting different cultivars of wheat and multiple cropping systems with maize. For each grid cell, the cropping period with the maximum reported area was selected for this study. We assumed that sowing was at the first day of the reported sowing month, also applied by Portmann *et al.* (2010), and harvest at the last day of the reported harvest month. Day of emergence was set equal to the sowing date and the harvest date was assumed to correspond to physiological maturity.

A dataset similar to MIRCA2000 was developed by Sacks *et al.* (2010) but they did not differentiate between rainfed and irrigated crops. Since cropping periods often differ between irrigated and rainfed crops we used MIRCA2000 for this study.

Point observations of winter wheat harvest dates

Information on the interannual variability of harvest dates was derived from the Pan-European Phenology Database (PEP725) which provides time series of harvest dates of winter wheat observed for several European countries. From this database we first selected countries that had data available for at least 10 successive years. Next, within these countries, we selected only locations with at least five observations within a 10-year period (the 10-year period was kept constant per country).

Photoperiod and temperature

Daily photoperiod (P , h day⁻¹) was calculated based on latitude and day of the year (Monteith & Unsworth, 1990) and monthly mean temperature data on a $30' \times 30'$ resolution was extracted from the Climate Research Unit TS3.0 dataset (Mitchell & Jones, 2005) using linear interpolation to obtain daily temperature values.

Computation of location-specific phenological parameter values

For the computation of the location-specific phenological parameter values, we assumed (1) that location-specific parameters represent cultivars that are best adapted to local climatic and photoperiodic conditions and (2) that farmers base their

cultivar selection on experiences with climatic conditions in previous years, using exponentially weighted moving averages of monthly temperatures ($\bar{T}_{m,y}$, °C; see Appendix S2). These location-specific parameters can be applied to regions not currently used for the cultivation of the crop and used to estimate how crop cultivar selection may change under climate change.

Vernalization requirements of winter wheat cultivars

Ewert *et al.* (1996) indicated that both the effectiveness of temperature on the vernalization process and the vernalization requirements are different among winter cultivars. Because of the scarcity of data we assumed equal temperature effectiveness for all cultivars and only varied the vernalization requirements (V_{sat} and V_b , in days; see Appendix S1) to characterize cultivars.

Tolerance to below-freezing temperatures is built up due to the exposure of winter wheat to vernalizing temperatures. Once the vernalization requirements are met the tolerance gradually disappears (Mahfoozi *et al.*, 2001). Vernalization requirements of cultivars grown at locations with long, cold winters are higher than those of cultivars grown at locations with milder winters (Iwaki *et al.*, 2001). Therefore, we used the temperatures of the five coldest months of the year to characterize winter duration and coldness, assuming that in winter wheat growing regions the frost period is 5 months long at most. By considering the five coldest months separately, influences of possible relatively warm months were minimized (Appendix S3). We excluded higher-latitude regions where MIRCA2000 indicated a sowing date during the spring, since this sowing date suggests the use of spring wheat cultivars.

Photoperiodism sensitivity

To simulate differences among wheat and maize cultivars with respect to photoperiod, we adapted P_{opt} values to local conditions as described below, while P_b was kept constant among the cultivars, using values from the literature (see Table S1, Appendix S1).

Wheat

Wheat cultivars originating from high latitudes (e.g. the UK or Finland) are true photoperiod-sensitive cultivars (Worland *et al.*, 1994), while modern cultivars grown in lower latitudes such as in the Mediterranean region and North Africa are photoperiod insensitive (Ortiz Ferrara *et al.*, 1998). Miralles and Slafer (1999) indicated that for wheat cultivars with different photoperiod sensitivities the optimum photoperiod differs significantly, ranging from c. 15 to 21 h day⁻¹. For Argentinean wheat cultivars the optimum photoperiod was found to be 13.4 h day⁻¹ (Miralles *et al.*, 2007), which approximately coincides with the maximum photoperiod (P_{max} , h day⁻¹; i.e. P at 21 December) on average in the main wheat growing area in Argentina. We characterized sensitivity to photoperiod by setting P_{opt} to the location-specific P_{max} , i.e. P at 21 June and 21 December in the Northern and Southern Hemisphere, respectively.

Maize

The difference in photoperiod between two successive days is smaller at lower latitudes than at higher latitudes. It is therefore plausible that tropical cultivars of short-day plants like maize are most sensitive to photoperiod (Summerfield *et al.*, 1997). Indeed, maize cultivars adapted to temperate regions (i.e. cool, long-day environments) show lower or no sensitivity to photoperiod compared with tropical cultivars (Bonhomme *et al.*, 1994; Birch *et al.*, 1998). Similar to wheat cultivars, maize cultivars also differ in their optimum photoperiod (Birch *et al.*, 1998). Rood and Major (1980) found optimum photoperiods varying from < 14 to 24 h day⁻¹. Location-specific P_{opt} values for maize were established as follows:

$$P_{\text{opt}} = \max \left[0, P_{\text{max}} \times \left(1 - \frac{1}{P_{\text{max}} - P_{\text{min}}} \right) \right] \quad (2)$$

where P_{min} (h day⁻¹) is the minimum photoperiod possible at a certain location, i.e. P at 21 December and 21 June in the Northern and Southern Hemisphere, respectively.

Heat units required from emergence to maturity

Heat units from emergence to maturity (HU_{req} , °C day) could not be derived from the literature in a general fashion applicable to global-scale studies because estimates from case studies are model specific, depending on whether sensitivity to photoperiod and vernalization was considered and depending on which parameters were used to describe this sensitivity. We therefore estimate HU_{req} with linear regression models between HU_{req} calculated by using observed MIRCA2000 crop calendar data as the dependent variable and, as the independent variable, available heat units during an estimated vegetative cropping period ($HU_{\text{sum veg-period}}$, °C day), in line with Bignon (1990). Due to the spatial resolution of MIRCA2000, sowing dates are also reported in grid cells which are too cold for growing maize. We therefore excluded grid cells with $HU_{\text{sum year}} < 750$ °C day from the regression analysis. To account for differences between temperate and tropical maize cultivars, we calculated separate linear regression models for maize cultivars grown in warm regions ($HU_{\text{sum year}} \geq 3000$ °C day) and for maize cultivars grown in cold regions ($HU_{\text{sum year}} < 3000$ °C day).

Regression models were established separately for a thermal model, taking into account only temperature effects, and a photo-thermal model, taking into account temperature effects combined with photoperiod effects, including the cultivar differences as described above. For winter wheat two additional regression models were distinguished: temperature effects combined with vernalization effects (vernal-thermal model) and temperature effects combined with vernalization and photoperiod effects (vernal-photo-thermal model). For these calculations, it was assumed that wheat and maize are sensitive to photoperiod until flowering, and winter wheat was assumed to be sensitive to vernalization until V_{sat} requirements are met. To calculate available heat units during the vegetative cropping period ($HU_{\text{sum veg-period}}$, °C day) we assumed a vegetative period

Table 1 Coefficients of determination from the linear regression analysis with the available heat units during an estimated vegetative cropping period ($HU_{\text{sum veg-period}}$), as well as the accompanying equations to compute location-specific HU_{req} values based on $HU_{\text{sum veg-period}}$. The regression models were applied at the global scale with the same set of parameters. For maize we estimated the vegetative period from May to July in the Northern Hemisphere and November to January in the Southern Hemisphere; for wheat from March to June in the Northern Hemisphere and September to December in the Southern Hemisphere. For maize we distinguished between warm and cold regions using $HU_{\text{sum year}}$, we considered a location to be warm if $HU_{\text{sum year}} \geq 3000$ °C day, if $HU_{\text{sum year}} < 3000$ °C day we considered it to be cold.

Crop	Equations to calculate HU_{req}	R^2
	Thermal model	$HU_{\text{sum veg-period}}$
Spring wheat	$HU_{\text{req}} = 1.06 \times HU_{\text{sum veg-period}} + 815.08$	0.75
Winter wheat	$HU_{\text{req}} = 1.18 \times HU_{\text{sum veg-period}} + 941.45$	0.45
Maize	Warm region: $HU_{\text{req}} = 1.4 \times HU_{\text{sum veg-period}} + 399.66$	0.44
	Cold region: $HU_{\text{req}} = 1.82 \times HU_{\text{sum veg-period}} - 150.51$	0.92
	Photo-thermal model	
Spring wheat	$HU_{\text{req}} = 0.91 \times HU_{\text{sum veg-period}} + 775.27$	0.64
Winter wheat	$HU_{\text{req}} = 0.66 \times HU_{\text{sum veg-period}} + 1126.49$	0.26
Maize	Warm region: $HU_{\text{req}} = 1.06 \times HU_{\text{sum veg-period}} + 145.04$	0.35
	Cold region: $HU_{\text{req}} = 1.52 \times HU_{\text{sum veg-period}} - 72.08$	0.91
	Vernal-thermal model	
Winter wheat	$HU_{\text{req}} = 0.99 \times HU_{\text{sum veg-period}} + 811.52$	0.44
	Vernal-photo-thermal model	
Winter wheat	$HU_{\text{req}} = 0.87 \times HU_{\text{sum veg-period}} + 907.47$	0.36

from May to July in the Northern Hemisphere and November to January in the Southern Hemisphere for maize and from March to June in the Northern Hemisphere and September to December in the Southern Hemisphere for wheat, respectively. Regression parameters and R^2 are shown in Table 1 for each of the models.

Assessment of different phenological models

Assessment of regression model skill

Three area-weighted indices of agreement were computed to assess the degree of agreement between simulated and observed cropping period lengths: the mean absolute error (MAE, months), the root mean square error (RMSE, months), and the Willmott coefficient of agreement (W , dimensionless, ranging from 0 to 1, with 1 showing perfect agreement) (Willmott, 1982). MAE and RMSE indicate the global average error between simulations and observations. In addition, W is a relative measure for the differences (Willmott, 1982):

$$\text{MAE} = \frac{\sum_{i=1}^N |O_i - S_i| \times A_i}{\sum_{i=1}^N A_i} \quad (3)$$

$$\text{RMSE} = \sqrt{\frac{\sum_{i=1}^N (O_i - S_i)^2 \times A_i}{\sum_{i=1}^N A_i}} \quad (4)$$

$$W = 1 - \frac{\sum_{i=1}^N (O_i - S_i)^2 \times A_i}{\sum_{i=1}^N (|S_i - \bar{O}| + |O_i - \bar{O}|)^2 \times A_i} \quad (5)$$

where S_i (months) is the simulated and O_i (months) the observed length of the cropping period in spatial unit i , $\bar{O}$ (months) is the mean observed length of the cropping period based on all spatial units, A_i (ha) is the cultivated area of the crop in spatial unit i , and N is the number of spatial units.

Comparison of simulated interannual variability in harvest dates

To assess if accounting for the effects of photoperiod and vernalization results in less interannual variability in the length of simulated wheat cropping periods than simulation of phenology based on thermal requirements alone, we used the different phenology models (thermal, photo-thermal, vernal-thermal, vernal-photo-thermal) that were parameterized with location-specific parameters derived from the linear regression analysis, to simulate the lengths of the cropping period for the period 1996 to 2005. As an indicator for interannual variability we calculated for each grid cell and model the differences between the latest and earliest of the simulated harvest dates in that period.

To compare the interannual variability in observed and simulated harvest dates we calculated per location selected from the

PEP725 database the standard deviation of the observed harvest dates (Fig. S2, Appendix S4). Next we calculated the country average of these standard deviations and the average standard deviations of simulated harvest dates from the different models.

RESULTS

Calculated phenological model parameters

Vernalization requirements

For cultivars grown in large parts of Russia, western Europe, northern USA, and north-east Asia the maximum possible required duration of exposure to vernalizing temperatures was computed. South of these regions lower vernalization requirements were simulated (Fig. 1).

Heat units required from emergence to maturity

The simulated heat units that are required for the phenological development from emergence to maturity show a clear trend from north to south: southern regions grow varieties that require more heat units from emergence to maturity than the varieties in northern regions (Fig. 2 shows the HU_{req} for the vernal-photo-thermal wheat model and the photo-thermal maize model). The range in HU_{req} is larger for wheat than for maize.

Comparison of simulated and observed cropping periods

Wheat

For wheat, the agreement between observed and simulated cropping periods was in the same range for the different




Figure 1 Simulated location-specific vernalization requirements (i.e. required duration of exposure to vernalizing temperatures; V_{sat} , days) for winter wheat (Mollweide projection).




Figure 2 Simulated location-specific heat units from emergence to maturity (HU_{req} , °C day) for (a) the vernal-photo-thermal wheat model and (b) the photo-thermal maize model (Mollweide projection). Patterns reflect that varieties with high heat unit requirements are grown in warm locations and varieties with low heat unit requirements are grown in cooler locations.

Table 2 Area-weighted means, coefficient of variation of spatial distribution and indices of agreement between simulated and observed cropping periods for wheat and maize.

Crop	Mean (months)*	Coefficient of variation (–)	Mean absolute error (months)*	Root mean square error (months)*	Willmott coefficient of agreement (–)
Observations					
Wheat	8.7	3.8	–	–	–
Maize	5.7	2.1	–	–	–
Thermal model					
Wheat	8.6	3.6	0.73	0.92	0.97
Maize	5.5	2.1	0.75	0.92	0.54
Photo-thermal model					
Wheat	8.7	3.6	0.77	0.96	0.97
Maize	5.5	2.1	0.89	1.03	0.42
Vernal-thermal model					
Wheat	8.7	3.7	0.66	0.82	0.98
Vernal-photo-thermal model					
Wheat	9.0	3.6	0.75	0.97	0.97

*Recalculation from days to months is done by assuming an equal number of days per month (i.e. $365/12 = 30.42$ days per month).

phenological models (Table 2), with the vernal-thermal model giving the highest agreement and the photo-thermal model the lowest agreement. The mean simulated lengths of the cropping periods were in the same range as the observed mean; the spatial heterogeneity was slightly lower in the simulated cropping periods in comparison with the observed cropping periods. Agreement between observed and simulated lengths of cropping periods of countries with large wheat cropping areas, such as Russia, Canada and Turkey, was high. Figure 3(a) gives results of

the vernal-photo-thermal model; scatterplots for the other models look similar and are not shown.

Maize

For maize, agreement between observed and simulated cropping periods was slightly higher based on the thermal model than when based on the photo-thermal model, but both were lower than for wheat. The simulated cropping periods underestimated the observed cropping period by approximately 1 week on average; also the spatial heterogeneity in cropping period was underestimated by the simulations (Table 2). The scatterplot (Fig. 3b for the photo-thermal model) indicates that in countries situated in warmer regions, for example Mexico and Nigeria, the lengths of the cropping periods were overestimated, while in cooler regions, for example the USA and Russia, the lengths of the cropping periods were underestimated.

Comparison of simulated interannual variability in harvest dates

As an indication of the interannual variability in simulated harvest dates we calculated, per gridcell, the differences between the latest and earliest of the 10 simulated harvest dates. For comparison we plotted in Fig. 4 the cumulative frequency distributions of these differences per model. The initially steeper slopes of the vernal-photo-thermal and photo-thermal models for wheat in Fig. 4(a) indicate that accounting for photoperiodism in particular results in less variability in harvest dates among years. Accounting for vernalization only slightly reduced the interannual variability in harvest dates. We also mapped the spatial distribution of the difference between the latest and earliest of the simulated harvest dates of the thermal




Figure 3 Scatterplots of observed versus simulated cropping periods per spatial unit of MIRCA2000 for (a) wheat, based on the vernal-photo-thermal model and (b) maize, based on the photo-thermal model. The solid line represents the 1:1 line; the radius of the circles is proportional to the cultivated area of the crop in the corresponding spatial unit.




Figure 4 Relative cumulative frequency distributions of the differences between the latest and earliest of the simulated harvest dates in the period 1996 to 2005 for (a) wheat and (b) maize. The lines indicate the different models used to simulate the length of the cropping period. The arrows in the wheat plot indicate that if phenological development is simulated including photoperiod effects, the differences in 75% of the grid cells between the latest and earliest simulated harvest dates within the 10-year period is at most 18 days, while for the thermal model this is at most 26 days.

(Fig. 5a) and vernal-photo-thermal (Fig. 5b) models in the period 1996 to 2005. To quantify the interannual variability in temperature conditions the coefficient of variation of the available $HU_{sum\ veg-period}$ within the same period was calculated (Fig. 5c). The maps indicate that particularly in areas with high interannual variability in temperature conditions, combined with cultivars sensitive to photoperiod, such as the north-eastern USA and northern and western Europe, interannual variability in harvest dates decreases due to the inclusion of photoperiod effects.

For maize we found that including the effects of photoperiodism only slightly changed the interannual variability of simulated harvest dates (Fig. 4b).

A direct quantitative comparison between the interannual variability in observed and simulated wheat harvest dates was not possible, for various reasons. First of all, the spatial scale of the observed and simulated harvest dates differed (point versus $0.5^\circ \times 0.5^\circ$ resolution). Secondly, besides variation in observed harvest dates due to variation in interannual weather conditions, it is likely that in the real-world observations additional variation occurs because of several other factors, for example delay in harvesting due to unfavourable soil conditions or lack of machinery for timely harvesting. Despite these differences, the values in Table 3 show that for four out of the six countries interannual variability in harvest dates was simulated more in

line with observed variations when using models including photoperiod than with thermal-only models.

DISCUSSION

Comparison of simulated and observed cropping periods at the global scale

We present in this study an evaluation of the effects of including photoperiod and temperature (directly and indirectly) effects in crop phenology simulations at the global scale. We developed algorithms to compute location-specific parameter values that account for differences among cultivars in vernalization requirements and sensitivity to photoperiod and temperature. Accounting for photoperiod and vernalization effects in the phenology model decreases the spatial heterogeneity of heat units required from emergence to physiological maturity for wheat, a result in line with Miralles and Slafer (1999) who reported that differences in the rate of development among wheat cultivars mainly originate from differences in sensitivity to photoperiod and vernalization requirements.

The accuracy of simulated cropping period lengths did not improve if the phenology models accounted for the effects of photoperiod and/or vernalization (Table 2). However, accounting for effects of photoperiod and vernalization decreased the




Figure 5 Differences between the latest and earliest of the simulated wheat harvest dates in the period 1996 to 2005 for (a) the thermal model and (b) the vernal-photo-thermal model. (c) Coefficient of variation of the available heat units during the estimated vegetative cropping period ($HU_{\text{sum veg-period}}$) in the period 1996 to 2005 for wheat (Mollweide projection). The coefficient of variation is used to characterize interannual variability in temperature conditions: regions with high coefficients of variation are characterized by high interannual variability in temperature conditions.

Table 3 Observed winter wheat data specifications and standard deviation of observed and simulated harvest dates of winter wheat. Figure. S2 gives the locations of the observations.

Country	Time period observations	Total number of observations	Number of locations with ≥ 5 years of observations	Standard deviation observed and simulated harvest dates (days)				
				Observations	Thermal model	Photo-thermal model	Vernal-thermal model	Vernal-photo-thermal model
Austria	1996–2005	178	24	8.08	13.81	10.00	11.01	9.20
Belgium	1989–1998	95	13	7.61	10.98	4.98	9.14	5.09
Croatia	1996–2005	40	4	9.28	8.33	5.11	6.92	5.04
Czech Republic	2001–2010	20	2	6.20	10.22	6.05	8.77	6.45
Germany	1995–2004	11,560	1352	8.70	10.85	5.44	10.47	5.82
Slovakia	2000–2009	456	56	8.98	9.14	5.66	7.43	5.81

simulated interannual variability in the length of cropping periods, especially for wheat (Fig. 4). As such, we concluded from the comparison with the PEP725 dataset that the interannual variability in wheat harvest dates was probably simulated more realistically by the models that include photoperiod. The comparison is complicated, however, by not taking into account in the simulations the variability in sowing dates.

Still, we consider that the reduced variability in cropping periods makes a strong case for considering photoperiod in phenology models which are applied for large spatial areas. For maize we could not determine whether the inclusion of photoperiodism has added value, because there is no observational dataset like PEP725 for maize available for tropical regions. Also growing seasons in the tropics are often

determined by non-climatic factors, so a comparison is deemed to be difficult here.

The indices of agreement (Table 2) and the scatterplots (Fig. 3) indicated that, in general, agreement between simulations and observations was higher for wheat than for maize. This reduced agreement, despite good fits of the regression models for maize especially in the cooler regions (Table 1), can be explained by the differences in harvest seasons between wheat and maize. In temperate regions, wheat is normally harvested during the warmest period of the year, while maize is harvested in autumn, when temperatures approach the base temperature of maize. As a consequence, an equal over- or underestimation of the heat unit requirements leads to higher deviations in simulated harvest dates for maize than for wheat.

Calculated vernalization requirements

The pattern of simulated vernalization requirements (Fig. 1) corresponds with information found in the literature, for example in the south-eastern border of the Australian wheat belt winter wheat is grown, while in rest of the belt spring wheat is grown (Fisher, 1999). Also the low computed vernalization requirements for wheat in west Asia (e.g. Yemen and Saudi Arabia) and North Africa are in line with results of previous studies (see, e.g., Ortiz Ferrara *et al.*, 1998). Kato and Yokoyama (1992) determined the vernalization requirements of traditional cultivars originating from various countries. They found vernalization requirements of approximately 31 days for landraces originating from western Turkey, Italy and Greece; 35 days for landraces from Afghanistan, Pakistan, Nepal and Bhutan; 56 days for landraces from Georgia, east Turkey and north and east Iran; 28 days for landraces from Armenia; and 7 and 14 days for landraces from Egypt and Ethiopia. This pattern is reflected in our results and builds trust in the relatively simple relationship between vernalization days and temperatures of the coldest five months of the year.

Simulated interannual variability in harvest dates

To evaluate the effect of vernalization and photoperiod on the interannual variability of harvest dates we compared time series of harvest dates, simulated with different phenology models, with observations from six European countries (Table 3). We found in particular that the consideration of photoperiod resulted in a decrease of the interannual variability of winter wheat harvest dates. However, the mean standard deviation of simulated winter wheat harvest dates was even smaller when using the photo-thermal model or the vernal-photo-thermal model as compared with the observations, in particular for the two countries where most of the observations were obtained (Germany and Slovakia). An explanation for this difference could be that the observations were collected by a network of voluntary observers so that errors (e.g. due to confusion of crops, data handling or a low frequency of field visits) in the observational dataset are likely. It is very difficult to distinguish 'true' from potentially 'wrong' observations, but the correspond-

ing outliers may contribute substantially to the standard deviation in the observed harvest dates. Therefore, standard deviation of harvest days is probably overestimated in the observations.

The small differences in interannual variability of simulated maize harvest dates between the thermal and photo-thermal models is a consequence of low sensitivity to photoperiod in areas with high interannual variability in temperature conditions (e.g. western Europe), while cultivars with high sensitivity to photoperiod are grown in areas with lower interannual variability in temperature (tropical regions).

Practical implications of our research

Our approach yields two major advantages of large-scale crop modelling studies. Firstly, the simple but robust relationship between cultivar parameters that determine the length of the growing period and simple climatic indicators allows for out-scaling of crop cultivar parameters to areas currently not used for cultivation of these crops. This is important to inform simulations of gridded crop models that feed data to economic land-use models (Müller & Robertson, 2014). Secondly, our approach also allows us to assess possible changes in cultivar choice under climate change, as crop cultivar parameters can be derived from climatic indicators and can thus inform climate adaptation studies and provide information that cannot be derived from observed cropping calendars (see Appendix S5 for a comparison between growing period lengths simulated for future climatic conditions based on non-adapted inputs and inputs based on our methodology). Moreover, our methodology can be applied in combination with the methodology developed by Waha *et al.* (2012) to determine sowing dates using climatic data independently of fixed cropping calendars such as MIRCA2000 or Sacks *et al.* (2010).

Limitations of our methodology and directions for further research

Only two datasets (Portmann *et al.*, 2010; Sacks *et al.*, 2010), with roughly the same data sources, report on global sowing and harvest dates, but only Portmann *et al.* (2010) distinguish between irrigated and rainfed crops. Literature reporting reasonable ranges of observed HU_{req} across the world is not available and case studies are difficult to compare, given differences in underlying phenology models. Calibration of the different phenology models was therefore carried out for HU_{req} values based on the dataset of Portmann *et al.* (2010), and evaluation of how well the global-scale application of the regression equations shown in Table 1 reproduced the spatial pattern in observed crop phenology was done with the same dataset. Being aware of this limitation, we still found it valid to test whether observed patterns of harvest dates could be reproduced when cultivar-specific parameters were computed based on average climatic conditions only. Clearly, we did not aim to evaluate the concept of heat units itself, rather our methodology to compute location-specific cultivar parameters used within the heat unit concept.

We assumed in this study that cultivar selection is made only on the basis of local photoperiod and temperature conditions. The choice of farmers whether to grow a certain cultivar with specific characteristics is, however, also dependent on numerous other factors, including particulars of the production system, such as multiple-cropping systems that require cultivars with a specific cropping period length (Tao *et al.*, 2014), the demand for specific cultivars for quality reasons, such as durum wheat in Italy (Dettori *et al.*, 2011), or to avoid pests and diseases (Kouressy *et al.*, 2008). Currently this complex system of cultivar choice is not well understood, therefore we could not include these factors in our methodology. Insights into the complex issues of cultivar choice will help to improve simulation of crop phenology for large spatial areas, but could also assist in improving studies related to possible measures for adapting to climate change (see, e.g., Olesen *et al.*, 2012). Finally, the interaction of simulated sowing dates (see, e.g., Waha *et al.*, 2012) with simulated cultivar traits, including the effects of among others irrigation and increasing soil temperatures, needs further evaluation.

Our approach allows accurate derivation of parameters for wheat phenology models, but for maize the approach has more limitations. This could be due to a more complex phenological response to temperature in maize (Kumudini *et al.*, 2014), oversimplistic assumptions about the development phase that is sensitive to photoperiod (Birch *et al.*, 1998), as well as to overestimation of the effect of photoperiod in tropical regions where the longest and shortest days are not very different (see equation 2 and Table 1). These limitations for maize need to be considered when employing our approach, and we suggest that the thermal model is used for maize. However, the thermal model approach may still be a valid alternative to assuming static variety distributions in long-term climate impact assessments or even simpler temperature-based response functions (Rosenzweig *et al.*, 2014).

A limitation of MIRCA2000 was the temporal resolution of 1 month. By assuming sowing to be on the first day and harvest to be on the last day, it is likely that we have overestimated the simulated and observed growing periods, possibly by 2 months. To further improve the simulation of crop phenology for the globe we stress the importance continuously expanding datasets such as MIRCA2000 over time and including more spatial detail. Moreover, to make it possible to understand the complex system of cultivar choice, we stress the importance of collecting cultivar characteristics, including the timing of vulnerable stages such as anthesis, and cultivar use for as many (contrasting) locations as possible.

ACKNOWLEDGEMENTS

We would like to thank the members of the PEP75 project for providing the point observations of the winter wheat harvest dates and Felix Portmann for making available the MIRCA2000 dataset. Comments of two anonymous referees greatly improved a previous version of the manuscript. C.M. acknowledges financial support from the MACMIT project (01LN1317A) and the

FACCE MACSUR project (031A103B) funded through the German Federal Ministry of Education and Research (BMBF).

REFERENCES

- Asseng, S., Ewert, F., Rosenzweig, C. *et al.* (2013) Uncertainty in simulating wheat yields under climate change. *Nature Climate Change*, **3**, 827–832.
- Bassu, S., Brisson, N., Durand, J.-L. *et al.* (2014) How do various maize crop models vary in their responses to climate change factors? *Global Change Biology*, **20**, 2301–2320.
- Bignon, J. (1990) *Agrometeorologie et physiologie du maïs grain dans la communauté Européenne*. Commission des Communautés Européennes, Luxembourg.
- Birch, C.J., Hammer, G.L. & Rickert, K.G. (1998) Temperature and photoperiod sensitivity of development in five cultivars of maize (*Zea mays* L.) from emergence to tassel initiation. *Field Crops Research*, **55**, 93–107.
- Bonhomme, R., Derieux, M. & Edmeades, G.O. (1994) Flowering of diverse maize cultivars in relation to temperature and photoperiod in multilocation field trials. *Crop Science*, **34**, 156–164.
- Craufurd, P.Q. & Wheeler, T.R. (2009) Climate change and the flowering time of annual crops. *Journal of Experimental Botany*, **60**, 2529–2539.
- Dettori, M., Cesaraccio, C., Motroni, A., Spano, D. & Duce, P. (2011) Using CERES-Wheat to simulate durum wheat production and phenology in southern Sardinia, Italy. *Field Crops Research*, **120**, 179–188.
- Ewert, F., Porter, J. & Honermeier, B. (1996) Use of AFRCWHEAT2 to predict the development of main stem and tillers in winter triticale and winter wheat in North East Germany. *European Journal of Agronomy*, **5**, 89–103.
- FAO (2012) FAOSTAT Available at: <http://faostat.fao.org/default.aspx> (accessed 17 April 2012).
- Fisher, R.A. (1999) Wheat cropping in Australia. *Wheat ecology and physiology of yield determination* (ed. by E.H. Satorre and G.A. Slafer), pp. 277–294. Food Products Press, New York.
- Gouesnard, B., Rebourg, C., Welcker, C. & Charcosset, A. (2002) Analysis of photoperiod sensitivity within a collection of tropical maize populations. *Genetic Resources and Crop Evolution*, **49**, 471–481.
- Hay, R.K.M. & Kirby, E.J.M. (1991) Convergence and synchrony – a review of the coordination of development in wheat. *Australian Journal of Agricultural Research*, **42**, 661–700.
- He, L., Asseng, S., Zhao, G., Wu, D., Yang, X., Zhuang, W., Jin, N. & Yu, Q. (2015) Impacts of recent climate warming, cultivar changes, and crop management on winter wheat phenology across the Loess Plateau of China. *Agricultural and Forest Meteorology*, **200**, 135–143.
- Iwaki, K., Haruna, S., Niwa, T. & Kato, K. (2001) Adaptation and ecological differentiation in wheat with special reference to geographical variation of growth habit and Vrn genotype. *Plant Breeding*, **120**, 107–114.

- Kato, K. & Yokoyama, H. (1992) Geographical variation in heading characters among wheat landraces, *Triticum aestivum* L., and its implication for their adaptability. *Theoretical and Applied Genetics*, **84**, 259–265.
- Kouressy, M., Dingkuhn, M., Vaksman, M. & Heinemann, A.B. (2008) Adaptation to diverse semi-arid environments of sorghum genotypes having different plant type and sensitivity to photoperiod. *Agricultural and Forest Meteorology*, **148**, 357–371.
- Kumudini, S., Andrade, F.H., Boote, K.J. *et al.* (2014) Predicting maize phenology: intercomparison of functions for developmental response to temperature. *Agronomy Journal*, **106**, 2087–2097.
- Mahfoozi, S., Limin, A.E. & Fowler, D.B. (2001) Influence of vernalization and photoperiod responses on cold hardiness in winter cereals. *Crop Science*, **41**, 1006–1011.
- Menzel, A., Sparks, T.H., Estrella, N. *et al.* (2006) European phenological response to climate change matches the warming pattern. *Global Change Biology*, **12**, 1969–1976.
- Miralles, D.J. & Slafer, G.A. (1999) Wheat development. *Wheat ecology and physiology of yield determination* (ed. by E.H. Satorre and G.A. Slafer), pp. 13–43. Food Products Press, New York.
- Miralles, D.J., Spinedi, M.V., Abeledo, L.G. & Abelleira, D. (2007) Variability on photoperiod responses in Argentinean wheat cultivars differing in length of crop cycle. *Wheat production in stressed environments* (ed. by H.T. Buck, J.E. Nisi and N. Salomón), pp. 599–609. Springer, Dordrecht.
- Mitchell, T.D. & Jones, P.D. (2005) An improved method of constructing a database of monthly climate observations and associated high-resolution grids. *International Journal of Climatology*, **25**, 693–712.
- Monteith, J.L. & Unsworth, M.H. (1990) *Principles of environmental physics*. Arnold, London.
- Müller, C. & Robertson, R.D. (2014) Projecting future crop productivity for global economic modeling. *Agricultural Economics*, **45**, 37–50.
- Olesen, J.E., Børgesen, C.D., Elsgaard, L., Palosuo, T., Rötter, R.P., Skjelvåg, A.O., Peltonen-Sainio, P., Börjesson, T., Trnka, M., Ewert, F., Siebert, S., Brisson, N., Eitzinger, J., Van Asselt, E.D., Oberforster, M. & Van der Fels-Klerx, H.J. (2012) Changes in time of sowing, flowering and maturity of cereals in Europe under climate change. *Food Additives and Contaminants: Part A*, **29**, 1527–1542.
- Ortiz Ferrara, G., Mosaad, M.G., Mahalakshmi, V. & Rajaram, S. (1998) Photoperiod and vernalisation response of Mediterranean wheats, and implications for adaptation. *Euphytica*, **100**, 377–384.
- PEP725 (2014) Pan European Phenology Database. Available at: <http://www.zamg.ac.at/pep725/> (accessed 22 July 2014).
- Porter, J.R. (1993) AFRCWHEAT2: a model of the growth and development of wheat incorporating responses to water and nitrogen. *European Journal of Agronomy*, **2**, 69–82.
- Porter, J.R. & Semenov, M.A. (2005) Crop responses to climatic variation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **360**, 2021–2035.
- Portmann, F.T., Siebert, S. & Doll, P. (2010) MIRCA2000-Global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and hydrological modeling. *Global Biogeochemical Cycles*, **24**, GB1011.
- Raven, P.H., Evert, R.F. & Eichhorn, S.E. (2005) *Biology of plants*. Freeman, New York.
- Rood, S.B. & Major, D.J. (1980) Responses of early corn inbreds to photoperiod. *Crop Science*, **20**, 679–682.
- Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A.C., Müller, C., Arneth, A., Boote, K.J., Folberth, C., Glotter, M., Khabarov, N., Neumann, K., Piontek, F., Pugh, T.A.M., Schmid, E., Stehfest, E., Yang, H. & Jones, J.W. (2014) Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the National Academy of Sciences USA*, **111**, 3268–3273.
- Sacks, W.J., Deryng, D., Foley, J.A. & Ramankutty, N. (2010) Crop planting dates: an analysis of global patterns. *Global Ecology and Biogeography*, **19**, 607–620.
- Slafer, G.A. & Rawson, H.M. (1994) Sensitivity of wheat phasic development to major environmental factors: a re-examination of some assumptions made by physiologists and modellers. *Functional Plant Biology*, **21**, 393–426.
- Summerfield, R.J., Ellis, R.H., Craufurd, P.Q., Aiming, Q., Roberts, E.H. & Wheeler, T.R. (1997) Environmental and genetic regulation of flowering of tropical annual crops. *Euphytica*, **96**, 83–91.
- Tao, F., Zhang, S., Zhang, Z. & Rötter, R.P. (2014) Maize growing duration was prolonged across China in the past three decades under the combined effects of temperature, agronomic management, and cultivar shift. *Global Change Biology*, **20**, 3686–3699.
- Trethowan, R., Morgunov, A., He, Z., De Pauw, R., Crossa, J., Warburton, M., Baytasov, A., Zhang, C., Mergoum, M. & Alvarado, G. (2006) The global adaptation of bread wheat at high latitudes. *Euphytica*, **152**, 303–316.
- Twine, T.E. & Kucharik, C.J. (2009) Climate impacts on net primary productivity trends in natural and managed ecosystems of the central and eastern United States. *Agricultural and Forest Meteorology*, **149**, 2143–2161.
- Waha, K., Van Bussel, L.G.J., Müller, C. & Bondeau, A. (2012) Climate-driven simulation of global crop sowing dates. *Global Ecology and Biogeography*, **21**, 247–259.
- Weir, A.H., Bragg, P.L., Porter, J.R. & Rayner, J.H. (1984) A winter wheat crop stimulation model without water or nutrient limitations. *Journal of Agricultural Science*, **102**, 371–382.
- Wheeler, T.R., Craufurd, P.Q., Ellis, R.H., Porter, J.R. & Vara Prasad, P.V. (2000) Temperature variability and the yield of annual crops. *Agriculture, Ecosystems and Environment*, **82**, 159–167.
- Willmott, C.J. (1982) Some comments on the evaluation of model performance. *Bulletin of the American Meteorological Society*, **63**, 1309–1313.
- Worland, A.J., Appendino, M.L. & Sayers, E.J. (1994) The distribution, in European winter wheats, of genes that influence

ecoclimatic adaptability whilst determining photoperiodic insensitivity and plant height. *Euphytica*, **80**, 219–228.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article at the publisher's web-site.

Appendix S1 Detailed description of the AFRCWHEAT2 model.

Appendix S2 Calculation of the exponentially weighted moving averages of monthly temperatures.

Appendix S3 Detailed description of the computation of the location-specific parameters.

Appendix S4 Locations selected from the PEP725 database.

Appendix S5 Simulated wheat growing period lengths for current (year 2000) and future climatic conditions.

BIOSKETCH

We concentrate in our research on simulation of crop growth at large spatial areas, because it helps us to understand challenges such as which regions have the greatest potential to increase crop production and which regions will be most vulnerable to climate change.

Author contributions: L.v.B., E.S., C.M. and F.E. conceived the idea of computing location-specific parameter values to account for cultivar differences at global scale. All authors were involved in developing the methodology and discussing the model outputs. L.v.B. wrote the model code, ran the models and prepared the manuscript and the supporting material. All authors were involved in reviewing and editing the manuscript.

Editor: Martin Sykes