

The Effects of Intraspecific Genetic Variation on the Dynamic of Predator-Prey Ecological Communities

Sam Fleischer, Pablo Chavarria

May 3, 2015

Preparing Undergraduates through Mentoring for PhDs (PUMP) Research Symposium

Advisor: Dr. Jing Li, Mathematics, CSU Northridge
Consultant: Dr. Casey terHorst, Biology, CSU Northridge
Supported By: Pacific Math Alliance, PUMP, CSU Northridge

Overview

- Motivation
 - Observations in Nature
 - Previous Models
- Our Expansions
 - Gaussian Attack Rate under Coevolution
 - Introduce Stabilizing Selection
 - General Ditrophic Expansion
- Discussion

Observations in Nature

- Predator/Prey interactions are prevalent in nature
 - Crab vs. gastropod [Saloniemi, 1993]
 - Classical Lotka-Volterra model
 - Genetic adaptation is insignificant




Observations in Nature

- Predator/Prey interactions are prevalent in nature
 - Crab vs. gastropod [Saloniemi, 1993]
 - Classical Lotka-Volterra model
 - Genetic adaptation is insignificant
- There is trait variation within species, which causes **variation in fundamental model parameters**
 - Relative strength of crab claw vs. gastropod shell [Saloniemi, 1993]
 - Incorporating trait variation provides **richer dynamics** than classical Lotka-Volterra models



Classical Lotka-Volterra Model



Variables

- $N \equiv$ Prey Density
- $M \equiv$ Predator Density


Parameters

- $\alpha \equiv$ Attack rate
- $r \equiv$ Prey birth rate
- $e \equiv$ Efficiency
- $d \equiv$ Predator death rate

$$\frac{dN}{dt} = N(r - \alpha M)$$

$$\frac{dM}{dt} = M(e\alpha N - d)$$

Classical Lotka-Volterra Model



Variables

- $N \equiv$ Prey Density
- $M \equiv$ Predator Density

Parameters

- $\alpha \equiv$ Attack rate \leftarrow **No variation!**
- $r \equiv$ Prey birth rate
- $e \equiv$ Efficiency
- $d \equiv$ Predator death rate

$$\frac{dN}{dt} = N(r - \alpha M)$$

$$\frac{dM}{dt} = M(e\alpha N - d)$$

Schreiber, Bürger, and Bolnick's Expansion

Assume the **Predator Species** has a normally distributed trait value.

$$p(m, \bar{m}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{(m - \bar{m})^2}{2\sigma^2} \right]$$

Parameters

- $\sigma^2 \equiv$ Predator Trait Variance

Variables

- $m \equiv$ Predator Trait Value

Schreiber, Bürger, and Bolnick's Expansion

Assume the **Predator Species** has a normally distributed trait value.

$$p(m, \bar{m}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{(m - \bar{m})^2}{2\sigma^2} \right]$$

Attack Rate is a Function of the **Predator's Trait Value**

$$a(m) = \alpha \exp \left[-\frac{(m - \theta)^2}{2\tau^2} \right]$$

Parameters

- $\sigma^2 \equiv$ Predator Trait Variance
- $\alpha \equiv$ Maximum attack rate
- $\tau \equiv$ Specialization Constant
- $\theta \equiv$ Optimal trait value

Variables

- $m \equiv$ Predator Trait Value

Schreiber, Bürger, and Bolnick's Expansion

Assume the **Predator Species** has a normally distributed trait value.

$$p(m, \bar{m}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{(m - \bar{m})^2}{2\sigma^2} \right]$$

Attack Rate is a Function of the **Predator's Trait Value**

$$a(m) = \alpha \exp \left[-\frac{(m - \theta)^2}{2\tau^2} \right]$$

Parameters

Variables

- $m \equiv$ Predator Trait Value
- (((No Prey Trait Value)))

- $\sigma^2 \equiv$ Predator Trait Variance
- $\alpha \equiv$ Maximum attack rate
- $\tau \equiv$ Specialization Constant
- $\theta \equiv$ Optimal trait value
↑ **No variation!**

Normally Distributed Trait Values

Assume **Prey** and **Predator** have normally distributed trait values.

$$p(n, \bar{n}) = \frac{1}{\sqrt{2\pi\beta^2}} \exp \left[-\frac{(n - \bar{n})^2}{2\beta^2} \right]$$

$$p(m, \bar{m}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{(m - \bar{m})^2}{2\sigma^2} \right]$$

Variables

- $n \equiv$ Prey Trait Value
- $\bar{n} \equiv$ **Average** Prey Trait Value
- $m \equiv$ Predator Trait Value
- $\bar{m} \equiv$ **Average** Predator Trait Value

Parameters

- $\beta^2 \equiv$ Prey Trait Variance
- $\sigma^2 \equiv$ Predator Trait Variance

Attack Rate

Attack Rate is a Gaussian Function of the **Prey's Trait Value** and the **Predator's Trait Value**

$$a(n, m) = \alpha \exp \left[-\frac{((m - n) - \theta)^2}{2\tau^2} \right]$$

Variables

- $n \equiv$ Prey Trait Value
- $\bar{n} \equiv$ **Average** Prey Trait Value
- $m \equiv$ Predator Trait Value
- $\bar{m} \equiv$ **Average** Predator Trait Value

Parameters

- $\alpha \equiv$ Maximum attack rate
- $\theta \equiv$ **Optimal trait difference**
- $\tau^2 \equiv$ Specialization Constant

Attack Rate

Attack Rate is a Gaussian Function of the **Prey's Trait Value** and the **Predator's Trait Value**

$$a(n, m) = \alpha \exp \left[-\frac{((m - n) - \theta)^2}{2\tau^2} \right]$$

Average Attack Rate

$$\begin{aligned} \bar{a}(\bar{n}, \bar{m}) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} a(n, m) \cdot p(n, \bar{n}) \cdot p(m, \bar{m}) \, dn \, dm \\ &= \frac{\alpha\tau}{\sqrt{\sigma^2 + \beta^2 + \tau^2}} \exp \left[-\frac{((\bar{m} - \bar{n}) - \theta)^2}{2(\sigma^2 + \beta^2 + \tau^2)} \right] \end{aligned}$$

Variables

- $n \equiv$ Prey Trait Value
- $\bar{n} \equiv$ Average Prey Trait Value
- $m \equiv$ Predator Trait Value
- $\bar{m} \equiv$ Average Predator Trait Value

Parameters

- $\alpha \equiv$ Maximum attack rate
- $\theta \equiv$ Optimal trait difference
- $\tau^2 \equiv$ Specialization Constant
- $\beta^2 \equiv$ Prey Trait Variance
- $\sigma^2 \equiv$ Predator Trait Variance

Fitness Assumptions

- Prey experiences **logistic growth** in absence of predator
- Predator experiences **exponential decay** in absence of prey

$$Y(N, n, M, m) = r \left(1 - \frac{N}{K} \right) - Ma(n, m)$$

$$W(N, n, M, m) = eNa(n, m) - d$$

Variables

- $N \equiv$ Prey Density
- $n \equiv$ Prey Trait Value
- $M \equiv$ Predator Density
- $m \equiv$ Predator Trait Value

Parameters

- $r \equiv$ Intrinsic Prey Growth Rate
- $K \equiv$ Prey Carrying Capacity
- $d \equiv$ Predator Death Rate
- $e \equiv$ Efficiency

Average Fitness

$$\begin{aligned}\overline{Y}(N, \bar{n}, M, \bar{m}) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} Y(N, n, M, m) \cdot p(m, \bar{m}) \cdot p(n, \bar{n}) \, dmdn \\ &= r \left(1 - \frac{N}{K} \right) - M\bar{a}(\bar{n}, \bar{m})\end{aligned}$$

$$\begin{aligned}\overline{W}(N, \bar{n}, M, \bar{m}) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} W(N, n, M, m) \cdot p(m, \bar{m}) \cdot p(n, \bar{n}) \, dmdn \\ &= eN\bar{a}(\bar{n}, \bar{m}) - d\end{aligned}$$

Variables

- $N \equiv$ Prey Density
- $\bar{n} \equiv$ **Average** Prey Trait Value
- $M \equiv$ Predator Density
- $\bar{m} \equiv$ **Average** Predator Trait Value


Parameters

- $\bar{r} \equiv$ Intrinsic Prey Growth Rate
- $K \equiv$ Prey Carrying Capacity
- $d \equiv$ Predator Death Rate
- $e \equiv$ Efficiency

Ecological Components

$$\frac{dN}{dt} = N \cdot \overline{Y}(N, \bar{n}, M, \bar{m}) = N \left[r \left(1 - \frac{N}{K} \right) - M \bar{a}(\bar{n}, \bar{m}) \right]$$

$$\frac{dM}{dt} = M \cdot \overline{W}(N, \bar{n}, M, \bar{m}) = M [e N \bar{a}(\bar{n}, \bar{m}) - d]$$



Variables

- $N \equiv$ Prey Density
- $\bar{n} \equiv$ **Average** Prey Trait Value
- $M \equiv$ Predator Density
- $\bar{m} \equiv$ **Average** Predator Trait Value

Parameters

- $\bar{r} \equiv$ Intrinsic Prey Growth Rate
- $K \equiv$ Prey Carrying Capacity
- $d \equiv$ Predator Death Rate
- $e \equiv$ Efficiency

Evolutionary Components

- The evolution of the mean trait value is always in the direction which increases the mean fitness in the population. [Lande, 1976]

$$\frac{d\bar{n}}{dt} = \beta_G^2 \frac{\partial \bar{Y}}{\partial \bar{n}} = \beta_G^2 \frac{M(\theta - (\bar{m} - \bar{n}))}{\sigma^2 + \beta^2 + \tau^2} \bar{a}(\bar{n}, \bar{m})$$

$$\frac{d\bar{m}}{dt} = \sigma_G^2 \frac{\partial \bar{W}}{\partial \bar{m}} = \sigma_G^2 \frac{eN(\theta - (\bar{m} - \bar{n}))}{\sigma^2 + \beta^2 + \tau^2} \bar{a}(\bar{n}, \bar{m})$$

Variables

- $N \equiv$ Prey Density
- $\bar{n} \equiv$ Mean Prey Character
- $M \equiv$ Predator Density
- $\bar{m} \equiv$ Mean Predator Character

Parameters

- $\beta_G^2 \equiv$ Prey genetic variance
- $\sigma_G^2 \equiv$ Predator genetic variance

The Complete 1×1 Model (One Predator Species, One Prey Species)

Ecological Components

$$\frac{dN}{dt} = N \cdot \overline{Y}(N, \bar{n}, M, \bar{m}) = N \left[r \left(1 - \frac{N}{K} \right) - M \bar{a}(\bar{m}, \bar{n}) \right]$$

$$\frac{dM}{dt} = M \cdot \overline{W}(N, \bar{n}, M, \bar{m}) = M [eN \bar{a}(\bar{m}, \bar{n}) - d]$$

Evolutionary Components

$$\frac{d\bar{n}}{dt} = \beta_G^2 \frac{\partial \overline{Y}}{\partial \bar{n}} = \beta_G^2 \frac{M(\theta - (\bar{m} - \bar{n}))}{\sigma^2 + \beta^2 + \tau^2} \bar{a}(\bar{n}, \bar{m})$$

$$\frac{d\bar{m}}{dt} = \sigma_G^2 \frac{\partial \overline{W}}{\partial \bar{m}} = \sigma_G^2 \frac{eN(\theta - (\bar{m} - \bar{n}))}{\sigma^2 + \beta^2 + \tau^2} \bar{a}(\bar{n}, \bar{m})$$

Equilibria - 1×1

Extinction

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = (0, 0, _, _)$$

Equilibria - 1×1

Extinction *Unstable*

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = (0, 0, _, _)$$

Equilibria - 1×1

Extinction *Unstable*

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = (0, 0, _, _)$$

Exclusion

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = (K, 0, \mu^*, \mu^* + \theta)$$

where μ^* is arbitrary

Equilibria - 1×1

Coexistence

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = \left(\frac{d\sqrt{A}}{e\alpha\tau}, \frac{r\sqrt{A}}{\alpha\tau} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right), \mu^*, \mu^* + \theta \right)$$

Equilibria - 1×1



Coexistence

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = \left(\frac{d\sqrt{A}}{e\alpha\tau}, \frac{r\sqrt{A}}{\alpha\tau} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right), \mu^*, \mu^* + \theta \right)$$



Necessary Condition for Asymptotically Stable Coexistence:

$$\frac{\sigma_G^2}{\beta_G^2} > \frac{r}{d} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right)$$



Figures - 1×1 - Stable Exclusion



Figures - 1×1 - Stable Coexistence



Figures - 1×1 - “Arms Race” Coexistence



Avoiding an “Arms Race” with Stabilizing Selection

Assume Prey Growth Rate is a Function of the **Prey's Trait Value**

$$r(n) = \rho \exp \left[-\frac{(n - \phi)^2}{2\gamma^2} \right]$$

Variables

- $n \equiv$ Prey Trait Value
- $\bar{n} \equiv$ **Average Prey Trait Value**

Parameters

- $\rho \equiv$ Maximum Growth Rate
- $\phi \equiv$ **Prey Optimum Trait Value**
- $\gamma^2 \equiv$ Stabilizing Selection Constant

Avoiding an “Arms Race” with Stabilizing Selection

Assume Prey Growth Rate is a Function of the **Prey's Trait Value**

$$r(n) = \rho \exp \left[-\frac{(n - \phi)^2}{2\gamma^2} \right]$$

Average Growth Rate

$$\begin{aligned} \bar{r}(\bar{n}) &= \int_{-\infty}^{\infty} r(n) \cdot p(n, \bar{n}) dn \\ &= \frac{\rho\gamma}{\sqrt{\beta^2 + \gamma^2}} \exp \left[-\frac{(\bar{n} - \phi)^2}{2\gamma^2} \right] \end{aligned}$$

Variables

- $n \equiv$ Prey Trait Value
- $\bar{n} \equiv$ Average Prey Trait Value

Parameters

- $\rho \equiv$ Maximum Growth Rate
- $\phi \equiv$ Prey Optimum Trait Value
- $\gamma^2 \equiv$ Stabilizing Selection Constant
- $\beta^2 \equiv$ Prey Trait Variance

Fitness Assumptions

- Prey experiences **logistic growth** in absence of predator
- Predator experiences **exponential decay** in absence of prey

$$Y(N, n, M, m) = r(n) \left(1 - \frac{N}{K}\right) - Ma(n, m)$$

$$W(N, n, M, m) = eNa(n, m) - d$$

Variables

- $N \equiv$ Prey Density
- $n \equiv$ Prey Trait Value
- $M \equiv$ Predator Density
- $m \equiv$ Predator Trait Value

Parameters

- $r \equiv$ Intrinsic Prey Growth Rate Function
- $K \equiv$ Prey Carrying Capacity
- $d \equiv$ Predator Death Rate
- $e \equiv$ Efficiency

The Complete 1×1 Model (One Predator Species, One Prey Species)

Ecological Components

$$\frac{dN}{dt} = N \cdot \bar{Y}(N, \bar{n}, M, \bar{m}) = N \left[\bar{r}(\bar{n}) \left(1 - \frac{N}{K} \right) - M \bar{a}(\bar{n}, \bar{m}) \right]$$

$$\frac{dM}{dt} = M \cdot \bar{W}(N, \bar{n}, M, \bar{m}) = M [eN \bar{a}(\bar{n}, \bar{m}) - d]$$

Evolutionary Components

$$\frac{d\bar{n}}{dt} = \beta_G^2 \frac{\partial \bar{Y}}{\partial \bar{n}} = \beta_G^2 \left[\bar{r}(\bar{n}) \left(1 - \frac{N}{K} \right) \frac{(\phi - \bar{n})}{\beta^2 + \gamma^2} + \frac{M(\theta - (\bar{m} - \bar{n}))}{\sigma^2 + \beta^2 + \tau^2} \bar{a}(\bar{m}, \bar{n}) \right]$$

$$\frac{d\bar{m}}{dt} = \sigma_G^2 \frac{\partial \bar{W}}{\partial \bar{m}} = \sigma_G^2 \frac{eN(\theta - (\bar{m} - \bar{n}))}{\sigma^2 + \beta^2 + \tau^2} \bar{a}(\bar{n}, \bar{m})$$

Equilibria - 1×1

Extinction

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = (0, 0, _, _)$$

Equilibria - 1×1

Extinction *Unstable*

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = (0, 0, _, _)$$

Equilibria - 1×1

Coexistence

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = \left(\frac{d\sqrt{A}}{e\alpha\tau}, \frac{\rho\gamma\sqrt{A}}{\alpha\tau\sqrt{B}} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right), \phi, \theta + \phi \right)$$

where $A = \sigma^2 + \beta^2 + \tau^2$ and $B = \beta^2 + \gamma^2$

Equilibria - 1×1

Coexistence

$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = \left(\frac{d\sqrt{A}}{e\alpha\tau}, \frac{\rho\gamma\sqrt{A}}{\alpha\tau\sqrt{B}} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right), \phi, \theta + \phi \right)$$

where $A = \sigma^2 + \beta^2 + \tau^2$ and $B = \beta^2 + \gamma^2$

Equilibria - 1×1

Coexistence





$$(N^*, M^*, \bar{n}^*, \bar{m}^*) = \left(\frac{d\sqrt{A}}{e\alpha\tau}, \frac{\rho\gamma\sqrt{A}}{\alpha\tau\sqrt{B}} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right), \phi, \theta + \phi \right)$$

where $A = \sigma^2 + \beta^2 + \tau^2$ and $B = \beta^2 + \gamma^2$





Necessary Condition for Asymptotically Stable Coexistence:

$$\frac{\sigma_G^2}{\beta_G^2} > \frac{\rho\gamma}{d\sqrt{B}} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right) \left(1 - \frac{A}{B} \right)$$





Figures - 1×1 - Stable Exclusion



Figures - 1×1 - Stable Coexistence



Figures - 1×1 - Stable Cycles (Red Queen Dynamics) [Kindrik, Kondrashov, 1994]



Question

What happens when the exclusion **AND**
coexistence stability criterion are **NOT** met?


Question

What happens when the exclusion **AND**
coexistence stability criterion are **NOT** met?

Answer

Stable Limit Cycles

Contour Plot - Coexistence Asymptotic Stability Criterion



$$f_{\text{stable}}(\text{system parameters}) = \frac{\sigma_G^2}{\beta_G^2} - \frac{\rho\gamma}{d\sqrt{B}} \left(1 - \frac{d\sqrt{A}}{Ke\alpha\tau} \right) \left(1 - \frac{A}{B} \right)$$



$f_{\text{stable}} > 0 \implies$ Coexistence is *stable*

$f_{\text{stable}} < 0 \implies$ Coexistence is *unstable*

$f_{\text{stable}} = 0 \implies$ Hopf Bifurcation



$\tau = 0.05$: Limit Cycle vs. Node

$$\frac{\sigma_G}{\beta_G} = 1.3 \Rightarrow f_{\text{stable}} < 0$$



Sam Fleischer, Pablo Chavarria

$$\frac{\sigma_G}{\beta_G} = 1.5 \Rightarrow f_{\text{stable}} > 0$$



The Effects of Intraspecific Genetic Variation on the Dynamic of Predator-Prey

Summary of the 1×1 Model

- 4-dimensional system of ODEs
 - 2 ODEs describing the change in **population size** over time
 - 2 ODEs describing the change in **trait value** over time

Summary of the 1×1 Model

- 4-dimensional system of ODEs
 - 2 ODEs describing the change in **population size** over time
 - 2 ODEs describing the change in **trait value** over time
- 3 equilibrium points
 - Extinction (Unstable)
 - Exclusion (Stable under certain conditions)
 - Coexistence (Stable under certain conditions)

Summary of the 1×1 Model

- 4-dimensional system of ODEs
 - 2 ODEs describing the change in **population size** over time
 - 2 ODEs describing the change in **trait value** over time
- 3 equilibrium points
 - Extinction (Unstable)
 - Exclusion (Stable under certain conditions)
 - Coexistence (Stable under certain conditions)
- Non-Equilibrium Dynamics
 - Constant Growth Rate
 - “Arms Race” Coexistence
 - Gaussian Growth Rate Function
 - Stable Limit Cycles

Expansion of Fitness Functions

Prey Fitness

$$Y(N, n, M, m) = r(n) \left(1 - \frac{N}{K} \right) - Ma(n, m)$$

Predator Fitness

$$W(N, n, M, m) = eNa(n, m) - d$$

Expansion of Fitness Functions

Prey Fitness

$$Y(N, n, M, m) = r(n) \left(1 - \frac{N}{K} \right) - Ma(n, m)$$

↓

$$Y_j(N_j, n_j, [M_i]_{i=1}^u, [m_i]_{i=1}^u) = r_j(n_j) \left(1 - \frac{N_j}{K_j} \right) - \sum_{i=1}^u M_i a_{ij}(n_j, m_i)$$

Predator Fitness

$$W(N, n, M, m) = eNa(n, m) - d$$

Notation: $[x_i]_{i=1}^u = x_1, \dots, x_u$

Expansion of Fitness Functions

Prey Fitness

$$Y(N, n, M, m) = r(n) \left(1 - \frac{N}{K}\right) - Ma(n, m)$$

↓

$$Y_j(N_j, n_j, [M_i]_{i=1}^u, [m_i]_{i=1}^u) = r_j(n_j) \left(1 - \frac{N_j}{K_j}\right) - \sum_{i=1}^u M_i a_{ij}(n_j, m_i)$$

Predator Fitness

$$W(N, n, M, m) = eNa(n, m) - d$$

↓

$$W_i([N_j]_{j=1}^v, [n_j]_{j=1}^v, M_i, m_i) = \sum_{j=1}^v \left[e_{ij} N_j a_{ij}(n_j, m_i) \right] - d_i$$

Notation: $[x_i]_{i=1}^u = x_1, \dots, x_u$

Average Fitness Calculation

$$\begin{aligned}
 \overline{Y}_j(N_j, \overline{n}_j, [M_i]_{i=1}^u, [\overline{m}_i]_{i=1}^u) \\
 &= \int_{\mathbb{R}^{u+1}} Y_j \cdot \prod_{i=1}^u \left[p_i(m_i, \overline{m}_i) \right] \cdot p(n, \overline{n}) \prod_{i=1}^u [dm_i] dn_j \\
 &= \overline{r}_j(\overline{n}_j) \left(1 - \frac{N_j}{K_j} \right) - \sum_{i=1}^u M_i \overline{a}_{ij}(\overline{n}_j, \overline{m}_i)
 \end{aligned}$$

$$\begin{aligned}
 \overline{W}_i(N_j, \overline{n}_j, [M_i]_{i=1}^u, [\overline{m}_i]_{i=1}^u) \\
 &= \int_{\mathbb{R}^{u+1}} W_i \cdot p_i(m_i, \overline{m}_i) \cdot \prod_{j=1}^v \left[p(n_j, \overline{n}_j) \right] dm_i \prod_{j=1}^v [dn_j] \\
 &= \sum_{j=1}^v \left[e_{ij} N_j \overline{a}_{ij}(\overline{n}_j, \overline{m}_i) \right] - d_i
 \end{aligned}$$

The Complete $u \times v$ Model - (u Predator Species, v Prey Species)

Ecological Components

$$\frac{dN_j}{dt} = N_j \overline{Y}_j = N_j \left[\overline{r}_j(\overline{n}_j) \left(1 - \frac{N_j}{K_j} \right) - \sum_{i=1}^u M_i \overline{a}_{ij}(\overline{n}_j, \overline{m}_i) \right]$$


$$\frac{dM_i}{dt} = M_i \overline{W}_i = M_i \left[\sum_{j=1}^v \left[e_{ij} N_j \overline{a}_{ij}(\overline{m}_i, \overline{n}_j) \right] - d_i \right]$$

Evolutionary Components



$$\begin{aligned} \frac{d\overline{n}_j}{dt} = \beta_{Gj}^2 \frac{\partial \overline{Y}_j}{\partial \overline{n}_j} = \beta_{Gj}^2 \left[\overline{r}_j(\overline{n}_j) \left(1 - \frac{N_j}{K_j} \right) \frac{(\phi_j - \overline{n}_j)}{\beta_j^2 + \gamma_j^2} \right. \\ \left. + \sum_{i=1}^u \left[\frac{M_i(\theta_{ij} - (\overline{m}_i - \overline{n}_j))}{\sigma_i^2 + \beta_j^2 + \tau_{ij}^2} \overline{a}_{ij}(\overline{m}_i, \overline{n}_j) \right] \right] \end{aligned}$$

$$\frac{d\overline{m}_i}{dt} = \sigma_{Gi}^2 \sum_{j=1}^v \left[\frac{e_{ij} N_j(\theta_{ij} - (\overline{m}_i - \overline{n}_j))}{\sigma_i^2 + \beta_j^2 + \tau_{ij}^2} \overline{a}_{ij}(\overline{m}_i, \overline{n}_j) \right]$$



The Complete $u \times v$ Model - (u Predator Species, v Prey Species)





Figures - 1×1 - Stable Exclusion



Figures - 1×1 - Stable Coexistence



Figures - 1×1 - “Arms Race” Coexistence



Future Work

- (1×2) Two predator species in competition for one prey
- (2×1) Two prey species in apparent competition via one generalist predator
- (2×2) One specialist predator competing with one generalist predator for two prey
- (2×3) Two specialist predators competing with one generalist predator for two prey species
- $(u \times v)$ The General Ditrophic Expansion
- Intraguild Predation and General Multitrophic Expansion

Thank You!

- PUMP (Preparing Undergraduates through Mentoring towards PhDs)
- The Pacific Math Alliance
- The National Math Alliance
- Dr. Helena Noronha, Dr. Ramin Vakilian, and all other PUMP organizers
- National Science Foundation
- California State University, Northridge
- Dr. Jing Li and Dr. Casey terHorst

Questions?