

We assume the reader is familiar with additive categories and (closed, symmetric) monoidal categories.

Definition 0.1. A *triangulated category* $(\mathcal{C}, \Sigma, \mathcal{D})$ is the data of:

- (1) An additive category \mathcal{C} .
- (2) An additive auto-equivalence $\Sigma : \mathcal{C} \rightarrow \mathcal{C}$ called the *shift functor*.
- (3) A collection \mathcal{D} of *distinguished triangles* in \mathcal{C} , where a is a sequence of arrows of the form

$$X \rightarrow Y \rightarrow Z \rightarrow \Sigma X.$$

Distinguished triangles are also sometimes called *cofiber sequences* or *fiber sequences*.

These data must satisfy the following axioms:

TR0 Given a commutative diagram

$$\begin{array}{ccccccc} X & \longrightarrow & Y & \longrightarrow & Z & \longrightarrow & \Sigma X \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow & & \cong \downarrow \\ X' & \longrightarrow & Y' & \longrightarrow & Z' & \longrightarrow & \Sigma X' \end{array}$$

where the vertical arrows are isomorphisms, if the top row is distinguished then so is the bottom.

TR1 For any object X in \mathcal{C} , the diagram

$$X \xrightarrow{\text{id}_X} X \rightarrow 0 \rightarrow \Sigma X$$

is a distinguished triangle.

TR2 For all $f : X \rightarrow Y$ there exists an object C_f (also sometimes denoted Y/X) called the *cofiber of f* and a distinguished triangle

$$X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X.$$

TR3 Given a solid diagram

$$\begin{array}{ccccccc} X & \longrightarrow & Y & \longrightarrow & Z & \longrightarrow & \Sigma X \\ f \downarrow & & \downarrow & & \vdots & & \downarrow \Sigma f \\ X' & \longrightarrow & Y' & \longrightarrow & Z' & \longrightarrow & \Sigma X' \end{array}$$

such that the leftmost square commutes and both rows are distinguished, there exists a dashed arrow $Z \rightarrow Z'$ which makes the remaining two squares commute.

TR4 A triangle

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{\Sigma} X$$

is distinguished if and only if

$$Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X \xrightarrow{-\Sigma f} \Sigma Y$$

is distinguished.

TR5 (Octahedral axiom) Given three distinguished triangles

$$X \xrightarrow{f} Y \xrightarrow{h} Y/X \rightarrow \Sigma X$$

$$Y \xrightarrow{g} Z \xrightarrow{k} Z/Y \rightarrow \Sigma Y$$

$$X \xrightarrow{g \circ f} Z \xrightarrow{l} Z/X \rightarrow \Sigma X$$

there exists a distinguished triangle

$$Y/X \xrightarrow{u} Z/X \xrightarrow{v} Z/Y \xrightarrow{w} \Sigma(Y/X)$$

such that the following diagram commutes

$$\begin{array}{ccccccc}
 X & \xrightarrow{g \circ f} & Z & \xrightarrow{k} & Z/Y & \xrightarrow{w} & \Sigma(Y/X) \\
 & \searrow f & \nearrow g & \searrow l & \nearrow v & \searrow & \nearrow \Sigma h \\
 & Y & & Z/X & & \Sigma Y & \\
 & \searrow h & \nearrow u & \searrow & \nearrow & \searrow \Sigma f & \\
 & Y/X & \xrightarrow{\quad} & \Sigma X & & &
 \end{array}$$

It turns out that the above definition is actually redundant; TR3 and TR4 follow from the remaining axioms (see Lemmas 2.2 and 2.4 in [1]). From now on, we fix a triangulated category $(\mathcal{C}, \Sigma, \mathcal{D})$. To start, recall the following definition:

Definition 0.2. A sequence

$$X_1 \rightarrow X_2 \rightarrow \cdots \rightarrow X_n$$

of arrows in \mathcal{C} is *exact* if, for any object A in \mathcal{C} , the induced sequence

$$[A, X_1] \rightarrow [A, X_2] \rightarrow \cdots \rightarrow [A, X_n]$$

is an exact sequence of abelian groups.

Proposition 0.3. Any distinguished triangle is an exact sequence (in the sense of Definition 0.2).

Proof. Suppose we have some distinguished triangle

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X.$$

Then we would like to show that given any object A in \mathcal{C} , the sequence

$$[A, X] \xrightarrow{f_*} [A, Y] \xrightarrow{g_*} [A, Z] \xrightarrow{h_*} [A, \Sigma X]$$

is exact. First we show exactness at $[A, Y]$. To see $\text{im } f_* \subseteq \ker g_*$, note it suffices to show that $g \circ f = 0$. Indeed, consider the commuting diagram

$$\begin{array}{ccccccc}
 X & \xlongequal{\quad} & X & \longrightarrow & 0 & \longrightarrow & \Sigma X \\
 \parallel & & \downarrow f & & & & \\
 X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X
 \end{array}$$

The top row is distinguished by axiom TR1. Thus by TR3, the following diagram commutes:

$$\begin{array}{ccccccc}
 X & \xlongequal{\quad} & X & \longrightarrow & 0 & \longrightarrow & \Sigma X \\
 \parallel & & \downarrow f & & \downarrow & & \parallel \\
 X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X
 \end{array}$$

In particular, commutativity of the second square tells us that $g \circ f = 0$, as desired. Conversely, we'd like to show that $\ker g_* \subseteq \text{im } f_*$. Let $\psi : A \rightarrow Y$ be in the kernel of g_* , so that $g \circ \psi = 0$. Consider the following commutative diagram:

$$\begin{array}{ccccccc}
 A & \longrightarrow & 0 & \longrightarrow & \Sigma A & \xrightarrow{-\Sigma \text{id}_A} & \Sigma A \\
 \psi \downarrow & & \downarrow & & & & \\
 Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X & \xrightarrow{-\Sigma f} & \Sigma Y
 \end{array}$$

The top row is distinguished by axioms TR1 and TR4. The bottom row is distinguished by axiom TR4. Thus by axiom TR3 there exists a map $\tilde{\phi} : \Sigma A \rightarrow \Sigma X$ such that the following diagram commutes:

$$\begin{array}{ccccccc} A & \longrightarrow & 0 & \longrightarrow & \Sigma A & \xrightarrow{-\Sigma \text{id}_A} & \Sigma A \\ \psi \downarrow & & \downarrow & & \tilde{\phi} \downarrow & & \Sigma \psi \downarrow \\ Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X & \xrightarrow{-\Sigma f} & \Sigma Y \end{array}$$

Now, since Σ is an equivalence, it is a full functor, so that in particular there exists some $\phi : A \rightarrow X$ such that $\tilde{\phi} = \Sigma \phi$. Then by faithfulness, we may pull back the right square to get a commuting diagram

$$\begin{array}{ccc} A & \xrightarrow{-\text{id}_A} & A \\ \phi \downarrow & & \downarrow \psi \\ X & \xrightarrow{-f} & Y \end{array}$$

Hence,

$$f_*(\phi) = f \circ \phi \stackrel{(*)}{=} -((-f) \circ \phi) = -(\psi \circ (-\text{id}_A)) \stackrel{(*)}{=} \psi \circ \text{id}_A = \psi,$$

where the equalities marked $(*)$ follow by bilinearity of composition in an additive category. Thus $\psi \in \text{im } f_*$, as desired, meaning $\ker g_* \subseteq \text{im } f_*$.

Now, we have shown that

$$[A, X] \xrightarrow{f_*} [A, Y] \xrightarrow{g_*} [A, Z] \xrightarrow{h_*} [A, \Sigma X]$$

is exact at $[A, Y]$. It remains to show exactness at $[A, Z]$. Yet this follows by the exact same argument given above applied to the sequence obtained from the shifted triangle (TR4)

$$Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X \xrightarrow{-\Sigma f} \Sigma Y. \quad \square$$

Lemma 0.4. *Suppose we have a commutative diagram*

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X \\ \downarrow j & & \downarrow k & & \downarrow \ell & & \downarrow \Sigma j \\ X' & \xrightarrow{f'} & Y' & \xrightarrow{g'} & Z' & \xrightarrow{h'} & \Sigma X' \end{array}$$

with both rows distinguished. Then if any two of the maps j , k , and ℓ are isomorphisms, then so is the third.

Proof. Suppose we are given any object A in \mathcal{C} , and consider the commutative diagram

$$\begin{array}{ccccccccccc} [A, X] & \xrightarrow{f_*} & [A, Y] & \xrightarrow{g_*} & [A, Z] & \xrightarrow{k_*} & [A, \Sigma X] & \xrightarrow{-\Sigma f_*} & [A, \Sigma Y] & \xrightarrow{-\Sigma g_*} & [A, \Sigma Z] & \xrightarrow{-\Sigma h_*} & [A, \Sigma^2 X] \\ \downarrow j_* & & \downarrow k_* & & \downarrow \ell_* & & \downarrow \Sigma j_* & & \downarrow \Sigma k_* & & \downarrow \Sigma \ell_* & & \downarrow \Sigma^2 j_* \\ [A, X'] & \xrightarrow{f'_*} & [A, Y'] & \xrightarrow{g'_*} & [A, Z'] & \xrightarrow{h'_*} & [A, \Sigma X'] & \xrightarrow{-\Sigma f'_*} & [A, \Sigma Y'] & \xrightarrow{-\Sigma g'_*} & [A, \Sigma Z'] & \xrightarrow{-\Sigma h'_*} & [A, \Sigma^2 X'] \end{array}$$

The rows are exact by **Proposition 0.3** and repeated applications of axiom TR4. It follows by the five lemma that if j and k are isomorphisms, then ℓ_* is an isomorphism. Similarly, if k and ℓ are isomorphisms then Σj_* is an isomorphism. Finally, if ℓ and j are isomorphisms, then Σk_* is an isomorphism. The desired result follows by faithfulness of Σ and the Yoneda embedding. \square

Proposition 0.5. *Given a map $f : X \rightarrow Y$ in a triangulated category $(\mathcal{C}, \Sigma, \Omega, \mathcal{D})$, the cofiber sequence of f is unique up to isomorphism, in the sense that given any two distinguished triangles*

$$X \xrightarrow{f} Y \rightarrow Z \rightarrow \Sigma X \quad \text{and} \quad X \xrightarrow{f} Y \rightarrow Z' \rightarrow \Sigma X,$$

there exists an isomorphism $Z \rightarrow Z'$ which makes the following diagram commute:

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \longrightarrow & Z & \longrightarrow & \Sigma X \\ \parallel & & \parallel & & \downarrow k & & \parallel \\ X & \xrightarrow{f} & Y & \longrightarrow & Z' & \longrightarrow & \Sigma X \end{array}$$

Proof. Suppose we have two distinguished triangles

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X \quad \text{and} \quad X \xrightarrow{f} Y \xrightarrow{g'} Z' \xrightarrow{h'} \Sigma X,$$

and consider the following commutative diagram

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X \\ \parallel & & \parallel & & & & \\ X & \xrightarrow{f} & Y & \xrightarrow{g'} & Z' & \xrightarrow{h'} & \Sigma X \end{array}$$

By axiom TR3, there exists some map $k : Z \rightarrow Z'$ which makes the following diagram commute:

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X \\ \parallel & & \parallel & & \downarrow k & & \parallel \\ X & \xrightarrow{f} & Y & \xrightarrow{g'} & Z' & \xrightarrow{h'} & \Sigma X \end{array}$$

Now, by Lemma 0.4, k is an isomorphism. □

Proposition 0.6. *Given an arrow $f : X \rightarrow Y$ in \mathcal{C} , there exists an object F_f called the fiber of f , and a distinguished triangle*

$$F_f \rightarrow X \xrightarrow{f} Y \rightarrow \Sigma F_f (\cong C_f).$$

Proof. Since Σ is an equivalence, there exists some functor $\Omega : \mathcal{C} \rightarrow \mathcal{C}$ and natural isomorphisms $\eta : \Omega\Sigma \Rightarrow \text{Id}_{\mathcal{C}}$ and $\varepsilon : \text{Id}_{\mathcal{C}} \Rightarrow \Sigma\Omega$. By axiom TR2, we have a distinguished triangle

$$X \xrightarrow{f} Y \xrightarrow{g} C_f \xrightarrow{h} \Sigma X.$$

Now, consider the commutative diagram

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & C_f & \xrightarrow{h} & \Sigma X \\ \parallel & & \parallel & & \downarrow \eta_{C_f} & & \parallel \\ X & \xrightarrow{f} & Y & \xrightarrow{\tilde{g}} & \Sigma\Omega C_f & \xrightarrow{\tilde{h}} & \Sigma X \end{array}$$

where $\tilde{g} = \eta_{C_f} \circ g$, and $\tilde{h} = \eta_{C_f} \circ h$. Since each vertical map is an isomorphism and the top row is distinguished, the bottom row is also distinguished by axiom TR0. Now, since Σ is an equivalence of categories, it is faithful, so that in particular there exists some map $k : \Omega C_f \rightarrow X$ such that $\Sigma k = -\tilde{h} \implies -\Sigma k = \tilde{h}$. Thus, we have a distinguished triangle of the form

$$X \xrightarrow{f} Y \xrightarrow{\tilde{g}} \Sigma\Omega C_f \xrightarrow{-\Sigma k} \Sigma X.$$

Finally, by axiom TR4, we get a distinguished triangle

$$\Omega C_f \xrightarrow{k} X \xrightarrow{f} Y \xrightarrow{\tilde{g}} \Sigma\Omega C_f,$$

so we may define the fiber of f to be ΩC_f . □

For our purposes, we will always be dealing with triangulated categories with a bit of extra structure, in the following sense:

Definition 0.7. An *adjointly triangulated category* $(\mathcal{C}, \Omega, \Sigma, \eta, \varepsilon, \mathcal{D})$ is the data of a triangulated category $(\mathcal{C}, \Sigma, \mathcal{D})$ along with a functor $\Omega : \mathcal{C} \rightarrow \mathcal{C}$ and natural isomorphisms $\eta : \text{Id}_{\mathcal{C}} \Rightarrow \Sigma\Omega$ and $\varepsilon : \Omega\Sigma \Rightarrow \text{Id}_{\mathcal{C}}$ such that $(\Omega, \Sigma, \eta, \varepsilon)$ forms an adjoint equivalence of \mathcal{C} . In other words, η and ε are natural isomorphisms which also are the unit and counit of an adjunction $\Omega \dashv \Sigma$, so they satisfy either of the following “zig-zag identities”:

$$\begin{array}{ccc} \Omega & \xrightarrow{\Omega\eta} & \Omega\Sigma\Omega \\ & \searrow & \downarrow \varepsilon\Omega \\ & & \Omega \end{array} \quad \begin{array}{ccc} \Sigma\Omega\Sigma & \xleftarrow{\eta\Sigma} & \Sigma \\ \Sigma\varepsilon \downarrow & & \nearrow \\ & & \Sigma \end{array}$$

(Satisfying one implies the other is automatically satisfied, see [2, Lemma 3.2]).

From now on, we will assume that \mathcal{C} is an *adjointly triangulated category* with inverse shift Ω , unit $\eta : \text{Id}_{\mathcal{C}} \Rightarrow \Sigma\Omega$, and counit $\varepsilon : \Omega\Sigma \Rightarrow \text{Id}_{\mathcal{C}}$.

Lemma 0.8. *Given a triangle*

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X,$$

it can be shifted to the left to obtain a distinguished triangle

$$\Omega Z \xrightarrow{\tilde{h}} X \xrightarrow{f} Y \xrightarrow{\tilde{\Omega}g} \Sigma\Omega Z,$$

where $\tilde{h} : \Omega Z \rightarrow X$ is the adjoint of $h : Z \rightarrow \Sigma X$ and $\tilde{\Omega}g : Y \rightarrow \Sigma\Omega Z$ is the adjoint of $\Omega g : \Omega Y \rightarrow \Omega Z$.

Proof. Note that unravelling definitions, then \tilde{h} and \tilde{g} are the compositions

$$\tilde{h} : \Omega Z \xrightarrow{\Omega h} \Omega\Sigma X \xrightarrow{\varepsilon_X} X \quad \text{and} \quad \tilde{\Omega}g : Y \xrightarrow{\eta_Y} \Sigma\Omega Y \xrightarrow{\Sigma\Omega g} \Sigma\Omega Z.$$

Now consider the following diagram:

$$(1) \quad \begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X \\ \parallel & & \parallel & & \eta_Z \downarrow & & \parallel \\ X & \xrightarrow{f} & Y & \xrightarrow{\tilde{\Omega}g} & \Sigma\Omega Z & \xrightarrow{\Sigma\tilde{h}} & \Sigma X \end{array}$$

The left square commutes by definition. To see that the middle square commutes, expanding definitions, note it is given by the following diagram:

$$\begin{array}{ccc} Y & \xrightarrow{g} & Z \\ \parallel & & \downarrow \eta_Y \\ Y & \xrightarrow{\eta_Y} \Sigma\Omega Y & \xrightarrow{\Sigma\Omega g} \Sigma\Omega Z \end{array}$$

and this commutes by naturality of η . To see that the right square commutes, consider the following diagram:

$$\begin{array}{ccc} Z & \xrightarrow{h} & \Sigma X \\ \eta_Z \downarrow & & \nwarrow \eta_{\Sigma X} \\ \Sigma\Omega Z & \xrightarrow{\Sigma\Omega h} \Sigma\Omega\Sigma X & \xrightarrow{\Sigma\varepsilon_X} \Sigma X \end{array}$$

By functoriality of Σ , the bottom composition is $\Sigma\tilde{h}$. The left region commutes by naturality of η . Commutativity of the right region is precisely one of the the zig-zag identities. Hence, since diagram (1) commutes, the vertical arrows are isomorphisms, and the bottom row is distinguished, we have that the top row is distinguished as well by axiom TR0. Then by axiom TR4, since $(f, \tilde{\Omega}g, \Sigma\tilde{h})$ is distinguished, so is the triangle

$$\Omega Z \xrightarrow{\tilde{h}} X \xrightarrow{f} Y \xrightarrow{\tilde{\Omega}g} \Sigma\Omega Z.$$

□

Lemma 0.9. *Given a distinguished triangle*

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X,$$

for any $n > 0$, the triangle

$$\Omega^n X \xrightarrow{(-1)^n \Omega^n f} \Omega^n Y \xrightarrow{(-1)^n \Omega^n g} \Omega^n Z \xrightarrow{(-1)^n \Omega^n h} \Omega^n \Sigma X \cong \Sigma \Omega^n X,$$

is distinguished, where the final isomorphism is given by the composition

$$\Omega^n \Sigma X = \Omega^{n-1} \Omega \Sigma X \xrightarrow{\Omega^{n-1} \varepsilon_X} \Omega^{n-1} X \xrightarrow{\eta_{\Omega^{n-1} X}} \Sigma \Omega \Omega^{n-1} X = \Sigma \Omega^n X.$$

Proof. We give a proof by induction. First we show the case $n = 1$. Note by [Lemma 0.8](#), we have that given a distinguished triangle

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X,$$

we can shift it to the left to obtain a distinguished triangle

$$\Omega Z \xrightarrow{\widetilde{h}} X \xrightarrow{f} Y \xrightarrow{\widetilde{\Omega g}} \Sigma \Omega Z,$$

where \widetilde{h} is the adjoint of $h : Z \rightarrow \Sigma X$ and $\widetilde{\Omega g}$ is the adjoint of $\Omega g : \Omega Y \rightarrow \Omega Z$. If we apply this shifting operation again, we get the distinguished triangle

$$\Omega Y \xrightarrow{\widetilde{\widetilde{\Omega g}}} \Omega Z \xrightarrow{\widetilde{h}} X \xrightarrow{\widetilde{\Omega f}} \Sigma \Omega Y,$$

where unravelling definitions, $\widetilde{\Omega f}$ is the right adjoint of $\Omega f : \Omega X \rightarrow \Omega Y$ and $\widetilde{\widetilde{\Omega g}}$ is the right adjoint of $\widetilde{\Omega g}$, which itself is the left adjoint of Ωg , so $\widetilde{\widetilde{\Omega g}} = \Omega g$. Hence we have a distinguished triangle

$$\Omega Y \xrightarrow{-\Omega g} \Omega Z \xrightarrow{\widetilde{h}} X \xrightarrow{\widetilde{\Omega f}} \Sigma \Omega Y.$$

We may again shift this triangle again and the above arguments yield the distinguished triangle

$$\Omega X \xrightarrow{-\Omega f} \Omega Y \xrightarrow{-\Omega g} \Omega Z \xrightarrow{\widetilde{\Omega(-\widetilde{h})}} \Sigma \Omega X,$$

where $\widetilde{\Omega(-\widetilde{h})}$ is the right adjoint of $\Omega(-\widetilde{h}) = -\Omega \widetilde{h} : \Omega \Omega Z \rightarrow \Omega X$. Explicitly unravelling definitions, $\Omega(-\widetilde{h}) = -\Omega \widetilde{h}$ is the composition

$$\begin{aligned} [\Omega Z \xrightarrow{\eta_{\Omega Z}} \Sigma \Omega \Omega Z \xrightarrow{\Sigma(-\Omega \widetilde{h})} \Sigma \Omega X] &= -[\Omega Z \xrightarrow{\eta_{\Omega Z}} \Sigma \Omega \Omega Z \xrightarrow{\Sigma \Omega \widetilde{h}} \Sigma \Omega X] \\ &= -[\Omega Z \xrightarrow{\eta_{\Omega Z}} \Sigma \Omega \Omega Z \xrightarrow{\Sigma \Omega \Omega h} \Sigma \Omega \Omega \Sigma X \xrightarrow{\Sigma \Omega \varepsilon_X} \Sigma \Omega X] \\ &= -[\Omega Z \xrightarrow{\Omega h} \Omega \Sigma X \xrightarrow{\varepsilon_X} X \xrightarrow{\eta_X} \Sigma \Omega X], \end{aligned}$$

where the first equality follows by additivity of Σ and additivity of composition, the second follows by further unravelling how \widetilde{h} is defined, and the third follows by naturality of η , which tells us the following diagram commutes:

$$\begin{array}{ccccc} \Omega Z & \xrightarrow{\Omega h} & \Omega \Sigma X & \xrightarrow{\varepsilon_X} & X \\ \downarrow \eta_{\Omega Z} & & \downarrow \eta_{\Omega \Sigma X} & & \downarrow \eta_X \\ \Sigma \Omega \Omega Z & \xrightarrow{\Sigma \Omega \Omega h} & \Sigma \Omega \Omega \Sigma X & \xrightarrow{\Sigma \Omega \varepsilon_X} & \Sigma \Omega X \end{array}$$

Thus indeed we have a distinguished triangle

$$\Omega X \xrightarrow{-\Omega f} \Omega Y \xrightarrow{-\Omega g} \Omega Z \xrightarrow{-\Omega h} \Omega \Sigma X \cong \Sigma \Omega X,$$

where the last isomorphism is $\eta_X \circ \varepsilon_X$, as desired.

Now, we show the inductive step. Suppose we know that given a distinguished triangle

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X,$$

that for some $n > 0$ the triangle

$$\Omega^n X \xrightarrow{(-1)^n \Omega^n f} \Omega^n Y \xrightarrow{(-1)^n \Omega^n g} \Omega^n Z \xrightarrow{(-1)^n h^n} \Sigma \Omega^n X,$$

is distinguished, where $h^n : \Omega^n Z \rightarrow \Sigma \Omega^n X$ is the composition

$$\Omega^n Z \xrightarrow{\Omega^n h} \Omega^n \Sigma X \xrightarrow{\Omega^{n-1} \varepsilon_X} \Omega^{n-1} X \xrightarrow{\eta_{\Omega^{n-1} X}} \Sigma \Omega^n X.$$

Then by applying the $n = 1$ case to this triangle, we get that the following triangle is distinguished

$$\Omega^{n+1} X \xrightarrow{-\Omega((-1)^n \Omega^n f)} \Omega^{n+1} Y \xrightarrow{-\Omega((-1)^n \Omega^n g)} \Omega^{n+1} Z \xrightarrow{-\Omega((-1)^n h^n)} \Omega \Sigma \Omega^n X \cong \Sigma \Omega^{n+1} X,$$

where the final isomorphism is the composition

$$\Omega \Sigma \Omega^n X \xrightarrow{\varepsilon_{\Omega^n X}} \Omega^n X \xrightarrow{\eta_{\Omega^n X}} \Sigma \Omega \Omega^n X = \Sigma \Omega^{n+1} X.$$

We claim that this is precisely the distinguished triangle given in the statement of the lemma for $n + 1$. First of all, note that $-\Omega((-1)^n \Omega^n f) = (-1)^{n+1} \Omega^{n+1} f$, $-\Omega((-1)^n \Omega^n g) = (-1)^{n+1} \Omega^{n+1} g$, and $-\Omega((-1)^n h^n) = (-1)^{n+1} \Omega h^n$ by additivity of Ω , so that the triangle becomes

$$(2) \quad \Omega^{n+1} X \xrightarrow{(-1)^{n+1} \Omega^{n+1} f} \Omega^{n+1} Y \xrightarrow{(-1)^{n+1} \Omega^{n+1} g} \Omega^{n+1} Z \xrightarrow{(-1)^{n+1} \Omega h^n} \Omega \Sigma \Omega^n X \cong \Sigma \Omega^{n+1} X.$$

Thus, in order to prove the desired characterization, it remains to show this diagram commutes:

$$\begin{array}{ccccc} \Omega^{n+1} Z & \xrightarrow{(-1)^{n+1} \Omega h^n} & \Omega \Sigma \Omega^n X & \xrightarrow{\varepsilon_{\Omega^n X}} & \Omega^n X \\ (-1)^{n+1} \Omega^{n+1} h \downarrow & & & & \downarrow \eta_{\Omega^n X} \\ \Omega^{n+1} \Sigma X & \xrightarrow{\Omega^n \varepsilon_X} & \Omega^n X & \xrightarrow{\eta_{\Omega^n X}} & \Sigma \Omega^{n+1} X \end{array}$$

(The top composition is the last two arrows in diagram (2), and the bottom composition is the last two arrows in the diagram in the statement of the lemma). Unravelling how h^n is constructed, by additivity of Ω it further suffices to show the outside of the following diagram commutes:

$$\begin{array}{ccccccc} \Omega^{n+1} Z & \xrightarrow{(-1)^{n+1} \Omega^{n+1} h} & \Omega^{n+1} \Sigma X & \xrightarrow{\Omega^n \varepsilon_X} & \Omega^n X & \xrightarrow{\Omega \eta_{\Omega^{n-1} X}} & \Omega \Sigma \Omega^n X \\ \downarrow (-1)^{n+1} \Omega^{n+1} h & & & & \parallel & & \downarrow \varepsilon_{\Omega^n X} \\ \Omega^{n+1} \Sigma X & \xrightarrow{\Omega^n \varepsilon_X} & \Omega^n X & \xrightarrow{\eta_{\Omega^n X}} & \Sigma \Omega^{n+1} X & & \Omega^n X \\ & & & & & \nearrow & \downarrow \eta_{\Omega^n X} \end{array}$$

The left rectangle and bottom right triangle commute by definition. Finally, commutativity of the top right trapezoid is precisely one of the zig-zag identities applied to $\Omega^{n-1} X$. Hence, we have shown the desired result. \square

Proposition 0.10. *Given a distinguished triangle*

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X,$$

let $\tilde{h} : \Omega Z \rightarrow X$ be the left adjoint of h . Then the following infinite sequence is exact:

$$\begin{array}{ccccccc}
 & & & & \dots & & \\
 & & & & \swarrow & & \\
 \Omega^{n+1}Z & \xleftarrow{(-1)^{n+1}\Omega^n\tilde{h}} & \Omega^n X & \xrightarrow{(-1)^n\Omega^n f} & \Omega^n Y & \xrightarrow{(-1)^n\Omega^n g} & \Omega^n Z \xrightarrow{(-1)^n\Omega^{n-1}\tilde{h}} \Omega^{n-1}X \\
 & & & & \swarrow & & \\
 & & & & \dots & & \\
 \Omega Z & \xleftarrow{-\tilde{h}} & X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \xrightarrow{h} \Sigma X \\
 & & & & \swarrow & & \\
 & & & & \dots & & \\
 \Sigma^{n-1}Z & \xleftarrow{(-1)^{n-1}\Sigma^n h} & \Sigma^n X & \xrightarrow{(-1)^n\Sigma^n f} & \Sigma^n Y & \xrightarrow{(-1)^n\Sigma^n g} & \Sigma^n Z \xrightarrow{(-1)^n\Sigma^n h} \Sigma^{n+1}X \\
 & & & & \swarrow & & \\
 & & & & \dots & &
 \end{array}$$

In particular, it remains exact even if we remove the signs.

Proof. Exactness of

$$X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X \xrightarrow{-\Sigma f} \Sigma Y$$

is **Proposition 0.3** and axiom TR4. By induction using axiom TR4, for $n > 0$ we get that each contiguous composition of three maps below is a distinguished triangle:

$$\Sigma^n X \xrightarrow{(-1)^n\Sigma^n f} \Sigma^n Y \xrightarrow{(-1)^n\Sigma^n g} \Sigma^n Z \xrightarrow{(-1)^n\Sigma^n h} \Sigma^{n+1} X \xrightarrow{(-1)^{n+1}\Sigma^{n+1} f} \Sigma^{n+1} Y,$$

thus the sequence is exact by **Proposition 0.3**. It remains to show exactness of the LES to the left of Y . It suffices to show that the row in the following diagram is exact for all $n > 0$:

$$(3) \quad \begin{array}{ccccccc}
 \Omega^n X & \xrightarrow{(-1)^n\Omega^n f} & \Omega^n Y & \xrightarrow{(-1)^n\Omega^n g} & \Omega^n Z & \xrightarrow{(-1)^n\Omega^{n-1}(\varepsilon_X \circ \Omega h)} & \Omega^{n-1} X \xrightarrow{(-1)^{n-1}\Omega^{n-1} f} \Omega^{n-1} Y \\
 & & & & \searrow & & \nearrow \\
 & & & & \Omega^n \Sigma X & & \Omega^{n-1} \varepsilon_X
 \end{array}$$

First of all, to see exactness at $\Omega^n Y$ and $\Omega^n Z$, consider the following commutative diagram:

$$\begin{array}{ccccccc}
 \Omega^n X & \xrightarrow{(-1)^n\Omega^n f} & \Omega^n Y & \xrightarrow{(-1)^n\Omega^n g} & \Omega^n Z & \xrightarrow{(-1)^n\Omega^{n-1}(\varepsilon_X \circ \Omega h)} & \Omega^{n-1} X \\
 \parallel & & \parallel & & \parallel & \nearrow \scriptstyle (-1)^n\Omega^n h & \nearrow \scriptstyle \Omega^{n-1}\varepsilon_X \\
 & & & & \Omega^n \Sigma X & & \downarrow \scriptstyle \eta_{\Omega^{n-1} X} \\
 \Omega^n X & \xrightarrow{(-1)^n\Omega^n f} & \Omega^n Y & \xrightarrow{(-1)^n\Omega^n g} & \Omega^n Z & \dashrightarrow & \Sigma \Omega^n X
 \end{array}$$

(here the dashed arrow is the morphism which makes the diagram commute). The bottom row is distinguished by **Lemma 0.9**. Then by axiom TR0, the top row is distinguished, and thus exact by **Proposition 0.3**. Thus we have shown exactness of (3) at $\Omega^n Y$ and $\Omega^n Z$. It remains to show exactness at $\Omega^{n-1} X$. In the case $n = 1$, we want to show exactness at X in the following diagram:

$$\begin{array}{ccccc}
 \Omega Z & \xrightarrow{-(\varepsilon_X \circ \Omega h)} & X & \xrightarrow{f} & Y \\
 \searrow \scriptstyle -\Omega h & & \nearrow \scriptstyle \varepsilon_X & & \\
 & \Omega \Sigma X & & &
 \end{array}$$

Unravelling definitions, $\varepsilon_X \circ \Omega h$ is precisely the adjoint $\tilde{h} : \Omega Z \rightarrow X$ of $h : Z \rightarrow \Sigma X$, in which case we have that the row in the above diagram fits into a distinguished triangle by [Lemma 0.8](#), and thus it is exact by [Proposition 0.3](#). To see exactness at $\Omega^{n-1}X$ in diagram (3), note that if we apply [Lemma 0.8](#) to the sequence [Lemma 0.9](#) for $n-1$, then we get that the following composition fits into a distinguished triangle, and is thus exact:

$$\Omega^n Z \xrightarrow{-k} \Omega^{n-1} X \xrightarrow{(-1)^{n-1} \Omega^{n-1} f} \Omega^{n-1} Y,$$

where $k : \Omega(\Omega^{n-1} Z) \rightarrow \Omega^{n-1} X$ is the adjoint of the composition

$$\Omega^{n-1} Z \xrightarrow{(-1)^{n-1} \Omega^{n-1} h} \Omega^{n-1} \Sigma X \xrightarrow{\Omega^{n-2} \varepsilon_X} \Omega^{n-2} X \xrightarrow{\eta_{\Omega^{n-2} X}} \Sigma \Omega^{n-1} X.$$

Further expanding how adjoints are constructed, k is the composition

$$\Omega^n Z \xrightarrow{(-1)^{n-1} \Omega^n h} \Omega^n \Sigma X \xrightarrow{\Omega^{n-1} \varepsilon_X} \Omega^{n-1} X \xrightarrow{\Omega \eta_{\Omega^{n-2} X}} \Omega \Sigma \Omega^{n-1} X \xrightarrow{\varepsilon_{\Omega^{n-1} X}} \Omega^{n-1} X.$$

Thus, in order to show exactness of (3) at $\Sigma^{n-1} X$, it suffices to show that $k = (-1)^{n-1} \Omega^{n-1} (\varepsilon_X \circ \Omega h)$. To that end, consider the following diagram:

$$\begin{array}{ccccc} \Omega^n Z & \xrightarrow{(-1)^{n-1} \Omega^n h} & \Omega^n \Sigma X & \xrightarrow{\Omega^{n-1} \varepsilon_X} & \Omega^{n-1} X & \xrightarrow{\Omega \eta_{\Omega^{n-2} X}} & \Omega \Sigma \Omega^{n-1} X \\ \downarrow (-1)^{n-1} \Omega^n h & & & & \searrow & & \downarrow \varepsilon_{\Omega^{n-1} X} \\ \Omega^n \Sigma X & \xrightarrow{\hspace{10em}} & \Omega^{n-1} X & & & & \end{array}$$

$\Omega^{n-1} \varepsilon_X$

The top composition is k , while the bottom composition is $(-1)^{n-1} \Omega^{n-1} (\varepsilon_X \circ \Omega h)$. The left region commutes by definition, while commutativity of the right region is precisely one of the zig-zag identities applied to $\Omega^{n-2} X$. Thus, we have shown that $-k = (-1)^n \Omega^{n-1} (\varepsilon_X \circ \Omega h)$, so (3) is exact at $\Omega^{n-1} X$, as desired. \square

Also important for our work is the concept of a *tensor triangulated category*, that is, a triangulated symmetric monoidal category in which the triangulated structures are compatible, in the following sense:

Definition 0.11. A *tensor triangulated category* is a triangulated symmetric monoidal category $(\mathcal{C}, \otimes, S, \Sigma, \mathcal{D})$ such that:

TT1 For all objects X and Y in \mathcal{C} , there are natural isomorphisms

$$e_{X,Y} : \Sigma X \otimes Y \xrightarrow{\cong} \Sigma(X \otimes Y).$$

TT2 For each object X in \mathcal{C} , the functor $X \otimes (-) \cong (-) \otimes X$ is an additive functor.

TT3 For each object X in \mathcal{C} , the functor $X \otimes (-) \cong (-) \otimes X$ preserves distinguished triangles, in that given a distinguished triangle/(co)fiber sequence

$$A \xrightarrow{f} B \xrightarrow{g} C \xrightarrow{h} \Sigma A,$$

then also

$$X \otimes A \xrightarrow{X \otimes f} X \otimes B \xrightarrow{X \otimes g} X \otimes C \xrightarrow{X \otimes h} \Sigma(X \otimes A)$$

and

$$A \otimes X \xrightarrow{f \otimes X} B \otimes X \xrightarrow{g \otimes X} C \otimes X \xrightarrow{h \otimes X} \Sigma(A \otimes X)$$

are distinguished triangles, where here we are being abusive and writing $X \otimes h$ and $h \otimes X$ to denote the compositions

$$X \otimes C \xrightarrow{X \otimes h} X \otimes \Sigma A \xrightarrow{\tau} \Sigma A \otimes X \xrightarrow{e_{A,X}} \Sigma(A \otimes X) \xrightarrow{\Sigma \tau} \Sigma(X \otimes A)$$

and

$$C \otimes X \xrightarrow{h \otimes X} \Sigma A \otimes X \xrightarrow{e_{A,X}} \Sigma(A \otimes X),$$

respectively.

TT4 Given objects X , Y , and Z in \mathcal{C} , the following diagram must commute:

$$\begin{array}{ccc} (\Sigma X \otimes Y) \otimes Z & \xrightarrow{e_{X,Y} \otimes Z} & \Sigma(X \otimes Y) \otimes Z \xrightarrow{e_{X \otimes Y, Z}} \Sigma((X \otimes Y) \otimes Z) \\ \alpha \downarrow & & \downarrow \Sigma \alpha \\ \Sigma X \otimes (Y \otimes Z) & \xrightarrow{e_{X,Y \otimes Z}} & \Sigma(X \otimes (Y \otimes Z)) \end{array}$$

TT5 The following diagram must commute

$$\begin{array}{ccccc} \Sigma S \otimes \Sigma S & \xrightarrow{e_{S,S}} & \Sigma(S \otimes \Sigma S) & \xrightarrow{\Sigma \lambda_{\Sigma S}} & \Sigma^2 S \\ \tau \downarrow & & & & \downarrow -1 \\ \Sigma S \otimes \Sigma S & \xrightarrow{e_{S,S}} & \Sigma(S \otimes \Sigma S) & \xrightarrow{\Sigma \lambda_{\Sigma S}} & \Sigma^2 S \end{array}$$

In other words, $\tau_{\Sigma S, \Sigma S} = -1$.

Usually, most tensor triangulated categories that arise in nature will satisfy additional coherence axioms (see axioms TC1–TC5 in [1]), but the above definition will suffice for our purposes. In what follows, we fix a tensor triangulated category $(\mathcal{C}, \otimes, S, \Sigma, e, \mathcal{D})$.

Definition 0.12. There are natural isomorphisms

$$e'_{X,Y} : X \otimes \Sigma Y \xrightarrow{\cong} \Sigma(X \otimes Y)$$

obtained via the composition

$$X \otimes \Sigma Y \xrightarrow{\tau} \Sigma Y \otimes X \xrightarrow{e_{Y,X}} \Sigma(Y \otimes X) \xrightarrow{\Sigma \tau} \Sigma(X \otimes Y).$$

Lemma 0.13. For all X and Y in \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc} \Sigma X \otimes \Sigma Y & \xrightarrow{-e_{X,\Sigma Y}} & \Sigma(X \otimes \Sigma Y) \\ e'_{\Sigma X, Y} \downarrow & & \downarrow \Sigma e'_{X,Y} \\ \Sigma(\Sigma X \otimes Y) & \xrightarrow{\Sigma e_{X,Y}} & \Sigma \Sigma(X \otimes Y) \end{array}$$

(note the sign on the top map).

Proof. Note there are natural isomorphisms

$$a_X : \Sigma X \xrightarrow{\cong} \Sigma S \otimes X$$

given by the composition

$$\Sigma X \xrightarrow{\Sigma \lambda_X} \Sigma(S \otimes X) \xrightarrow{e_{S,X}^{-1}} \Sigma S \otimes X.$$

Furthermore, under the isomorphism $a : \Sigma \cong \Sigma S \otimes -$, $e_{X,Y} : \Sigma X \otimes Y \cong \Sigma(X \otimes Y)$ corresponds to the associator $(\Sigma S \otimes X) \otimes Y \cong \Sigma S \otimes (X \otimes Y)$. Indeed, consider the following diagram:

$$\begin{array}{ccc} \Sigma X \otimes Y & \xrightarrow{e_{X,Y}} & \Sigma(X \otimes Y) \\ \Sigma \lambda_{X \otimes Y} \downarrow & \swarrow \Sigma(\lambda_X \otimes Y) & \downarrow \Sigma \lambda_{X \otimes Y} \\ \Sigma(S \otimes X) \otimes Y & \xrightarrow{e_{S \otimes X, Y}} \Sigma((S \otimes X) \otimes Y) & \xrightarrow{\Sigma \alpha} \Sigma(S \otimes (X \otimes Y)) \\ e_{S,X}^{-1} \otimes Y \downarrow & & \downarrow e_{S,X \otimes Y}^{-1} \\ (\Sigma S \otimes X) \otimes Y & \xrightarrow{\alpha} & \Sigma S \otimes (X \otimes Y) \end{array}$$

The two vertical composites are $a_X \otimes Y$ and $a_{X \otimes Y}$, respectively. The top trapezoid commutes by naturality of e . The triangle commutes by coherence for a monoidal category. Finally, commutativity of the bottom rectangle is axiom TT4 for a tensor triangulated category.

Similarly, under the isomorphism $a : \Sigma \cong \Sigma S \otimes -$, $e'_{X,Y} : X \otimes \Sigma Y \cong \Sigma(X \otimes Y)$ corresponds to the map $X \otimes \Sigma S \otimes Y \xrightarrow{\tau_{X,\Sigma S \otimes Y}} X \otimes Y \otimes \Sigma S$. To see this, consider the following diagram:

$$\begin{array}{ccccccc}
 X \otimes \Sigma Y & \xrightarrow{\tau_{X,\Sigma Y}} & \Sigma Y \otimes X & \xrightarrow{e_{Y,X}} & \Sigma(Y \otimes X) & \xrightarrow{\Sigma \tau_{Y,X}} & \Sigma(X \otimes Y) \\
 X \otimes \Sigma \lambda_Y \downarrow & & \Sigma \lambda_Y \otimes X \downarrow & & \Sigma(\lambda_Y \otimes X) \downarrow & & \Sigma \lambda_{X \otimes Y} \downarrow \\
 X \otimes \Sigma(S \otimes Y) & \xrightarrow{\tau_{X,\Sigma(S \otimes Y)}} & \Sigma(S \otimes Y) \otimes X & \xrightarrow{e_{S \otimes Y, X}} & \Sigma(S \otimes Y \otimes X) & \xrightarrow{\Sigma(S \otimes \tau_{Y,X})} & \Sigma(S \otimes X \otimes Y) \\
 X \otimes e_{S,Y}^{-1} \downarrow & & e_{S,Y} \otimes X \uparrow & & e_{S,Y \otimes X} \nearrow & & e_{S,X \otimes Y}^{-1} \downarrow \\
 & & \Sigma S \otimes Y \otimes X & & & & \\
 X \otimes \Sigma S \otimes Y & \xrightarrow{\tau_{X,\Sigma S \otimes Y}} & & \xrightarrow{\tau_{X,\Sigma S \otimes Y}} & \Sigma S \otimes X \otimes Y & & \\
 & & \nearrow \tau_{X,\Sigma S \otimes Y} & & \searrow \Sigma S \otimes \tau_{Y,X} & &
 \end{array}$$

Here we are taking the associators to be isomorphisms, by coherence for monoidal categories. The top horizontal composition is $e'_{X,Y}$, by definition. The vertical edge compositions are $X \otimes a_Y$ and $a_{X \otimes Y}$. The top left rectangle commutes by naturality of τ . The top middle rectangle commutes by naturality of e . The top right triangle commutes by naturality of τ . The bottom left trapezoid commutes by naturality of τ . The small middle triangle commutes by axiom TT4 for a tensor triangulated category. The bottom triangle commutes by coherence for a symmetric monoidal category. Finally, the remaining region on the right commutes by naturality of e . Thus, in order to show the diagram in the statement of the lemma commutes, it suffices to show the following diagram commutes:

$$\begin{array}{ccc}
 \Sigma S \otimes X \otimes \Sigma S \otimes Y & \xrightarrow{-\alpha} & \Sigma S \otimes X \otimes \Sigma S \otimes Y \\
 \tau_{\Sigma S \otimes X, \Sigma S \otimes Y} \downarrow & & \downarrow \Sigma S \otimes \tau_{X, \Sigma S \otimes Y} \\
 \Sigma S \otimes \Sigma S \otimes X \otimes Y & \xrightarrow{\alpha} & \Sigma S \otimes \Sigma S \otimes X \otimes Y
 \end{array}$$

To see this diagram commutes, consider the following diagram:

$$\begin{array}{ccc}
 \Sigma S \otimes X \otimes \Sigma S \otimes Y & \xrightarrow{\alpha} & \Sigma S \otimes X \otimes \Sigma S \otimes Y \xrightarrow{\Sigma S \otimes \tau_{X, \Sigma S \otimes Y}} \Sigma S \otimes \Sigma S \otimes X \otimes Y \\
 \tau_{\Sigma S \otimes X, \Sigma S \otimes Y} \downarrow & & \downarrow \tau_{\Sigma S, \Sigma S \otimes X \otimes Y} \\
 \Sigma S \otimes \Sigma S \otimes X \otimes Y & \xrightarrow{\alpha} & \Sigma S \otimes \Sigma S \otimes X \otimes Y
 \end{array}$$

The diagram commutes by coherence for a symmetric monoidal category. The desired result follows by applying axiom TT5 for a tensor triangulated category, which tells us that $\tau_{\Sigma S, \Sigma S} = -1$, and additivity of $- \otimes -$ and composition. \square

Lemma 0.14. *Let $A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} D$ be any sequence isomorphic to a distinguished triangle. Then given any E in \mathcal{C} , the sequence*

$$E \otimes A \xrightarrow{E \otimes a} E \otimes B \xrightarrow{E \otimes b} E \otimes C \xrightarrow{E \otimes c} E \otimes D$$

is exact.

Proof. Suppose we have a commutative diagram

$$\begin{array}{ccccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z & \xrightarrow{h} & \Sigma X \\ \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow & & \delta \downarrow \\ A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & D \end{array}$$

where the top row is distinguished and the vertical arrows are isomorphisms. Then the following diagram commutes by functoriality of $- \otimes -$:

$$\begin{array}{ccccccc} E \otimes X & \xrightarrow{E \otimes f} & E \otimes Y & \xrightarrow{E \otimes g} & E \otimes Z & \xrightarrow{E \otimes' h} & \Sigma(E \otimes X) \\ \downarrow E \otimes \alpha & & \downarrow E \otimes \beta & & \downarrow E \otimes \gamma & \nearrow E \otimes h & \nwarrow E \otimes \delta \\ & & & & & E \otimes \Sigma X & \\ & & & & & \nwarrow E \otimes \delta & \nearrow (E \otimes \delta) \circ e'_{E,X} \\ E \otimes A & \xrightarrow{E \otimes a} & E \otimes B & \xrightarrow{E \otimes b} & E \otimes C & \xrightarrow{E \otimes c} & E \otimes D \end{array}$$

The top triangle is distinguished by axiom TT3 for a tensor triangulated category, thus exact by [Proposition 0.3](#), so that the bottom triangle is also exact since the vertical arrows are isomorphisms and each square commutes. \square

Definition 0.15. An *adjointly tensor triangulated* category is a tensor triangulated category $(\mathcal{C}, \otimes, S, \Sigma, e, \mathcal{D})$ along with the structure of an adjointly triangulated category $(\mathcal{C}, \Omega, \Sigma, \eta, \varepsilon, \mathcal{D})$.

From now on, we fix an adjointly tensor triangulated category $(\mathcal{C}, \otimes, S, \Omega, \Sigma, \eta, \varepsilon, e, \mathcal{D})$. First of all, note we may define natural isomorphisms $o_{X,Y} : \Omega X \otimes Y \xrightarrow{\cong} \Omega(X \otimes Y)$ and $o'_{X,Y} : X \otimes \Omega Y \xrightarrow{\cong} \Omega(X \otimes Y)$ as the compositions

$$o_{X,Y} : \Omega X \otimes Y \xrightarrow{\varepsilon_{\Omega X \otimes Y}^{-1}} \Omega \Sigma(\Omega X \otimes Y) \xrightarrow{\Omega e_{\Omega X, Y}^{-1}} \Omega(\Sigma \Omega X \otimes Y) \xrightarrow{\Omega(\eta_X^{-1} \otimes Y)} \Omega(X \otimes Y)$$

and

$$o'_{X,Y} : X \otimes \Omega Y \xrightarrow{\tau_{X, \Omega Y}} \Omega Y \otimes X \xrightarrow{o_{Y, X}} \Omega(Y \otimes X) \xrightarrow{\Omega \tau_{Y, X}} \Omega(X \otimes Y).$$

These are both clearly natural by naturality of ε , e , η , and τ .

Proposition 0.16. *Suppose we have a distinguished triangle*

$$X \rightarrow Y \rightarrow Z \rightarrow \Sigma X$$

in \mathcal{C} . Then given any object E in \mathcal{C} , the long exact sequence given in [Proposition 0.10](#) remains exact after applying $E \otimes -$.

Proof. Recall that in the proof of [Proposition 0.10](#) we showed that the sequence was exact by showing that any two consecutive maps were isomorphic to a part of a distinguished triangle. Then the desired result follows from [Lemma 0.14](#). \square