

In what follows, we fix an abelian group A . We will freely use the theory and results of ??

Definition 0.1. An A -graded spectral sequence $(E_r, d_r)_{r \geq r_0}$ is the data of:

- A collection of A -graded abelian groups $\{E_r^*\}_{r \geq r_0}$
- A collection of A -graded homomorphisms $d_r : E_r \rightarrow E_r$ for $r \geq r_0$ (of possibly nonzero degree) such that $d_r \circ d_r = 0$
- For each $r \geq r_0$, an A -graded isomorphism $E_{r+1} \cong \ker d_r / \operatorname{im} d_r$ of degree 0 (where $\ker d_r$ and $\operatorname{im} d_r$ are canonically A -graded by ??, and their quotient is canonically A -graded by ??).

Typically we call a \mathbb{Z}^2 -graded spectral sequence a *bigraded* spectral sequence, and a \mathbb{Z}^3 -graded spectral sequence is a *trigraded* spectral sequence.

For our purposes, we will only care about spectral sequences which arise from A -graded *unrolled exact couples*. In what follows, we follow [1], with minor modifications for our setting, in which everything is A -graded.

Definition 0.2. An A -graded *unrolled exact couple* $(D, E; i, j, k)$ is a diagram of A -graded abelian groups and A -graded homomorphisms (of possibly non-zero degree)

$$\begin{array}{ccccccc} \cdots & \longrightarrow & D^{s+2} & \xrightarrow{i} & D^{s+1} & \xrightarrow{i} & D^s & \xrightarrow{i} & D^{s-1} & \longrightarrow & \cdots \\ & & \downarrow j & \swarrow k & \downarrow j & \swarrow k & \downarrow j & \swarrow k & \downarrow j & & \\ & & E^{s+2} & & E^{s+1} & & E^s & & E^{s-1} & & \end{array}$$

in which each triangle $D^{s+1} \xrightarrow{i} D^s \xrightarrow{j} E^s \xrightarrow{k} D^{s+1}$ is an exact sequence. We require that each occurrence of i (resp. j , k) is of the same degree. In other words, an unrolled exact couple can be described as a tuple $(D, E; i, j, k)$ of $\mathbb{Z} \times A$ -graded abelian groups and $\mathbb{Z} \times A$ -graded maps $i : D \rightarrow D$, $j : D \rightarrow E$, and $k : E \rightarrow D$, such that the \mathbb{Z} -degrees of i , j , and k are -1 , 0 , and 1 , respectively. Usually i and one of j or k will be of A -degree 0.

Given an A -graded unrolled exact couple $(D, E; i, j, k)$, we may define an associated $\mathbb{Z} \times A$ -graded spectral sequence as follows: Given some $s \in \mathbb{Z}$ and some $r \geq 1$, we first define the following subgroups of E_s :

$$Z_r^s := k^{-1}(\operatorname{im}[i^{r-1} : D^{s+r} \rightarrow D^{s+1}]) \quad \text{and} \quad B_r^s := j(\ker[i^{r-1} : D^s \rightarrow D^{s-r+1}])$$

where we adopt the convention that i^0 is simply the identity. These are furthermore A -graded subgroups of E_s (by ?? and ??). In this way, for each $s \in \mathbb{Z}$, we get an infinite sequence of A -graded subgroups:

$$0 = B_1^s \subseteq B_2^s \subseteq B_3^s \subseteq \cdots \subseteq \operatorname{im} j = \ker k \subseteq \cdots \subseteq Z_3^s \subseteq Z_2^s \subseteq Z_1^s = E^s.$$

Now, for each $s \in \mathbb{Z}$ and $r \geq 1$, we define the A -graded abelian group

$$E_r^s := Z_r^s / B_r^s,$$

so that in particular $E_1^s = E^s$ for all $s \in \mathbb{Z}$, as $Z_1^s = k^{-1}(D^{s+1}) = E^s$ and $B_1^s = j(\ker \operatorname{id}_{D^s}) = j(0) = 0$. Now we can define differentials $d_r^s : E_r^s \rightarrow E_r^{s+r}$ to be the composition

$$E_r^s = Z_r^s / B_r^s \xrightarrow{k} \operatorname{im}[i^{r-1} : D^{s+r} \rightarrow D^{s+1}] \xrightarrow{j \circ i^{-(r-1)}} Z_r^{s+r} / B_r^{s+r} = E_r^{s+r},$$

where given some $e \in Z_r^s = k^{-1}(\operatorname{im} i^{r-1})$, the first arrow takes a class $[e] \in E_r^s$ represented by some $e \in Z_r^s$ to the element $k(e)$, which lives in $\operatorname{im} i^{r-1}$ by definition, and the second arrow takes $i^{r-1}(d)$ to the class $[j(d)]$. Note the first map is well-defined, as given $b \in B_r^s = j(\ker[i^{r-1}])$, we

have $k(b) = 0$, as $b \in \text{im } j = \ker k$. To see the second map is well-defined, first note that given $d \in D^{s+r}$, that

$$k(j(d)) = 0 \in \text{im}[i^{r-1} : D^{s+2r} \rightarrow D^{s+r+1}],$$

so that

$$j(d) \in k^{-1}(\text{im}[i^{r-1} : D^{s+2r} \rightarrow D^{s+r+1}]) = Z_r^{s+r},$$

as desired, so that given $d \in D^{s+r}$, $j(d) \in Z_r^{s+r}$, so it makes sense to discuss the class $[j(d)] \in Z_r^{s+r}/B_r^{s+r} = E_r^{s+r}$. Secondly, if $i^{r-1}(d) = i^{r-1}(d')$ for some $d, d' \in D^{s+r}$, then

$$j(d) - j(d') = j(d - d') \in j(\ker[i^{r-1} : D^{s+r} \rightarrow D^{s+1}]) = B_r^{s+r},$$

so that $[j(d)] = [j(d')]$ in E_r^{s+r} , as desired. It is straightforward to check that these maps are also A -graded homomorphisms, so that by unravelling definitions d_r^s is an A -graded homomorphism of degree $\deg k - (r-1) \cdot \deg i + \deg j$ (so that in the standard case $\deg i = 0$, d_r^s simply has degree $\deg k + \deg j$).

These differentials square to zero, in the sense that for each $s \in \mathbb{Z}$ and $r \geq 1$ we have that $d_r^{s+r} \circ d_r^s : E_r^s \rightarrow E_r^{s+2r}$ is the zero map. Indeed, suppose we are given some class $[e] \in E_r^s$ represented by an element $e \in E^s$, so $k(e) = i^{r-1}(d)$ for some $d \in D^{s+r}$. Then

$$d_r^{s+r}(d_r^s([e])) = d_r^{s+r}([j(d)]) = [j(i^{-(r-1)}(k(j(d))))] = [j(i^{-(r-1)}(0))] = 0,$$

where the second-to-last equality follows by the fact that $k \circ j = 0$. Note that by unravelling definitions, $d_1^s = j \circ k$.

We claim that $\ker d_r^s = Z_{r+1}^s/B_r^s$. First of all, let $[e] \in E_r^s = Z_r^s/B_r^s$, so that $[e]$ is represented by some $e \in E^s$ with $k(e) = i^{r-1}(d)$ for some $d \in D^{s+r}$. Then if $[e] \in \ker d_r^s$, by definition this means $j(d) \in B_r^{s+r} = j(\ker[i^{r-1} : D^{s+r} \rightarrow D^{s+1}])$, so $j(d) = j(d')$ for some $d' \in D^{s+r}$ with $i^{r-1}(d') = 0$. Thus $d - d' \in \ker j = \text{im } i$, so there exists some $d'' \in D^{s+r+1}$ such that $i(d'') = d - d'$. Then

$$k(e) = i^{r-1}(d) = i^{r-1}(i(d'') + d') = i^r(d'') + i^{r-1}(d'),$$

but since $i^{r-1}(d') = 0$, we have $k(e) \in \text{im}[i^r : D^{s+r+1} \rightarrow D^{s+1}]$, so that $e \in Z_{r+1}^s$, meaning $[e] \in Z_{r+1}^s/B_r^s$, as desired. On the other hand, suppose we are given some class $[e] \in Z_{r+1}^s/B_r^s$, represented by $e \in Z_{r+1}^s$ with $k(e) \in \text{im}[i^r : D^{s+r+1} \rightarrow D^{s+1}]$. Then if we write $k(e) = i^r(d) = i^{r-1}(i(d))$, then $d_r^s([e]) = [j(i(d))] = 0$ (since $j \circ i = 0$), as asserted.

Finally, we claim that the image of $d_r^{s-r} : E_r^{s-r} \rightarrow E_r^s$ is B_{r+1}^s/B_r^s . First, let $e \in Z_r^{s-r}$, so $k(e) = i^{r-1}(d)$ for some $d \in D^s$. Then we'd like to show that $d_r^s([e]) = [j(d)]$ belongs to B_{r+1}^s/B_r^s . It suffices to show that $d \in \ker[i^r : D^s \rightarrow D^{s-r}]$. To see this, note that

$$i^r(d) = i(i^{r-1}(d)) = i(k(e)) = 0,$$

since $i \circ k = 0$. Hence we've shown $\text{im } d_r^{s-r} \subseteq B_{r+1}^s/B_r^s$. Conversely, let $j(d) \in B_{r+1}^s$, so $d \in D^s$ and $i^r(d) = 0$. Then we'd like to show that $[j(d)] \in B_{r+1}^s/B_r^s$ is in the image of d_r^{s-r} . To see this, note that

$$i^r(d) = 0 \implies i^{r-1}(d) \in \ker i = \text{im } k,$$

so there exists some $e \in E_r^{s-r}$ such that $k(e) = i^{r-1}(d)$, so $e \in Z_r^{s-r}$. Unravelling definitions, it follows that $d_r^{s-r}([e]) = [j(d)]$, so $[j(d)]$ is indeed in the image of d_r^{s-r} , as desired.

To recap, we have constructed for each $s \in \mathbb{Z}$ and $r \geq 1$ an A -graded abelian group E_r^s along with differentials $d_r^s : E_r^s \rightarrow E_r^{s+r}$. Furthermore, if we take homology in the middle term of the following sequence

$$E_r^{s-r} \xrightarrow{d_r^{s-r}} E_r^s \xrightarrow{d_r^s} E_r^{s+r},$$

we get

$$\ker d_r^s / \operatorname{im} d_r^{s-r} = \frac{Z_{r+1}^s / B_r^s}{B_{r+1}^s / B_r^s} \cong Z_{r+1}^s / B_{r+1}^s = E_{r+1}^s.$$

Thus, we get a spectral sequence:

Proposition 0.3. *We may associate a $\mathbb{Z} \times A$ -graded spectral sequence $r \mapsto (E_r, d_r)$ to the A -graded unrolled exact couple $(D, E; i, j, k)$ by defining $E_r := \bigoplus_{s \in \mathbb{Z}} E_r^s$ and the differentials*

$$d_r : E_r \rightarrow E_r$$

are those constructed above, which have $\mathbb{Z} \times A$ -degree $(r, \deg j - (r-1) \cdot \deg i + \deg k)$.

Definition 0.4 (Exact couple). An A -graded *exact couple* is a tuple $\mathcal{E} = (D, E; i, j, k)$, where D and E are A -graded abelian groups and i, j , and k are A -graded homomorphisms (of possibly nonzero degree)

$$\begin{array}{ccc} D_* & \xrightarrow{i} & D_* \\ & \swarrow k & \searrow j \\ & E_* & \end{array}$$

which form an *exact triangle*, in the sense that kernel = image at each vertex.

Definition 0.5 (Derived couple). Given an exact couple $(D, E; i, j, k)$ as in the above definition, the composition $j \circ k : E \rightarrow E$ itself satisfies

$$(j \circ k) \circ (j \circ k) = j \circ (k \circ j) \circ k = j \circ 0 \circ k = 0,$$

so we may form the A -graded homology group $H(E) := \ker(j \circ k) / \operatorname{im}(j \circ k)$. Then we may form the triangle \mathcal{E}'

$$\begin{array}{ccc} i(D) & \xrightarrow{i'} & i(D) \\ & \swarrow k' & \searrow j' \\ & H(E) & \end{array}$$

where i' is the restriction of i to $i(D)$, while j' and k' are given by

$$j'(i(d)) = [j(d)] \quad \text{and} \quad k'([e]) = k(e).$$

The map j' is well-defined since if $i(d) = i(d')$ then $i(d - d') = 0$, so that $d - d' \in \ker i = \operatorname{im} k$, meaning $d - d' = k(e)$ for some $e \in E$, so that

$$j(d) - j(d') = j(d - d') = j(k(e)) \in \operatorname{im}(j \circ k)$$

is a boundary, so that $[j(d)] = [j(d')]$. Similarly k' is well defined since if $[e] = [e']$ then $e - e' \in \operatorname{im}(j \circ k)$, which implies $e - e' = j(k(e''))$ for some $e'' \in E$, so that

$$k(e) - k(e') = k(e - e') = k(j(k(e''))) = 0,$$

where the last equality follows by the fact that $k \circ j = 0$. Further note that $i(D)$ and $H(E)$ are A -graded by ?? and ??, in which case by unravelling definitions, each of i', j' , and k' are A -graded homomorphisms with

$$\deg i' = \deg i, \quad \deg j' = \deg j - \deg i, \quad \text{and} \quad \deg k' = \deg k.$$

We call \mathcal{E}' the *derived couple* of \mathcal{E} . A diagram chase (left to the reader, or see [2, Lemma 1.10]) yields that \mathcal{E}' is an exact couple.

If we iterate the process of taking the exact couple r times, the result is called the r^{th} derived couple \mathcal{E}_r of \mathcal{E} .

$$\begin{array}{ccc} D_r & \xrightarrow{i} & D_r \\ & \swarrow k \quad \searrow j^{(r)} & \\ & E_r & \end{array}$$

Here $D_r = i^r(D)$ is a subgroup of D , and $E_r = H(E_{r-1})$ is a subquotient of E . The maps i and k are induced from the i and k of \mathcal{E} , while $j^{(r)}$ sends $[i^r(d)]$ to $[j(d)]$. In particular, by induction it can be seen that $\deg j^{(r)} = \deg j - r \cdot \deg i$, and the degrees of i and k remain unchanged as we take successive derived couples.

Definition 0.6 (The spectral sequence associated to an exact couple). An A -graded exact couple $\mathcal{E} = (D, E; i, j, k)$ gives rise to a spectral sequence $(E_r, d_r)_{r \geq 0}$, where $E_0 = E$, $d_0 = j \circ k$, and for $r > 0$, E_r is defined above and d_r is the composition $j^{(r)} \circ k$.

In practice, we will always shift everything up a degree by re-defining $E_r := E_{r-1}$ and $d_r := d_{r-1}$, so we get a spectral sequence $(E_r, d_r)_{r \geq 1}$ with $E^1 = E$ and $d^1 = j \circ k$. Then it follows that the differential $d^r = j^{(r-1)} \circ k$ has degree

$$\deg j^{(r-1)} + \deg k = \deg j - (r-1) \cdot \deg i + \deg k.$$

Remark 0.7. Given an exact couple $\mathcal{E} = (D, E; i, j, k)$, we can define A -graded subgroups $Z_r = k^{-1}(i^r(D)) \subseteq E$ and $B_r = j(\ker(i^r)) \subseteq E$ for $r \geq 1$. By induction, it is straightforward to check that we have inclusions

$$B_1 \subseteq B_2 \subseteq B_3 \subseteq \cdots \subseteq \operatorname{im} j = \ker k \subseteq \cdots \subseteq Z_3 \subseteq Z_2 \subseteq Z_1$$

and that the maps

$$Z_r \rightarrow E_r$$

sending an element e to its class $[[\cdots [e] \cdots]]$ has kernel B_r , so we have identifications $E_r = Z_r/B_r$ as A -graded abelian groups. Let $e \in Z_r$, so $k(e) = i^r(d)$ for some $d \in D$. Then under this identification, it can be seen that the map $d_{r+1} : Z_r/B_r \rightarrow Z_r/B_r$ sends the coset $e + B_r \in Z_r/B_r$ to the coset $j(d) + B_r$, and that $\ker d_r = Z_r$ and $\operatorname{im} d_r = B_r$ for all $r \geq 1$.

Henceforth, we fix an A -graded exact couple $\mathcal{E} = (D, E; i, j, k)$, and we let $(E_r, d_r)_{r \geq 1}$ denote the associated A -graded spectral sequence. We make the identifications given by the above remark, so we assume E_r is the A -graded abelian group Z_r/B_r for all $r \geq 1$, so that in particular for all $a \in A$ we have identifications $E_r^a = Z_r^a/B_r^a$ (by ??).