

---

1 A simulation-based assessment of the relation between Stone Age  
2 sites and relative sea-level change along the Norwegian Skagerrak  
3 coast

4 Isak Roalkvam<sup>1,\*</sup>

5 29 April, 2022

6 **Abstract**

7 A central premise for Norwegian Stone Age archaeology is that most coastal sites were located on or  
8 close to the contemporary shoreline when they were in use. By reconstructing the trajectory of relative  
9 sea-level change, this offers a dating method termed shoreline dating, separate from both radiometric and  
10 typological methods which has been widely applied. However, while the potentially immense benefits  
11 of an additional source of temporal data is undeniable, this also comes with a series of uncertainties  
12 which, due to the relative rarity of the method, has been under limited scrutiny compared to more  
13 traditional dating techniques in archaeology. This paper attempts to remedy this by quantifying the  
14 spatial relationship between Stone Age sites located beneath the marine limit and the prehistoric shoreline  
15 along the Norwegian Skagerrak coast. This is done by combining independent temporal data on the use  
16 of the sites in the form of  $^{14}\text{C}$ -dates and the reconstruction of local shoreline displacement through Monte  
17 Carlo simulation. The findings largely confirm previous evaluations of this relationship, indicating that  
18 sites older than the Late Neolithic tend to have been located on or close to the shoreline when they were  
19 occupied. Drawing on the quantitative nature of the results, a new and formalised method for the shoreline  
20 dating of sites in the region is proposed, and compared to previous applications of the technique.

21 <sup>1</sup> University of Oslo, Institute of Archaeology, Conservation and History

22 \* Correspondence: Isak Roalkvam <isak.roalkvam@iakh.uio.no>

23 **Highlights**

- 24 • Simulates the spatial relation between sites and the prehistoric shoreline  
25 • Quantification of horizontal, topographic and vertical distance  
26 • Confirms close association between sites and the contemporaneous shoreline  
27 • Proposes a formalised method for the shoreline dating of pre-Late Neolithic sites

28 Keywords: Shoreline dating; Stone Age; Settlement patterns; Scandinavia; Relative sea-level change

29 **1 Introduction**

30 The post-glacial relative sea-level fall that characterises large areas of Fennoscandia is fundamental to  
31 its archaeology. This follows not only from the dramatic changes to the landscape that this process has  
32 represented throughout prehistory, but also from the fact that if archaeological phenomena were situated  
33 close to the contemporary shoreline when they were in use, a reconstruction of the trajectory of shoreline  
34 displacement can be used to date these phenomena based on their altitude relative to the present day sea-level.  
35 This method, also called shoreline dating, has long history of use in the region and is frequently applied  
36 to assign an approximate date to diverse archaeological phenomena such as rock art, grave cairns, various  
37 harbour and sea-side constructions and, as is the focus of this study, Stone Age sites (e.g. Åkerlund 1996;

<sup>38</sup> Bjerck 2005; Nordqvist 1995; Schmitt et al. 2009; Sognnes 2003; Tallavaara and Pesonen 2020; Wikell et al. 2009).

<sup>40</sup> The close association between Stone Age settlements in the northern parts of Scandinavia and shifting prehistoric shorelines was established by the end of the 19th century (De Geer 1896), and was first applied as a dating method at the turn of the century (Brøgger 1905; Hollender 1901). Shoreline dating has been fundamental to Norwegian Stone Age archaeology ever since (e.g. Berg-Hansen 2009; Bjerck 1990, 2008a; Breivik 2014; Johansen 1963; Mikkelsen 1975; Mjærum 2022; Nummedal 1923; Shetelig 1922; Solheim and Persson 2018). The method is used both independently, and to compliment other sources of temporal data such as typological indicators or radiometric dates. However, given the coarse and fuzzy resolution of established typological frameworks, the vast amount of surveyed sites that only contain generic lithic debitage that could hail from any part of the period, and as the conditions for the preservation of organic material is typically poor in Norway, dating with reference to shoreline displacement is often the only and most precise method by which one can hope to date the sites. Shoreline dating is consequently fundamental to our understanding of the Norwegian Stone Age. This is both because it is central to the temporal framework on which our understanding of the period is based, but also because the method is only applicable so long as the societies in question have continuously settled on or close to the contemporary shoreline. Consequently, adherence or deviation from this pattern also has major implications for the socio-economic foundations of the societies in question.

<sup>56</sup> Despite its important role for Norwegian Stone Age archaeology, the applicability of dating by reference to shoreline displacement has only been evaluated using relatively coarse methods. The aim of this paper is to provide a systematic and comprehensive review of the degree to which radiocarbon dates correspond with the dates informed by our current knowledge of shoreline displacement in a larger area of south-eastern Norway, using a more refined methodological approach. The goal is to quantify the degree to which the assumption of shore-bound settlement holds through the Stone Age, and in turn have this inform an improved method for shoreline dating. As presented in more detail below, this problem involves the combined evaluation of three major analytical dimensions. One is the questions of when the sites were in use, the second pertains to the reconstruction of the contemporaneous sea-level, and the third follows from the fact that the relation between site and shoreline is inherently spatial. Taking inspiration from studies that have integrated various sources of spatio-temporal uncertainty through Monte Carlo simulation (e.g. Bevan et al. 2013; Crema et al. 2010; Crema 2012, 2015; Yubero-Gómez et al. 2016), a similar approach is adopted here.

## <sup>68</sup> 2 Background

<sup>69</sup> Relative sea-level (RSL) can be defined as the mean elevation of the surface of the sea relative to land, or, more formally, the difference in elevation between the geoid and the surface of the Earth as measured from the Earth's centre (Shennan 2015). Variation in this relative distance follow from a range of effects (e.g. Milne et al. 2009). Of central importance here is eustasy and istostasy. The eustatic sea-level is understood as the sea-level if the water has been evenly distributed across the Earth's surface without adjusting for variation in the rigidity of the Earth, its rotation, or the self-gravitation inherent to the water body itself (Shennan 2015). The eustatic sea-level is mainly impacted by glaciation and de-glaciation, which can bind or release large amounts of water into the oceans (Mörner 1976). Istostasy, on the other hand, pertains to adjustments in the crust to regain gravitational equilibrium relative to the underlying viscous mantle. This can be the result of glacial istostasy, which follows from glaciation and de-glaciation and corresponding loading and unloading of weight, as well as from erosion of the crust, which causes its weight to be redistributed. These effects thus causes the lithosphere to either subside due to increased weight, or to rebound and lift upwards due to lower weight (Milne 2015).

<sup>82</sup> Following the end of the Weichselian and the final retreat of the Fennoscandian Ice Sheet (e.g. Hughes et al. 2016; Stroeven et al. 2016, see Figure 1), the isostatic rebound has been so severe that most areas of Norway have been subject to a continuous relative sea-level regression, despite corresponding eustatic sea-level rise (e.g. Mörner 1979; Svendsen and Mangerud 1987). In other words, the RSL has been dropping throughout prehistory. As this process is the result of glacial loading, the rate of uplift is more severe towards the centre

87 of the ice sheet. Thus, some areas on the outer coast have had a more stable RSL or been subject to marine  
 88 transgression (e.g. Romundset et al. 2015; Svendsen and Mangerud 1987). These conditions are directly  
 89 reflected in the archaeological record. In areas where the sea-level has been stable over longer periods of  
 90 time, people have often reused coastal site locations multiple times and over long time-spans, creating a  
 91 mix of settlement phases that are difficult to disentangle (e.g. Hagen 1963; Reitan and Berg-Hansen 2009).  
 92 Transgression phases, on the other hand, can lead to complete destruction of the sites, bury them in marine  
 93 sediments, or in the outermost periphery, leave them still submerged today (Bjerck 2008a; Glørstad et al.  
 94 2020). This can lead to a hiatus in the archaeological record for certain sub-phases in the impacted areas.  
 95 Comparatively, given a continuous and still ongoing shoreline regression from as high as c. 220 m above  
 96 present sea-level in the inner Oslo fjord, any one location in south-eastern Norway has only been shore-bound  
 97 within a relatively limited time-span, and the sites have not been impacted by any transgressions (Hafsten  
 98 1957, 1983; Romundset et al. 2018; Sørensen 1979). This makes the region especially useful for evaluating  
 99 the assumption of a shore-bound settlement pattern over a long and continuous time-span.




Figure 1: Deglaciation at 1000 year intervals from c. 17–8 kyr BCE. The study area defined later in the text  
 is marked with a red outline (deglaciation data from Hughes et al. 2016, but see also Romundset et al. 2019  
 in relation to the study area).

100 The method of shoreline dating has been met with scepticism as related to the fundamental premise that  
 101 most sites would have been consistently shore-bound, been characterised as a relative dating method for sites  
 102 located at different elevations within a constrained geographical area, or been argued to offer no more than  
 103 an earliest possible date for when a site could have been in use (see review by Nordqvist 1999). The most

common application in Norway has arguably been to use shoreline dating to provide an approximate date for the occupation of the sites, often in combination with other dating methods (see for example chapters in Glørstad 2002, 2003, 2004; Jakslund 2001, 2012a, 2012b; Jakslund and Persson 2014; Melvold and Persson 2014; Reitan and Persson 2014; Reitan and Sundström 2018; Solheim 2017 and below). Recently the method has also been used independently to date a larger number sites to get a general impression of site frequency over time. This is done by aggregating point estimates of shoreline dates in 100, 200 or 500 year bins (Breivik 2014; Breivik and Bjerck 2018; Fossum 2020; Mjærum 2022; Nielsen 2021; Solheim and Persson 2018; see also Jørgensen et al. 2020; Tallavaara and Pesonen 2020). In his review, Nordqvist (1999) argues that there can be little doubt concerning the general applicability of the method – what is less clear is the level of reliability and chronological resolution that it can offer (see also Johansen 1963).

The shore-bound settlement location of prehistoric hunter-fisher-gatherers in Norway is generally believed to follow both from the exploitation of aquatic resources and from movement and communication, which would have been efficient on waterways (Bjerck 1990, 2017; Brøgger 1905:166; also discussed by Berg-Hansen 2009; Bergsvik 2009). The same logic has also been extended to the hinter- and inland regions, where sites are to be predominantly located along rivers and lakes (Brøgger 1905:166; Glørstad 2010:57–87; but see also Gundersen 2013; Mjærum 2018; Schülke 2020). This is to take a dramatic turn at the transition to the Late Neolithic, around 2400 BCE, with the introduction of the Neolithic proper (Prescott 2020; cf. Solheim 2021). The introduction of a comprehensive Neolithic cultural package, including a shift to agro-pastoralism and the introduction of the farm is to have led site locations to be more withdrawn from the shoreline (e.g. Bakka and Kaland 1971; Østmo 2008:223; Prescott 2020). That is not to say that waterways and aquatic resources were no longer exploited, but rather that these activities would not have been as tightly integrated with settlement and tool-production areas as in preceding periods (Glørstad 2012). At an earlier stage, at the transition to the Early Neolithic (c. 3900 BCE), pottery is introduced to the sites, and there are some indications of an initial uptake of agriculture at some sites in the Oslo fjord region. However, this appears to be small in scale and is believed to be combined with a continued and predominantly hunter-gatherer life-way, possibly followed by a complete de-Neolithisation in the Middle Neolithic (Hinsch 1955; Nielsen et al. 2019; Østmo 1988:225–227). Nielsen (2021) has recently argued that the initial uptake of agriculture in Early Neolithic south-eastern Norway is combined with a more complex settlement pattern, and that a simple foraging/agricultural dichotomy would underplay the variation present in the Early and Middle Neolithic settlement data (see also e.g. Amundsen et al. 2006; Østmo 1988; Solheim 2012:74). Seen in relation to the question of interest here, the empirical expectation for the above outlined development would thus be a predominantly shore-bound settlement in the Mesolithic, possibly followed by a more varied association between sites and the shore-line with the transition to the Early Neolithic around 3900 BCE, and finally a decisive shift with the Late Neolithic c. 2400 BCE.

Based on the generally accepted premise that most pre-Late Neolithic sites in south-eastern Norway located lower than the marine limit were situated on or close to the contemporaneous shoreline, it is common to err on the side of a shore-bound site location unless there is strong evidence to suggest otherwise. This is for example reflected in survey projects, which are often guided by both a digital and mental reconstruction of past sea-levels (e.g. Berg-Hansen 2009; Eskeland 2017). Similarly, following an excavation, if typological indicators in the assemblages correspond with available shoreline displacement curves, a shore-bound site location is often assumed, even if the typologically informed date-span is too wide to decisively verify this. It is also common to combine this with a qualitative consideration of the landscape surrounding the sites, and an evaluation of the degree to which the site location would have been sensible if the site was not shore bound (e.g. Jakslund 2014; Johansen 1963; Nummedal 1923). This can for example pertain to accessibility. If the site is situated on a ledge in a steep and jagged area of the present day landscape it would make intuitive sense that the site was in use when the ocean reached closer to its elevation, as the site would have been accessible by means of watercraft. Although it appears that the arguments for such site locations are sensible and can for the most part be assumed to hold, comprehensive evaluations and attempts at quantification of this tendency are relatively few (see also Ilves and Darmark 2011).

One of the more extensive evaluations of the relationship between archaeological radiocarbon dates and RSL-change was done by Solheim and colleagues (Breivik et al. 2018; Solheim 2020), who compared 102 radiocarbon dates from 33 Mesolithic sites on the western side of the Oslo fjord to the displacement curve

for the Larvik area. They found an overlap between the probability density of the radiocarbon dates with the shoreline displacement curve for 86.5% of the sites. However, where there was a discrepancy, the main occupation of the sites are still believed to have been shore-bound rather than associated with the deviating  $^{14}\text{C}$ -dates. This is based on typological and technological characteristics of the assemblages. Whether these mismatches represent later shorter visits that are responsible for the younger radiocarbon dates, or whether these dates are entirely erroneous can be difficult to evaluate (e.g. Persson 2008; Schülke 2020). However, this distinction is not deemed critical here, as what is of interest is settlements and tool-production areas as evidenced by artefact inventories or multiple site features. Not remnants of stays as ephemeral to only be discernible by isolated features or dubious  $^{14}\text{C}$ -dates. The evaluation of the relevance of radiocarbon dates to settlement activity will here therefore be entirely dependent on, and follow the discretion of the original excavation reports.

Other previous evaluations of the correspondence between radiocarbon- and RSL-informed dates have typically followed the same structure as that of Breivik et al. (2018), involving a visual inspection of radiocarbon probability density functions plotted against local shoreline displacement curves based on the elevation of the site (e.g. Åkerlund et al. 1995; Åstveit 2018; Solheim 2020; see also Bjerck 2008b; Kleppe 1985; Ramstad 2009). This approach has a couple of limitations. First of all, the displacement curves are sometimes applied directly to larger study areas, with only some studies having taken the variable uplift-rates into account when performing this comparison (e.g. Åstveit 2018; Fossum 2020; Møller 1987; Persson 2008). Secondly, with this method, the wider the uncertainty range associated with either radiocarbon date or displacement curve, the higher the probability that the confidence intervals overlap, and the higher the probability that we conclude in favour of our hypothesis. This thus leads to an inferential framework that favours uncertainty, which is hardly desirable. In statistical terms this follows from the fact that while one cannot conclude that two dates are different if their confidence intervals overlap, this does not necessarily mean that they are the same. The question thus necessitates a flip from a null-hypothesis of no significant difference, to one of equivalence (e.g. Lakens et al. 2018), as the question of interest is effectively one of synchronicity between events (cf. Parnell et al. 2008). Another limitation of this often-employed method is that it only takes into account the vertical distance between the sites and the sea-level. While this is the main parameter of interest for shoreline dating, the practical implications of a vertical difference in RSL will be highly dependent on local topography and bathymetry. RSL-change can have more dramatic consequences in a landscape characterised by a low relief, as the horizontal displacement of the shoreline will be greater. Taking the spatial nature of the relationship between site and shoreline into account will consequently help get more directly at the behavioural dimension of this relation, and help move the analysis beyond a purely instrumental consideration of the applicability of shoreline dating.

### 3 Data

To get at the relationship between sites and the contemporaneous shoreline, this analysis was dependent on a study area with good control of the trajectory of prehistoric shoreline displacement. While there is displacement data available for other areas of south-eastern Norway (e.g. Hafsten 1957; Sørensen 1979, 1999; and recent compilation by Creel et al. 2022), considerable methodological developments in recent years means that the most well-established displacement curves are from the region stretching from Horten county in the north-east, to Arendal in the south-west. This area has newly compiled displacement curves for Horten (**romundset2021?**), Larvik (Sørensen et al. in prep; Sørensen, Henningsmoen, et al. 2014; Sørensen, Høeg, et al. 2014), Tvedstrand (Romundset 2018; Romundset et al. 2018), and Arendal (Romundset 2018).

The employed shoreline displacement data is based on the so-called isolation basin method (e.g. Kjemperud 1986; Romundset et al. 2011), which involves extracting cores from a series of basins situated on bedrock at different elevations beneath the marine limit, and dating the transition from marine to lacustrine sediments. Each basin thus represent a high precision sea-level index point (SLIP) which are combined using what has been termed the isobase method to devise a continuous time series for RSL-change, projected to a common isobase (see Creel et al. 2022:5). Furthermore, to minimise the impact of variable uplift rates, the basins are located in a as constrained area of the landscape as possible. Following from the morphology of the

retreating ice sheet, the uplift is more severe towards the north-east, meaning that this needs to be adjusted for in the case that any basins are located any significant distance from the common isobase perpendicular to this gradient (Figure 2). The SLIPs indicate the isolation of the basins from the highest astronomical tide, which is adjusted to mean sea-level in the compilation of the displacement curves, based on the present day tidal range. This is assumed to have been the same throughout the Holocene (Sørensen, Henningsmoen, et al. 2014:44). The highest astronomical tide in the study area reaches around 30cm above mean sea-level (Norwegian Mapping Authority 2021:30cm at the standard port Helgeroa in Larvik). Furthermore, the confidence bands of the displacement curves and their trajectory are quite complex constructs, and are the integrated result of both expert knowledge and more objectively quantifiable parameters. The reason for this is in part that the curves do not only contain uncertainty as related to radiometric dates, which are well defined, but also hold potential error as related to the interpretation and analysis of sediment cores, the nature and condition of the basin outlets and the adjustment to a common isobase, to name but a few (e.g. Romundset et al. 2011, 2019; for an alternative approach see Creel et al. 2022). For more details and evaluations done for the compilation of each curve, the reader is therefore referred to the individual publications.

The archaeological data compiled for the analysis consists of excavated Stone Age sites with available spatial data from the coastal region between Horten county in the north-east, to Arendal in the south-west (Figure 2). These number 155 sites. Of these, 91 sites are associated with a total of 547 radiocarbon dates. Of these, in turn, 67 sites are related to the 259 radiocarbon dates that fall within the Stone Age (9500–1700 BCE), with 95% probability. These sites and  $^{14}\text{C}$ -dates form the basis for the analysis. Spatial data in the form of site limits and features, as defined by the excavating archaeologists, were retrieved from local databases at the Museum of Cultural History—the institution responsible for archaeological excavations in the region. In the compiled dataset, each radiocarbon date has been associated with the site features or excavation unit from where they originate, or, where these weren't available, the spatial limit of the entire site. Due to somewhat variable practices between excavations, what available spatial geometry best represents the site limit was decided based on an evaluation of the excavation reports. This means that the limits are variably given as that defined during initial survey, area de-turfed before excavation, area stripped with excavator following the excavation, manually excavated area, or convex hull polygons generated around the site features.

Three of the sites have been associated with agriculture, either directly or in the form building structures. The first is Nordby 1 at which the  $^{14}\text{C}$ -dates are associated with a Late Neolithic long-house (Gjerpe and Bukkemoen 2008). The Middle Neolithic phase at Kvastad A2 (Stokke and Reitan 2018) and Late Neolithic phase at Nauen A (Persson 2008) are both directly related to farming activities. Both of these sites also have radiocarbon dates and lithic inventory associated with Mesolithic forager activities. Following from the expected deviance from the settlement patterns that are to characterise forager sites, these agricultural phases are highlighted in the analysis below. Finally, Nielsen (2021) has recently suggested that Early and Middle Neolithic features from the otherwise younger sites Bratsberg (Wenn 2012) and Larønningen (Røberg 2012) could be related to early agricultural activity in the Oslo fjord region. Due to the uncertain and somewhat speculative nature of this suggestion, these are omitted here.

The elevation data used for the analysis is a digital terrain model (DTM) freely available from the Norwegian Mapping Authority (Norwegian Mapping Authority 2018, <https://hoydedata.no>). It was here opted for the 10m resolution DTM rather than the higher-resolution 1m version. In addition to resulting in considerably less processing time, the higher resolution elevation model is more vulnerable to smaller-scale modern disturbances that the 10m version is not impacted by. The 10m resolution DTM of the study area is a down-sampled version of the 1m version and has a height accuracy with a systematic error of 0.1m (Norwegian Mapping Authority 2018). All data and R programming code (R Core Team 2021) required to run the analyses, as well as the derived data are freely available in an online repository at <https://osf.io/7f9su/>, organised as a digital research compendium following Marwick et al. (2018).




Figure 2: A) Distribution of the 67 analysed sites relative to the isobases of the displacement curves. The isobases have a direction of  $327^\circ$  (Romundset et al. 2018), B) Displacement curves. Note the increasing<sup>7</sup> steepness of the curves towards the north-east.

---

## 252 4 Methods

253 Shoreline dating is based on the spatial relationship between two phenomena, occupation of sites and shoreline  
254 displacement, each associated with their own range of temporal uncertainty. The first task was therefore to  
255 ascribe likely date ranges and associated uncertainty to these dimensions. To take account of the gradient in  
256 the isostatic rebound, the trajectory of shoreline displacement was first interpolated to each site location based  
257 on the distance to the isobases of the displacement curves, using inverse distance weighting (e.g. Conolly  
258 2020; Conolly and Lake 2006:94–97). This was done for each year along the entirety of the curves, weighting  
259 the interpolation by the squared inverse of the distances. The result of this process is shown for an example  
260 site in Figure 3. For the date ranges associated with the sites, all radiocarbon dates were first individually  
261 calibrated using the IntCal20 calibration curve (Reimer et al. 2020) using OxCal v4.4.4 (Bronk Ramsey 2009)  
262 through the oxcAAR package for R (Hinz et al. 2021). Radiocarbon dates associated with each site were  
263 then grouped if they overlapped with 99.7% probability, meaning these were effectively taken to represent  
264 the same event, here termed settlement or site phase. In the case where there are multiple dates believed to  
265 belong to a single settlement phase, these were modelled using the Boundary function in OxCal and then  
266 summed. Multiple phases at a single site were treated as independent of each other.

267 The excavation of archaeological sites typically follow from residential and commercial development, as well as  
268 the expansion of infrastructure. As the data collection for the employed DTM was begun by the Norwegian  
269 Mapping Authority in 2016, the area of the DTM immediately surrounding the sites has sometimes been  
270 severely impacted by disturbances after the excavation. In addition to employing 10m resolution DTM to  
271 alleviate some of these issues, this also necessitated some additional editing of the elevation raster. This  
272 involved manually defining the extent of problem areas such as railways, highways, quarries and the like.  
273 The DTM values on these were then set to missing, and new elevation values were interpolated from the  
274 surrounding terrain. This was done using regularised spline interpolation with tension (e.g. Conolly 2020),  
275 using the default settings of r.fillnulls from GRASS GIS (GRASS Development Team 2017) in R through the  
276 package rgrass7 (Bivand 2021). In addition to code and original spatial data being available in the digital  
277 research compendium for this paper, the analysis of each individual site is presented in the supplementary  
278 material where it has been noted when the area surrounding a site has been edited in this manner.

279 Armed with a likely date range for the occupation(s) of each site, an estimated trajectory of relative sea-level  
280 change at that location, and a DTM edited to remove substantial modern disturbances, the simulations were  
281 performed. A single simulation run involved first drawing a single year from the posterior density estimate of  
282 a given occupation phase of a site (Figure 4). This year then has a corresponding likely elevation range for  
283 the contemporaneous shoreline from which an elevation value was drawn uniformly, using intervals of 5cm.  
284 The sea-level was then raised to this elevation on the DTM by defining all elevation values at or below this  
285 altitude as missing. Polygons were then created from the resulting areas with missing values. The horizontal  
286 distance was then found by measuring the shortest distance between site and sea polygons, and the vertical  
287 distance by subtracting the elevation of the sea-level from the lowest elevation of the site polygon. The topographic  
288 distance between site and sea was also found by measuring the distance while taking into account  
289 the slope of the terrain on the DTM. This was done using the topoDistance package for R (Wang 2019).  
290 The topographic distance was measured between the site polygon and the horizontally closest point on the  
291 shoreline. This means that the distance is not necessarily measured as the closest topographic distance to the  
292 shoreline, but rather as the shortest topographic path to the horizontally closest point on the shoreline. Not  
293 finding the topographically closest point significantly reduced the computational cost of the analysis, and is  
294 deemed unlikely to have a considerable impact on the results given the distances considered. The shortest  
295 topographic path was found using the Moore neighbourhood of eight cells (e.g. Conolly and Lake 2006:253;  
296 Herzog 2013). In the case where the sea polygons intersects the site polygon, all distance measures were  
297 set to zero. In the case that the sea polygons completely contain the site, the horizontal and topographic  
298 distance measures were made negative, and the vertical distance was instead measured to the highest point  
299 on the site polygon. While it is safe to assume that an archaeological site was not occupied when it was  
300 located beneath sea-level, a negative result can reflect the inherent uncertainty in this procedure, and might  
301 also help identify discrepancies in displacement data or radiocarbon dates. Negative values were therefore  
302 retained with the exception of for the sites Gunnarsrød 5 and Pjonkerød R1, where the negative values are  
303 believed to result from modern disturbances in the DTM rather than the  $^{14}\text{C}$ -dates or displacement curves

304 (see supplementary material for more details).

305 This process was repeated 1000 times for each phase for each site. The choice of 1000 simulation runs follows  
306 from an evaluation of when the mean distances between site and shoreline converged when running 5000  
307 iterations of the simulation on the site Hovland 5, available in the supplementary material (cf. Crema et al.  
308 2010:1125). Hovland 5 was chosen for this evaluation as it has a fairly imprecise date and is located in area  
309 of quite complex topography.

## 310 5 Simulation results

311 Overall, as is indicated by the measures for central tendency and the almost solid line along the 0m mark on  
312 the y-axes, the simulations show that the sites tend to have been situated close to the shoreline when they  
313 were in use (Figure 6). Some of the sites are situated considerable distances from the shoreline when the dates  
314 believed to be erroneous in the original reports are included (Figure 6A), but if one accepts the interpretation  
315 that these do not date the main occupation of the sites, as is indicated by the artefact inventories, Figure  
316 6B gives considerable support to the notion that the sites were in use when they were situated on or close  
317 to the contemporaneous shoreline. The distances for some of the earliest sites appears somewhat high, but  
318 this can likely be explained as the result of the steepness of the displacement curves for the earliest part of  
319 the Holocene (Figure 2B), which leads the uncertainty of the  $^{14}\text{C}$ -dates to give a wider possible elevation  
320 range for the simulated sea-level. Another immediately striking result is the apparent deviation from the  
321 shoreline towards the end of the Stone Age. From around 2500 BCE several sites are situated a considerable  
322 distance from the shoreline, and while a couple remain horizontally and topographically close, most appear  
323 to be elevated a considerable distance from the sea-level, as indicated on the plot for vertical distance. There  
324 are also a couple of sites located some distance from the shoreline just after 4000 BCE. While the sample size  
325 is limited, this would thus be in line with a development that sees an increase in settlements located in the  
326 immediate inland around this time.

327 The negative values around 8000 BCE originate from the sites Løvås 1, 2 and 3. These are recently excavated,  
328 well-dated sites situated in a relatively undisturbed area of the landscape. While there would be a danger of  
329 circularity of having archaeological sites inform a reconstruction RSL-change, and in turn use these to evaluate  
330 the degree of shore-bound settlement, the sites do clearly represent a upper limit for the sea-level, as they  
331 would not have been in use when located under water. It could therefore seem that the Løvås sites represent  
332 a case where the archaeological material indicates a slight discrepancy in the geological reconstruction of  
333 shoreline displacement in the area.

334 Accepting that shoreline dating appears to loose utility around the transition to the Late Neolithic, as  
335 indicated by the clear deviation in site location from the shoreline after this, the results for from Figure 6B is  
336 given again in Figure 7A, excluding all simulation results younger than 2500 BCE. Furthermore, all negative  
337 values have here been set to zero, under the assumption that these result from uncertainty or errors in the  
338 data, and not actual site locations. The resulting best point estimate for the vertical distance between sites  
339 and shoreline for the pre-Late Neolithic is given by the median at 4m, while 95% of the values fall within the  
340 range 0–18m. That is, for 95% of the cases, the shoreline was simulated to be situated on or down to 18m  
341 below the site location. While these values remain the same when only the Mesolithic dates are included  
342 (Figure 7B), the mean and standard deviation are slightly constrained. Furthermore, while the median for  
343 horizontal and topographic distance is only 10m across all plots in Figure 7, the variation in the statistics for  
344 dispersion is greater, illustrating the point that minor variations in vertical distance can have substantial  
345 consequences for these distance measures, depending on the surrounding topography.

346 An exponential function has been fit to the distributions for vertical distance using maximum likelihood  
347 estimation (Figure 7). While it makes theoretical sense that a process of exponential decay explains this  
348 relationship, it is also clear that this does not perfectly match the data. However, this can at least in part be  
349 related to methodological factors, where the accumulation of distance-values on the 0m mark likely follow  
350 from forcing negative values to zero, from the resolution of the spatial data, and from defining intersecting  
351 sea- and site polygon as having a distance of zero. If one accepts this, the probability density function for




Figure 3: Example site Hegna vest 1 (Fossum 2017). A) Location of the site on the edited 10m resolution DTM. The red outline is the site limit. B) Radiocarbon dates associated with the site. Fill colour indicates what dates are assumed to belong to the same settlement phase. Multiple dates are modelled using the Boundary function in OxCal and then summed. The red outline indicates that the date does not match the typological indicators in the artefact assemblage of the site. C) The location of the site within the study area relative to isobases of the employed displacement curves. D) Displacement curve interpolated to the site location.




Figure 4: Example of a single simulation run on the site Hegna vest 1. A) The simulation starts by drawing a single year from the posterior density estimate. B) This then corresponds to an elevation range on the interpolated displacement curve. A single elevation is drawn uniformly from this range using 5cm intervals. C) The sea-level is then adjusted on the DTM to this elevation and the various distance measures are found. D) The numerical result of the simulation run.




Figure 5: The result of 1000 simulation runs for each of the two groups of dates on the site Hegna vest 1. The first column of plots shows the radiocarbon probability density function from where dates were drawn during simulation. The second column displays the result of simulating the raised sea-level 1000 times. The more opaque the colour, the more times the sea-level was simulated to that location. The third column shows violin plots of the different distance measures across all simulations.

---



**A****B**

Figure 6: The result of running the analysis across all sites. Each data point is plotted with some transparency, meaning that the more intense the colour, the more often those values occurred. Results associated with agricultural activities are plotted in grey. The first row A) shows the result of including all dates to the Stone Age, including those seen as otherwise unrelated to the main occupation of the sites. The second row B) shows the result of excluding these. The table under each plot lists some corresponding statistics for central tendency and dispersion.

---



**A****B**

Figure 7: Histograms showing the simulated distance from the shoreline using radiocarbon dates corresponding to the site inventories. Negative values have been set to zero. A) Simulated results older than 2500 BCE, and B) simulated results older than 4000 BCE.

---

352 exponential decay can be used to characterise the vertical distance between sites and the shoreline, and be  
353 used to inform a method for shoreline dating that takes this into account.

354 **6 Shoreline dating**

355 The procedure for shoreline dating to be outlined is thus aimed at determining the likely age of the occupation  
356 of a site based on its altitude above present day sea-level, with reference to shoreline displacement and  
357 the likely elevation of the site above the sea-level when it was in use. For simplicity, this is conceptually  
358 treated a single event and thus the possibility of multiple or continuous phases of occupation is not treated  
359 explicitly. This leads the problem to become analogous to that of the calibration of a radiocarbon date  
360 (Bronk Ramsey 2009, see also Figure 8). First, finding the elevation of the sea-level at the time the site was  
361 in use is dependent on the present day elevation of the site  $\alpha$  and the distance between site and the shoreline  
362  $D$ . Based on the simulation results above, the distance from the elevation of the site to the contemporaneous  
363 shoreline is defined by the probability density function for exponential decay:

$$p(\alpha - D) = \lambda e^{-\lambda(\alpha - D)} \quad (1)$$

364 where  $\lambda$  is the decay ratio. This can then be coupled with the trajectory of relative sea-level change to find  
365 the corresponding calendar date  $T$ . Here this is defined by a uniform probability density function over the  
366 range between the lower  $T_l$  and upper  $T_u$  bounds of the displacement curve that as been interpolated to the  
367 site location:

$$p(T|\alpha - D) = U[T_{l|\alpha-D}, T_{u|\alpha-D}] \quad (2)$$

368 Finding the probability for the date of the site then becomes a matter of transferring the probability of the  
369 distance between site and shoreline and that of the displacement curve to calendar dates:

$$p(T|\alpha - D) = p(T|\alpha - D)p(\alpha - D) \quad (3)$$

370 We can then get rid of parameter  $D$  by summarising over all possible distances between site and the shoreline.  
371 Given its elevation, the probability of the date for the occupation of the site is then:

$$p(T|\alpha) = \sum_D p(T|\alpha - D)p(\alpha - D) \quad (4)$$

372 An example implementation of this approach is given in Figure 9, where  $\lambda = 0.173$ . This is the decay ratio  
373 identified when considering all of the pre-Late Neolithic simulation results (Figure 7A). For the numerical  
374 implementation,  $D$  is here defined as a sequence of increments of 0.001m from the site elevation  $\alpha$ . In Figure 9  
375 the outlined procedure is used to shoreline date all of the sites from where this relationship was derived, with  
376 the Late Neolithic sites also included for illustrative purposes. Following from having defined the distance  
377 between intersecting sea- and site polygons as zero during simulations, the sites were all dated using the mean  
378 elevation of the site polygons to allow for some variation in elevation over the site limits. The synchronicity  
379 between radiocarbon and shoreline dates was then evaluated using the method presented by Parnell et al.  
380 (2008). Here, 100,000 age samples drawn from the probability density function of each shoreline date were  
381 subtracted from 100,000 age samples drawn from the corresponding modelled  $^{14}\text{C}$ -dates. The resulting range  
382 of the 95% highest density region (HDR, Hyndman 1996) was then checked to see if it crosses zero, in which  
383 case the dates are considered to be in agreement (Figure 10). When excluding the earliest occupation phase  
384 at Gunnarsrød 5, the deviation of which is to be expected based on issues with the DTM (see above), the  
385 shoreline date correspond to the radiocarbon dates in 58 out of 68 cases (84%). Only including dates modelled  
386 to be older than 2500 BCE with 95% probability, i.e. older than the Late Neolithic, improves this to 56 out  
387 of 61 cases (92%). When only including dates older than 4000 BCE with 95% probability, i.e. only Mesolithic

388 site phases, the success rate is further increased to 46/49 (94%). The three failed Mesolithic shoreline dates  
 389 are from the early sites Langemyr and Kvastad A2, with the likely implication that a lower decay ratio than  
 390 what is used for characterising the distance between site and shoreline for all sites in aggregate should be  
 391 used for sites known to be from the earliest part of the Mesolithic (see also Figure 6).




Figure 8: Shoreline dating of Hegna vest 1. The dashed line marks the mean elevation of the site polygon which is used to inform  $\alpha$  in the dating of the site. The exponential function decays with ratio  $\lambda$  from Figure 7A. The resulting shoreline date in grey is underlined with the 95% HDR in black.

## 392 7 Re-dating previously shoreline dated sites

393 To further explore the implementation for shoreline dating presented above, excavated and shoreline dated  
 394 Stone Age sites within the study area where  $^{14}\text{C}$ -dates are not available or these are not believed to date the  
 395 main occupation of the sites have been subjected to the outlined approach (Figure 11). The resulting dates  
 396 are compared to those originally proposed in the excavation reports for the sites (the numerical results are  
 397 available in the supplementary material). To avoid issues with recent disturbances on the DTM, the sites  
 398 have been dated based on the mean of the altitudes provided in the report for each site.

399 The comparison with previously reported dates is an illustrative, but unfair exercise for a few reasons. First  
 400 of all the dates provided in the reports are typically stated to be a very rough estimate, and are sometimes  
 401 given as a point estimate with an undefined, but implied or explicit uncertainty range. Secondly, seeing as  
 402 these reports are from various dates in time, many are based on now outdated data on RSL-change. Finally,  
 403 they are sometimes only meant to indicate a lower bound for when the sites could have been in use. Overall,  
 404 the results could, with some danger of circularity, suggest that shoreline dating has generally been applied  
 405 with a fairly reasonable degree of success, seeing as these dates have typically been interpreted and informed  
 406 research in an approximate manner (although see e.g. Roalkvam 2022). That being said, the results do also  
 407 indicate that shoreline dating has at times been applied with an exaggerated degree of precision. While  
 408 the implications of a more stable RSL-change for shoreline dating are well known, this also appears to be




Figure 9: The result of backwards shoreline dating the sites with radiocarbon dates corresponding to the artefact inventory using the method proposed here. The shoreline dates are plotted in grey and underlined<sup>17</sup> with the 95% HDR in black. These are plotted against the modelled radiocarbon dates, which are given colour from oldest to youngest occupation phase for each site, defined by non-overlapping dates at 99.7% probability.




Figure 10: Evaluation of the agreement between the shoreline dates and radiocarbon dates given in Figure 9. When the range of the 95% HDR for age difference crosses zero, the shoreline and radiocarbon dates are considered to be in agreement. Line segments with vertical bars indicate the HDR does not cross zero and that the dates do not correspond. The division and colour coding at the top of the plots reflect the division of site phases given in Figure 9.

409 somewhat under-appreciated in the practical implementation of the method. The results also highlight the  
410 spatial and temporal contingency of the method, illustrated by the variation in the range of the 95% HDRs  
411 for the dates. In some cases the method provides a very precise date range and in others it offers little more  
412 than a *terminus post quem*. This is dependent on the steepness of the displacement curves, leading to the  
413 general pattern of older sites situated towards the north-east getting more precise dates (cf. Figure 2B).  
414 Furthermore, as some of the date ranges extend well beyond major chronological divisions, even into the  
415 Iron Age, they could be severely and securely constrained with only cursory reference to typology. While  
416 this would be trivial in some cases, the nature and uncertainty inherent to the method still means that this  
417 is arguably a required exercise that should be explicitly performed. This also points to the possibility of  
418 drawing on other temporal data, for example within a Bayesian framework, to further improve the precision  
419 of the dates that can be achieved with shoreline dating.

420 Not least following from the fact that relatively few Preboreal  $^{14}\text{C}$ -dates associated with anthropogenic  
421 activity have been achieved in Norway (Åstveit 2018; Damlien and Solheim 2018; Kleppe 2018), the shoreline  
422 dating of the earliest sites is essential for understanding the pioneer settlement and the initial colonisation of  
423 the Scandinavian peninsula (e.g. Bang-Andersen 2012; Berg-Hansen 2018; Breivik 2014; Fuglestvedt 2012;  
424 Glørstad 2016). The shoreline dated Preboreal sites from the Brunlanes-project are among the earliest known  
425 sites in Norway (Jaksland 2012a, 2012b; Jaksland and Persson 2014). These have a distinct Early Mesolithic  
426 artefact inventory and are situated in a steep area of the landscape where it would be difficult to envision use  
427 of the sites after the sea retreated any significant distance from their location. In the original publication  
428 of the sites, Jaksland (2014) provides a thorough discussion of shoreline dating in general, and as used for  
429 the dating of the Brunlanes sites specifically. A comparison of his results and the ones achieved using the  
430 above-outlined approach are given in Figure 12A. The sites have been dated using what Jaksland (2014) gives  
431 as the lowest elevation of finds at each site, and by employing a exponential decay ratio of 0.13, to allow for  
432 more deviance in the distance between site and shoreline. This corresponds to the decay ratio for results  
433 older than 7000 BCE in Figure 7.

434 The small discrepancies between the achieved results mainly follow from the fact that a slightly updated  
435 version of the local displacement curve is applied here (cf. Sørensen et al. in prep). Jaksland's dates are  
436 given a flat 200 and 50 year uncertainty range starting from what he gives as the earliest possible date. The  
437 200 year uncertainty range is given if the sites were to be considered in isolation, while the argument for  
438 the uncertainty range of only 50 years is based on the location of the sites relative to each other. Since  
439 they are located in such a constrained and steep area of the landscape, the difference in elevation between  
440 the sites is argued to establish their relative date and thus constrain the uncertainty ranges so that they  
441 don't overlap. This information is not integrated in the approach outlined here, but could justify further  
442 reducing the uncertainty ranges. Although their accuracy is of course ultimately dependent on the veracity of  
443 the geological reconstruction, the high rate of RSL-change in this period does result in very precise dates.  
444 Above it was suggested that additional temporal data could be combined with the method to improve its  
445 accuracy and precision. This example, on the other hand, highlights the fact that the spatial nature of the  
446 method means that a consideration of the surrounding terrain and other sites can also help in increasing the  
447 precision of the method if this can be used to exclude certain sea-levels as unlikely for when a site was in use.  
448 One approach could also be to assess the spatial implication of a proposed shoreline date by simulating the  
449 adjusted sea-levels, as is done for Pauder 1 in Figure 12B, followed for example by a visual evaluation of the  
450 topography or by evaluating the distance and steepness of the slope to the shoreline. If this is developed  
451 further, it could conceivably be possible to exclude certain elevations as unlikely for the position of the  
452 shoreline when the site was in use. Such approaches would make less of an impact in this setting, where the  
453 95% HDR is already quite constrained, but could considerably improve the precision of the method in cases  
454 where RSL-change has been less severe (cf. Figure 11).

## 455 8 Concluding remarks

456 The most immediate contribution of this paper is what must be considered a confirmation of previous research  
457 into the relation between coastal Norwegian Stone Age sites and the prehistoric shoreline. This is indicated

$$\lambda = 0.173$$




Figure 11: Re-dating excavated and previously shoreline dated sites in the study area without radiocarbon dates or with radiocarbon dates that do not correspond to the artefact inventories. The 95% HDRs in grey<sup>20</sup> are compared to the dates originally proposed by the excavation reports in red.




Figure 12: Shoreline dating of the Brunlanes sites using site altitudes provided by Jakobsson (2014:tab.4). A) The result of applying the approach to shoreline dating outlined above. The shoreline date in grey is underlined with the 95% HDR in black. Dates provided by Jakobsson (2014) are plotted in red. The box indicates a 50 year uncertainty range which in combination with the red line extends 200 years. B) Map showing the centroids of the Paurer sites and Sky 1. The sea-level has been simulated using the probability density associated with the shoreline date for Paurer 1 (see also map in Jakobsson 2014:fig.12a). Paurer 1 is the red point.

458 by the close relationship between sites and the shoreline up until the transition to the Neolithic at c. 4000  
459 BCE, after which a couple of sites become situated some distance from the sea, followed by a more decisive  
460 break at the transition to the Late Neolithic at c. 2500 BCE. This development is in clear agreement with  
461 the literature. Furthermore, based on the quantitative nature of these findings, an initial formulation of  
462 a refined method for the shoreline dating of pre-Late Neolithic Stone Age sites has been proposed. Apart  
463 from taking the distance between sites and the isobases of the displacement curves into consideration when  
464 dating the sites, this involves accounting for the distance between the sites and the shoreline. When no  
465 other information is available, it can at present be recommended to use the empirically derived exponential  
466 decay ratio of 0.173 (Figure 11A) to characterise this relationship. Furthermore, while this remains to be  
467 formalised and explored further, it was also showed how the accuracy of the method can be improved by  
468 including more information, both with reference to the topographic location of the sites and other temporal  
469 data. As the precision of the method is both geographically and temporally contingent due to the trajectory  
470 of RSL-change, where older sites situated towards the north-east in the study area will get a more precise  
471 date, the impact of such additional information will also vary.

472 Future investigations and radiocarbon dates from Stone Age sites in the region can not only be used to  
473 further evaluate and adjust the findings reported here, but a larger sample size could also lay the foundations  
474 for refining the method by identifying subsets of sites for which the application of the method could be  
475 adjusted. Given it's behavioural nature, it would for example seem likely that dimensions such as the  
476 nature and purpose of visits to the sites will have implications for how close to the shoreline they were  
477 located. Furthermore, other dimensions related to the topographic location of the sites could be similarly  
478 explored. This for example pertains to the exposure of sites to wave action, which is likely to have been  
479 of concern (Roalkvam 2020), and which presumably has implications for how close to the shoreline people  
480 settled (Blankholm 2020; Helskog 1978). This is also related to the fact that while the mean sea-level is  
481 used for dating the sites, a consideration of the tidal range could possibly also have implications for the site  
482 location relative to the shoreline, depending on the topography (Helskog 1978). The potential of exploring  
483 dimensions such as these was also hinted at here with the estimation and cursory treatment of the horizontal  
484 and topographic distance to the shoreline. If patterns related to such locational patterns can be discerned  
485 and unpicked, this will not least be useful for improving the shoreline dating of sites which have only been  
486 surveyed and where little information beyond their location is available.

487 Some limitations and sources of likely variation and uncertainty that have not been considered should also be  
488 mentioned. First of all the sample size is quite strained and the future addition of more sites might alter the  
489 picture considerably. Secondly, the validity of the method was evaluated by applying it to the data from  
490 where the input parameters were derived. Fitting and evaluating a model using the exact same data will  
491 likely exaggerate its performance. Thirdly, the DTM has only been corrected for major modern disturbances.  
492 This means that other forms of erosion, although likely not that prevalent, has not been taken into account.  
493 Fourthly, the DTM has a vertical error which could also benefit from being integrated in the analysis (cf.  
494 Lewis 2021). Fifthly, the displacement curves were here interpolated to all site locations without accounting  
495 for increased uncertainty as one moves further away from the isobases of the displacement curves. This is  
496 also related to the fact that the RSL data can be handled in different ways than with the isobase method  
497 that has been used for the compilation of the employed displacement curves (cf. Creel et al. 2022). Sixthly,  
498 neither the question of how site limits are defined nor the elevation range over which these extend was given  
499 much consideration (cf. Mjærum 2022). Finally, the radiocarbon dates and division of settlement phases  
500 at each site was here simply done by treating radiocarbon dates not overlapping at 99.7% as representing  
501 unrelated occupation events. This could also be handled differently (e.g. Bronk Ramsey 2009, 2015). While  
502 each of these factors will have variable impact on the final results, they clearly represent dimensions which  
503 would all benefit from further consideration and which means that some of the precision following from the  
504 outlined approach is likely to be spurious.

505 Given that shoreline dating is contingent on regularities in human behaviour it should naturally be applied  
506 with care. Furthermore, formulating and visualising the method along the lines of how radiocarbon dates  
507 are treated, as was done here, does stand the chance of giving a veneer of radiometric accuracy that is  
508 not warranted. That being said, the best chance we have of not throwing away precious temporal data, or  
509 exaggerating our handle on it, is arguably to rigorously evaluate the method using independent data such as

---

510 radiocarbon dates, by offering a precise formulation of how it could be applied, by specifying what sources of  
511 uncertainty are accounted for and by making this process open through the dissemination of underlying data  
512 and programming code.

513 Finally, this analysis employed a simulation approach to integrate multiple sources of spatio-temporal  
514 uncertainty. Here this was simply used to inform the question of the distance between sites and the shoreline.  
515 However, this method and general framework can be extended to a wide range of use-cases where one needs  
516 to visualise, and quantitatively or qualitatively evaluate the relationship between archaeological phenomena,  
517 the prehistoric shoreline, and the uncertainty inherent in this reconstruction.

---

## 518 9 References

- 519 Åkerlund, Agneta  
520 1996 *Human responses to shore displacement: Living by the sea in Eastern Middle Sweden during the Stone*  
521 *Age*. Riksantikvarieämbetet, Stockholm.
- 522 Åkerlund, Agneta, Jan Risberg, Urve Miller, and Per Gustafsson  
523 1995 On the applicability of the  $^{14}\text{C}$  method to interdisciplinary studies on shore displacement and settlement  
524 location. *PACT* 49:53–84.
- 525 Amundsen, Øystein, Stig Knutsen, Axel Mjærum, and Gaute Reitan  
526 2006 Nøkleby i Ski – en tidligeolittisk jordbruksboplass? *Primitive tider* 9:85–96.
- 527
- 528 Åstveit, Leif Inge  
529 2018 The Early Mesolithic of Western Norway. In *Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 231–274.  
530 Equinox, Sheffield.
- 531 Bakka, Egil, and Peter Emil Kaland  
532 1971 Early farming in Hordaland, western Norway. Problems and approaches in archaeology and pollen  
533 analysis. *Norwegian Archaeological Review* 4:1–17. DOI:10.1080/00293652.1971.9965136.
- 534 Bang-Andersen, Sveinung  
535 2012 Colonizing Contrasting Landscapes. The Pioneer Coast Settlement and Inland Utilization in  
536 Southern Norway 10,000–9500 Years Before Present. *Oxford Journal of Archaeology* 31:103–120.  
DOI:10.1111/j.1468-0092.2012.00381.x.
- 537 Berg-Hansen, Inger Marie  
538 2009 *Steinalderregistrering. Metodologi og forskningshistorie i Norge 1900–2000 med en feltstudie fra Lista i*  
539 *Vest-Agder*. Museum of Cultural History, University of Oslo, Oslo.
- 540 2018 Continuity and Change in Late- and Post-glacial Social Networks: Knowledge Transmission and  
541 Blade Production Methods in Ahrensburgian and Early Mesolithic North West Europe. In *The Early*  
*Settlement of Northern Europe. Transmission of Knowledge and Culture*, edited by Kjel Knutsson,  
Helena Knutsson, Jan Apel, and Håkon Glørstad, pp. 63–98. Equinox, Sheffield.
- 542 Bergsvik, Knut Andreas  
543 2009 Caught in the middle: functional and ideological aspects of Mesolithic shores in Norway. In *Mesolithic*  
*Horizons: Papers presented at the Seventh International Conference on the Mesolithic in Europe, Belfast 2005*, edited by Sinéad B. McCartan, Rick Schulting, Graeme Warren, and Peter Woodman,  
pp. 602–609. Oxbow Books, Oxford.
- 544 Bevan, Andrew, Enrico R. Crema, Xiuzhen Li, and Alessio Palmisano  
545 2013 Intensities, Interactions, and Uncertainties: Some New Approaches to Archaeological Distributions.  
In *Computational Approaches to Archaeological Spaces*, edited by Andrew Bevan and Mark Lake, pp.  
546 27–52. Left Coast Press, Walnut Creek.
- 547 Bivand, Roger  
548 2021 *rgrass7: Interface Between GRASS 7 Geographical Information System and R*. R package version  
0.2-6.
- 549 Bjerck, Hein Bjartmann  
550 1990 Mesolithic site types and settlement patterns at Vega, Northern Norway. *Acta Archaeologica* 60:1–32.
- 551 2005 Strandlinjedatering. In *Norsk arkeologisk leksikon*, edited by Einar Østmo and Lotte Hedeager, pp.  
363–364. Pax, Oslo.
- 552 2008a Norwegian Mesolithic Trends: A Review. In *Mesolithic Europe*, edited by Geoff Bailey and Penny  
Spikins, pp. 60–106. Cambridge University Press, Cambridge.
- 553 2008b Innledende betraktninger. In *NTNU Vitenskapsmuseets arkeologiske undersøkelser Ormen Lange*  
Nyhamna, edited by Hein Bjartmann Bjerck, Leif Inge Åstveit, Trond Meling, Jostein Gundersen,  
Guro Jørgensen, and Staale Normann, pp. 548–551. Tapir Akademisk Forlag, Trondheim.

- 560 2017 Settlements and Seafaring: Reflections on the Integration of Boats and Settlements Among Marine  
561 Foragers in Early Mesolithic Norway and the Yámana of Tierra del Fuego. *The Journal of Island and  
Coastal Archaeology* 12(2):276–299. DOI:10.1080/15564894.2016.1190425.
- 562 Blankholm, Hans Peter  
563 2020 In the wake of the wake. An investigation of the impact of the Storegga tsunami on the human  
settlement of inner Varangerfjord, northern Norway. *Quaternary International* 549:65–73. DOI:<https://doi.org/10.1016/j.quaint.2018.05.050>.
- 564 Breivik, Heidi Mjelva  
565 2014 Palaeo-oceanographic development and human adaptive strategies in the Pleistocene–Holocene transition: A study from the Norwegian coast. *The Holocene* 24:1478–1490. DOI:10.1177/0959683614544061.
- 566 Breivik, Heidi Mjelva, Guro Fossum, and Steinar Solheim  
567 2018 Exploring human responses to climatic fluctuations and environmental diversity: Two stories  
from Mesolithic Norway. *Quaternary International* 465. Impacts of gradual and abrupt environmental  
changes on Late glacial to Middle Holocene cultural changes in Europe:258–275.  
DOI:10.1016/j.quaint.2016.12.019.
- 568 Breivik, Heidi, and Hein Bjartmann Bjerck  
569 2018 Early Mesolithic Central Norway: A Review of Research History, Settlements, and Tool Tradition. In  
*Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change*,  
570 edited by Hans Peter Blankholm, pp. 169–206. Equinox, Sheffield.
- 571 Brøgger, Waldemar Christofer  
572 1905 *Strandliniens Beliggenhed under Stenalderen i Det Sydøstlige Norge*. Norges geologiske undersøkelse,  
573 Kristiania.
- 574 Bronk Ramsey, Christopher  
575 2009 Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* 51(1):337–360.  
DOI:10.1017/S0033822200033865.
- 576 2015 Bayesian Approaches to the Building of Archaeological Chronologies. In *Mathematics and Archaeology*,  
577 edited by Juan A. Barcelo and Igor Bogdanovic, pp. 272–292. CRC Press, Boca Raton.
- 578 Conolly, James  
579 2020 Spatial interpolation. In *Archaeological Spatial Analysis: A Methodological Guide*, edited by Mark  
Gillings, Piraye Hacigüzeller, and Gary Lock, pp. 118–134. Routledge, London & New York.
- 580 Conolly, James, and Mark Lake  
581 2006 *Geographical Information Systems in Archaeology*. Cambridge University Press, Cambridge.
- 582 Creel, Roger C., Jacqueline Austermann, Nicole S. Khan, William J. D’Andrea, Nicholas Balascio, Blake  
583 Dyer, Erica Ashe, and William Menke  
584 2022 Postglacial relative sea level change in Norway. *Quaternary Science Reviews* 282:107422.  
DOI:10.1016/j.quascirev.2022.107422.
- 585 Crema, Enrico R.  
586 2012 Modelling Temporal Uncertainty in Archaeological Analysis. *Journal of Archaeological Method and  
Theory* 19(3):440–461. DOI:10.1007/s10816-011-9122-3.
- 587 2015 Time and Probabilistic Reasoning in Settlement Analysis. In *Mathematics and Archaeology*, edited by  
Juan A. Barcelo and Igor Bogdanovic, pp. 314–334. CRC Press, Boca Raton.
- 588 Crema, Enrico R., Andrew Bevan, and Mark W. Lake  
589 2010 A probabilistic framework for assessing spatio-temporal point patterns in the archaeological record.  
*Journal of Archaeological Science* 37(5):1118–1130. DOI:10.1016/j.jas.2009.12.012.
- 590 Damlien, Hege, and Steinar Solheim  
591 2018 The Pioneer Settlement of Eastern Norway. In *Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 335–367.  
Equinox, Sheffield.
- 592 De Geer, Gerard

- 604 1896 *Om Skandinaviens geografiska utveckling efter Istiden*. P. A. Norstedt & Söner, Stockholm.
- 605
- 606 Eskeland, Knut Fossdal
- 607 2017 *Rapport, arkeologisk registrering. E18 Langangen Rugtvedt, 16/06999, Porsgrunn og Bamble kommune.*
- 608 Skien.
- 609 Fossum, Guro
- 610 2020 Specialists facing climate change. The 8200 cal BP event and its impact on the coastal settlement in the inner Oslo fjord, southeast Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement with the Coast from the Atlantic to the Baltic Sea*, edited by Almut Schülke, pp. 179–201. Routledge, London & New York.
- 611
- 612 Fuglestvedt, Ingrid
- 613 2012 The Pioneer Condition on the Scandinavian Peninsula: the Last Frontier of a ‘Palaeolithic Way’ in Europe. *Norwegian Archaeological Review* 45(1):1–29. DOI:10.1080/00293652.2012.669998.
- 614
- 615 Gjerpe, Lars Erik, and Grethe Bjørkan Bukkemoen
- 616 2008 Nordby 1 – Toskipede hus fra neolitikum-bronsealder og boplasspor fra jernalder. In *E18-prosjektet Vestfold. Bind 3. Hus, boplass- og dyrkningspor*, edited by Lars Erik Gjerpe, pp. 7–38. University of Oslo, Museum of Cultural History, Oslo.
- 617
- 618 Glørstad, Håkon (editor)
- 619 2002 *Svinesundprosjektet. Bind 1. Utgravninger avsluttet i 2001*. University of Oslo, Museum of Cultural History, Oslo.
- 620
- 621 (editor)
- 622 2003 *Svinesundprosjektet . Bind 2. Utgravninger avsluttet i 2002*. University of Oslo, Museum of Cultural History, Oslo.
- 623
- 624 (editor)
- 625 2004 *Svinesundprosjektet. Bind 3. Utgravninger avsluttet i 2003*. University of Oslo, Museum of Cultural History, Oslo.
- 626
- 627 2010 *The Structure and History of the Late Mesolithic Societies in the Oslo Fjord Area 6300-3800 BC*. Bricoleur Press, Lindome.
- 628
- 629 2012 Historical ideal types and the transition to the Late Neolithic in South Norway. In *Becoming European. The transformation of third millennium Northern and Western Europe*, edited by Christopher Prescott and Håkon Glørstad, pp. 82–99. Oxbow Books, Oxford & Oakville.
- 630
- 631 2016 Deglaciation, sea-level change and the Holocene colonization of Norway. *Geological Society, London, Special Publications* 411:9–25. DOI:10.1144/SP411.7.
- 632
- 633 Glørstad, Håkon, Jostein Gundersen, Frode Kvalø, Pål Nymoen, David Simpson, and Birgitte Skar
- 634 2020 Submerged Stone Age from a Norwegian Perspective. In *The Archaeology of Europe’s Drowned Landscapes*, edited by Geoff Bailey, Nena Galanidou, Hans Peeters, Hauke Jöns, and Moritz Mennenga, pp. 125–140. Springer, Cham.
- 635
- 636 GRASS Development Team
- 637 2017 *Geographic Resources Analysis Support System (GRASS) Software, Version 7.2*. Open Source Geospatial Foundation.
- 638
- 639 Gundersen, Jostein
- 640 2013 Verken fjord eller fjell – steinalderen i det kystnære innlandet. Gamle og nye funn fra Notodden i Telemark. *Viking* 76:35–62.
- 641
- 642 Hafsten, Ulf
- 643 1957 De senkvartære strandlinje-forskyvningene i Oslotrakten belyst ved pollenanalytiske undersøkelser. *Norwegian Journal of Geography* 16(1-8):74–99. DOI:10.1080/00291955708622137.
- 644
- 645 1983 Shore-level changes in South Norway during the last 13,000 years, traced by biostratigraphical methods and radiometric datings. *Norwegian Journal of Geography* 37(2):63–79. DOI:10.1080/00291958308552089.
- 646
- 647 Hagen, Anders

- 648 1963 Problemkompleks Fosna. Opphav – kontakt med kontinentale grupper – forholdet til Komsa. In *Boplatsproblem vid Kattegat och Skagerack*, pp. 53–59. Göteborg och Bohusläns fornminnesförening &  
649 Institutionen för nordisk fornkunskap, Gothenburg University, Gothenburg.
- 650 Helskog, Knut  
651 1978 Late Holocene sea-level changes seen from prehistoric settlements. *Norwegian Journal of Geography*  
652 32:111–119. DOI:10.1080/00291957808552032.
- 653 Herzog, Irmela  
654 2013 The Potential and Limits of Optimal Path Analysis. In *Computational Approaches to Archaeological*  
655 *Spaces*, edited by Andrew Bevan and Mark Lake, pp. 179–211. Left Coast Press, Walnut Creek.
- 656 Hinsch, Erik  
657 1955 Traktbegerkultur – Megalitkultur. En studie av Øst-Norges eldste neolitiske gruppe. *Universitetets*  
658 *Oldsaksamling Årbok* 1951–1953:10–177.
- 659 Hinz, Martin, Clemens Schmid, Daniel Knitter, and Carolin Tietze  
660 2021 *oxcAAR: Interface to 'OxCal' radiocarbon calibration. R package version 1.1.0.*
- 661 Hollender, Artur  
662 1901 Om sveriges nivåförändringar efter människans invandring. *Geologiska Föreningen i Stockholm Förhan-*  
663 *dlingar* 23(4):1118–1130. DOI:10.1080/00293652.1975.9965220.
- 664 Hughes, Anna L. C., Richard Gyllencreutz, Øystein S. Lohne, Jan Mangerud, and John Inge Svendsen  
665 2016 The last Eurasian ice sheets – a chronological database and time-slice reconstruction, DATED-1.  
666 *Boreas* 45(1):1–45. DOI:https://doi.org/10.1111/bor.12142.
- 667 Hyndman, Rob J  
668 1996 Computing and Graphing Highest Density Regions. *The American Statistician* 50(2):120–126.
- 669 Ilves, Kristin, and Kim Darmark  
670 2011 Some Critical and Methodological Aspects of Shoreline Determination: Examples from the Baltic Sea  
671 Region. *Journal of Archaeological Method and Theory* 18:147–165. DOI:10.1007/s10816-010-9084-x.
- 672 Jakslund, Lasse (editor)  
673 2001 *Vinterbrolokalitetene - En kronologisk sekvens fra mellom- og senmesolitikum i Ås, Akershus*. University  
674 of Oslo, Museum of Cultural History, Oslo.  
675 (editor)  
676 2012a *E18 Brunlanesprosjektet. Bind II. Undersøkte lokaliteter fra tidligmesolitikum*. University of Oslo,  
677 Museum of Cultural History, Oslo.  
678 (editor)  
679 2012b *E18 Brunlanesprosjektet. Bind III. Undersøkte lokaliteter fra tidligmesolitikum og senere*. University  
680 of Oslo, Museum of Cultural History, Oslo.  
681 2014 Kulturhistorisk sammenstilling. In *E18 brunlanesprosjektet. Bind i. Forutsetninger og kulturhistorisk*  
682 *sammenstilling*, edited by Lasse Jakslund and Per Persson, pp. 11–46. University of Oslo, Museum of  
683 Cultural History, Oslo.  
684 Jakslund, Lasse, and Per Persson (editors)  
685 2014 *E18 Brunlanesprosjektet. Bind I. Forutsetninger og kulturhistorisk sammenstilling*. University of Oslo,  
686 Museum of Cultural History, Oslo.  
687 Johansen, Erling  
688 1963 Kyst(fangst)boplassenes strandbundenhet og strandlinjekronologien. In *Boplatsproblem vid Kattegat*  
689 *och Skagerack*, pp. 90–92. Göteborg och Bohusläns fornminnesförening & Institutionen för nordisk  
690 fornkunskap, Gothenburg University, Gothenburg.
- 691 Jørgensen, Erlend Kirkeng, Petro Pesonen, and Miikka Tallavaara  
692 2020 Climatic changes cause synchronous population dynamics and adaptive strategies among coastal  
693 hunter-gatherers in Holocene northern Europe. *Quaternary Research*:1–16. DOI:10.1017/qua.2019.86.
- 694 Kjemperud, Alfred

- 695 1986 Late Weichselian and Holocene shoreline displacement in the Trondheimsfjord area, central Norway.  
696 *Boreas* 15(1):61–82. DOI:10.1111/j.1502-3885.1986.tb00744.x.
- 697 Kleppe, Else Johansen
- 698 1985 *Archaeological Data on Shore Displacements in Norway*. Norges geografiske oppmåling, Hønefoss.
- 699
- 700 Kleppe, Jan Ingolf
- 701 2018 The Pioneer Colonization of Northern Norway. In *Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 13–57. Equinox, Sheffield.
- 702 Lakens, Daniël, Anne M. Scheel, and Peder M. Isager
- 704 2018 Equivalence Testing for Psychological Research: A Tutorial. *Advances in Methods and Practices in Psychological Science* 1(2):259–269. DOI:10.1177/2515245918770963.
- 705
- 706 Lewis, Joseph
- 707 2021 Probabilistic Modelling for Incorporating Uncertainty in Least Cost Path Results: a Postdictive Roman Road Case Study. *Journal of Archaeological Method and Theory* 28(3):911–924. DOI:10.1007/s10816-021-09522-w.
- 708
- 709 Marwick, Ben, Carl Boettiger, and Lincoln Mullen
- 710 2018 Packaging Data Analytical Work Reproducibly Using R (and Friends). *The American Statistician* 72(1):80–88. DOI:10.1080/00031305.2017.1375986.
- 711
- 712 Melvold, Stine, and Per Persson (editors)
- 713 2014 *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og Porsgrunn. Bind 1. Tidlig- Og mellommesolittiske lokaliteter i Vestfold og Telemark*. Portal forlag, Kristiansand.
- 714
- 715 Mikkelsen, Egil
- 716 1975 Mesolithic in South-Eastern Norway. *Norwegian Archaeological Review* 8(1):1118–1130.  
717 DOI:10.1080/11035890109445866.
- 718 Milne, Glenn A
- 719 2015 Glacial isostatic adjustment. In *Handbook of sea-level research*, edited by Ian Shennan, Antony J Long, and Benjamin P Horton, pp. 421–437. Wiley, Chichester.
- 720
- 721 Milne, Glenn A., W. Roland Gehrels, Chris W. Hughes, and Mark E. Tamisiea
- 722 2009 Identifying the causes of sea-level change. *Nature Geoscience* 2(7):471–478. DOI:10.1038/ngeo544.
- 723
- 724 Mjærum, Axel
- 725 2018 Hinterland discoveries: Middle Mesolithic woodland utilization and the case of the site Eidsberg, eastern Norway. *Current Swedish Archaeology* 26(1):159–188. DOI:10.37718/CSA.2018.11.
- 726
- 727 2022 A Matter of Scale: Responses to Landscape Changes in the Oslo Fjord, Norway, in the Mesolithic. *Open Archaeology* 8(1):62–84. DOI:10.1515/opar-2022-0225.
- 728
- 729 Møller, Jakob J
- 730 1987 Shoreline relation and prehistoric settlement in northern norway. *Norwegian Journal of Geography* 41:45–60. DOI:<http://dx.doi.org/10.1080/00291958708552171>.
- 731
- 732 Mörner, Nils-Axel
- 733 1976 Eustasy and Geoid Changes. *The Journal of Geology* 84(2):123–151. DOI:10.1086/628184.
- 734
- 735 1979 The Fennoscandian Uplift and Late Cenozoic Geodynamics: Geological Evidence. *GeoJournal* 3(3):287–318. DOI:10.1007/BF00177634.
- 736
- 737 Nielsen, Svein Vatsvåg
- 738 2021 Early farming in Southeastern Norway: New evidence and interpretations. *Journal of Neolithic Archaeology* 23:83–113. DOI:10.12766/jna.2021.4.
- 739
- 740 Nielsen, Svein Vatsvåg, Per Persson, and Steinar Solheim
- 741 2019 De-Neolithisation in southern Norway inferred from statistical modelling of radiocarbon dates. *Journal of Anthropological Archaeology* 53:82–91. DOI:10.1016/j.jaa.2018.11.004.

- 742
- 743 Nordqvist, Bengt
- 744 1995 The Mesolithic settlement of the west coast of Sweden - with special emphasis on chronology and topography of coastal settlements. In *Man and Sea in the Mesolithic. Coastal settlement above and below present sea level*, edited by Anders Fischer, pp. 185–196. Oxbow Books, Oxford.
- 745
- 746 1999 The Chronology of the Western Swedish Mesolithic and Late Paleolithic: Old Answers in Spite of New Methods. In *The Mesolithic of Central Scandinavia*, edited by Joel Boaz, pp. 235–253. University of Oslo, Oslo.
- 747
- 748 Norwegian Mapping Authority
- 749 2018 *Produktspesifikasjon. Nasjonal modell for høydedata fra laserskanning (FKB-laser). FKB-laser\_v30.*
- 750
- 751 2021 *Tidevannstabeller for den norske kyst med Svalbard samt Dover, England.*
- 752
- 753 Nummedal, Anders
- 754 1923 Om flintpladsene. *Norwegian Journal of Geography* 7(2):89–141.
- 755
- 756 Østmo, Einar
- 757 1988 *Etableringen av jordbrukskultur i Østfold i steinalderen.* The University Collection of National Antiquities, University of Oslo, Oslo.
- 758
- 759 2008 *Auve. En fangstboplass fra yngre steinalder på Vesterøya i Sandefjord. I. Den arkeologiske del.* Museum of Cultural History, University of Oslo, Oslo.
- 760
- 761 Parnell, A. C., J. Haslett, J. R. M. Allen, C. E. Buck, and B. Huntley
- 762 2008 A flexible approach to assessing synchronicity of past events using Bayesian reconstructions of sedimentation history. *Quaternary Science Reviews* 27(19–20):1872–1885. DOI:10.1016/j.quascirev.2008.07.009.
- 763
- 764 Persson, Per
- 765 2008 Nauen 5.2 – Stenåldersboplatter och fossil åkermark. In *E18-prosjektet Vestfold. Bind 2. Steinalderboplasser, boplasspor, graver og dyrkningsspor*, edited by Lars Erik Gjerpe, pp. 163–198. University of Oslo, Museum of Cultural History, Oslo.
- 766
- 767 Prescott, Christopher
- 768 2020 Interpreting Complex Diachronic "Neolithic"-Period Data in Norway. In *Farmers at the Frontier – A Pan European Perspective on Neolithisation*, edited by Kurt J. Gron, Lasse Sørensen, and Peter Rowley-Conwy, pp. 381–400. Oxbow Books, Oxford.
- 769
- 770 R Core Team
- 771 2021 *R: A language and environment for statistical computing.* R Foundation for Statistical Computing, Vienna, Austria.
- 772
- 773 Ramstad, Morten
- 774 2009 Eldre steinalder på Melkøya, representativitet, strandlinjer og transgresjon. In *Undersøkelsene på Melkøya. Melkøyaprosjektet – Kulturhistoriske registreringer og utgravninger 2001 og 2002*, edited by Anders Hesjedal, Morten Ramstad, and Anja R. Niemi, pp. 491–495. Tromsø museum, Universitetsmuseet, Tromsø.
- 775
- 776 Reimer, Paula J., William E. N. Austin, Edouard Bard, Alex Bayliss, Paul G. Blackwell, Christopher Bronk Ramsey, Martin Butzin, Hai Cheng, R. Lawrence Edwards, Michael Friedrich, Pieter M. Grootes, Thomas P. Guilderson, Irka Hajdas, Timothy J. Heaton, Alan G. Hogg, Konrad A. Hughen, Bernd Kromer, Sturt W. Manning, Raimund Muscheler, Jonathan G. Palmer, Charlotte Pearson, Johannes van der Plicht, Ron W. Reimer, David A. Richards, E. Marian Scott, John R. Southon, Christian S. M. Turney, Lukas Wacker, Florian Adolphi, Ulf Büntgen, Manuela Capano, Simon M. Fahrni, Alexandra Fogtmann-Schulz, Ronny Friedrich, Peter Köhler, Sabrina Kudsk, Fusa Miyake, Jesper Olsen, Frederick Reinig, Minoru Sakamoto, Adam Sookdeo, and Sahra Talamo
- 784 2020 The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* 62(4):725–757. DOI:10.1017/RDC.2020.41.
- 785
- 786 Reitan, Gaute, and Inger Marie Berg-Hansen

- 787 2009 *Lundevågenprosjektet, delrapport 1. Sammenfattende rapport. Lunde, 6/1, 6/35 og Skjolnes 7/23, 7/27, Farsund kommune, Vest-Agder*. Oslo.
- 788
- 789 Reitan, Gaute, and Per Persson (editors)
- 790 2014 *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og Porsgrunn. Bind 2. Seinmesolittiske, neolittiske og yngre lokaliteter i Vestfold og Telemark*. Portal forlag, Kristiansand.
- 791
- 792 Reitan, Gaute, and Lars Sundström (editors)
- 793 2018 *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*. Cappelen Damm Akademisk, Oslo.
- 794
- 795 Roalkvam, Isak
- 796 2020 Algorithmic classification and statistical modelling of coastal settlement patterns in Mesolithic south-eastern Norway. *Journal of Computer Applications in Archaeology* 3(1):288–307. DOI:<https://doi.org/10.5334/jcaa.60>.
- 797
- 798 2022 Exploring the composition of lithic assemblages in Mesolithic south-eastern Norway. *Journal of Archaeological Science: Reports* 42:103371. DOI:10.1016/j.jasrep.2022.103371.
- 799
- 800 Røberg, Frank Halvar N.
- 801 2012 *Bosettings- og aktivitetsspor. Larønningen, 221/2138. Skien, Telemark*. University of Oslo, Museum of Cultural History, Oslo.
- 802
- 803 Romundset, Anders
- 804 2018 Postglacial shoreline displacement in the Tvedstrand-Arendal area. In *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*, edited by Gaute Reitan and Lars Sundström, pp. 463–478. Cappelen Damm Akademisk, Oslo.
- 805
- 806 Romundset, Anders, Stein Bondevik, and Ole Bennike
- 807 2011 Postglacial uplift and relative sea level changes in Finnmark, northern Norway. *Quaternary Science Reviews* 30(19-20):2398–2421. DOI:10.1016/j.quascirev.2011.06.007.
- 808
- 809 Romundset, Anders, Ola Fredin, and Fredrik Høgaas
- 810 2015 A Holocene sea-level curve and revised isobase map based on isolation basins from near the southern tip of Norway. *Boreas* 44:383–400. DOI:10.1111/bor.12105.
- 811
- 812 Romundset, Anders, Thomas R. Lakeman, and Fredrik Høgaas
- 813 2018 Quantifying variable rates of postglacial relative sea level fall from a cluster of 24 isolation basins in southern Norway. *Quaternary Science Reviews* 197:175–192. DOI:10.1016/j.quascirev.2018.07.041.
- 814
- 815 2019 Coastal lake records add constraints to the age and magnitude of the Younger Dryas ice-front oscillation along the Skagerrak coastline in southern Norway. *Journal of Quaternary Science* 34(2):112–124. DOI:<https://doi.org/10.1002/jqs.3085>.
- 816
- 817 Schmitt, Lou, Stephan Larsson, Jan Burdukiewicz, John Ziker, Krister Svedhage, Jeanette Zamon, and Steffen Holger
- 818
- 819 2009 Chronological Insights, Cultural Change, and Resource Exploitation on the West Coast of Sweden During the Late Palaeolithic/Early Mesolithic Transition. *Oxford Journal of Archaeology* 28:1–27. DOI:10.1111/j.1468-0092.2008.00317.x.
- 820
- 821 Schülke, Almut
- 822 2020 First visit or revisit? Motivations of mobility and the use and reuse of sites in the changing coastal areas of Mesolithic southeastern Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement with the Coast from the Atlantic to the Baltic Sea*, edited by Almut Schülke, pp. 359–393. Routledge, London & New York.
- 823
- 824 Shennan, Ian
- 825 2015 Handbook of sea-level research: Framing research questions. In *Handbook of Sea-Level Research*, edited by Ian Shennan, Antony J Long, and Benjamin P Horton, pp. 3–25. Wiley, Chichester.
- 826
- 827 Shetelig, Haakon
- 828 1922 *Primitive Tider i Norge – En oversikt over stenalderen*. John Griegs Forlag, Bergen.
- 829

- 830 Sognnes, Kalle  
831 2003 On shoreline dating of rock art. *Acta Archaeologica* 74:189–209.
- 832
- 833 Solheim, Steinar  
834 2012 Lokal praksis og fremmed opphav. Arbeidsdeling, sosiale relasjoner og differensiering i østnorsk tidlige neolitikum. Unpublished PhD thesis, Oslo.  
835
- 836 (editor)  
837 2017 *E18 Røgtvedt-Døradal. Arkeologiske undersøkelser av lokaliteter fra steinalder og jernalder i Bamble kommune, Telemark fylke*. Portal forlag, Kristiansand.  
838
- 839 2020 Mesolithic coastal landscapes. Demography, settlement patterns and subsistence economy in southeastern Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement with the Coast from the Atlantic to the Baltic Sea*, edited by Almut Schülke. Routledge, London & New York.  
840
- 841 2021 Timing the Emergence and Development of Arable Farming in Southeastern Norway by Using Summed Probability Distribution of Radiocarbon Dates and a Bayesian Age Model. *Radiocarbon* 63(5):1–22. DOI:10.1017/RDC.2021.80.  
842
- 843 Solheim, Steinar, and Per Persson  
844 2018 Early and mid-Holocene coastal settlement and demography in southeastern Norway: Comparing distribution of radiocarbon dates and shoreline-dated sites, 8500–2000 cal. BCE. *Journal of Archaeological Science: Reports* 19:334–343. DOI:10.1016/j.jasrep.2018.03.007.  
845
- 846 Sørensen, Rolf  
847 1979 Late Weichselian deglaciation in the Oslofjord area, south Norway. *Boreas* 8(2):241–246. DOI:<https://doi.org/10.1111/j.1502-3885.1979.tb00806.x>.  
848
- 849 1999 En <sup>14</sup>C datert og dendrokronologisk kalibrert strandforksyvningskurve for sørøst-Norge. In *Museumslandskap. Artikkelsamling til Kerstin Griffin på 60-årsdagen. Bind A*, edited by Lotte Selsing and Grete Lillehammer, pp. 227–242. AmS-rapport 12A. Museum of Archaeology, Stavanger.  
850
- 851 Sørensen, Rolf, Kari E. Henningsmoen, Helge I. Høeg, and Veronika Gälman  
852 2014 Holocene landhevningsstudier i sørøst-Vestfold og sørøstre Telemark – Revidert kurve. In *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og Porsgrunn. Bind 1*, edited by Stine Melvold and Per Persson, pp. 36–47. Portal, Kristiansand.  
853
- 854 Sørensen, Rolf, Kari E. Henningsmoen, Helge I. Høeg, and Veronika Gälman  
855 in prep Holocen vegetasjonshistorie og landhevning i sørøst-Vestfold og sørøstre Telemark. In *The Stone Age in Telemark. Archaeological results and scientific analysis from Vestfoldbaneprosjektet and E18 Røgtvedt-Døradal*, edited by Per Persson and Steinar Solheim.  
856
- 857 Sørensen, Rolf, Helge I. Høeg, Kari E. Henningsmoen, Göran Skog, Solveig F. Labowsky, and Bjørg Stabell  
858 2014 Utviklingen av det senglasiale og tidlig preboreale landskapet og vegetasjonen omkring steinalderboplassene ved Pauler. In *E18 Brunlanesprosjektet. Bind I. Forutsetninger og kulturhistorisk sammenstilling*, edited by Lasse Jaksland and Per Persson, pp. 171–213. University of Oslo, Museum of Cultural History, Oslo.  
859
- 860 Stokke, Jo-Simon Frøshaug, and Gaute Reitan  
861 2018 Kvastad A2. Lokalitet med funn fra tidlig- og mellommesolitikum og dyrkningsspor fra mellom- og senneolitikum. In *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*, edited by Gaute Reitan and Lars Sundström, pp. 375–407. Cappelen Damm Akademisk, Oslo.  
862
- 863 Stroeven, Arjen P., Clas Hättestrand, Johan Kleman, Jakob Heyman, Derek Fabel, Ola Fredin, Bradley W. Goodfellow, Jonathan M. Harbor, John D. Jansen, Lars Olsen, Marc W. Caffee, David Fink, Jan Lundqvist, Gunhild C. Rosqvist, Bo Strömborg, and Krister N. Jansson  
864
- 865 2016 Deglaciation of Fennoscandia. *Quaternary Science Reviews* 147:91–121. DOI:10.1016/j.quascirev.2015.09.016.  
866
- 867 Svendsen, John Inge, and Jan Mangerud  
868 1987 Late Weichselian and Holocene sea-level history for a cross-section of western Norway. *Journal of Quaternary Science* 2(2):113–132. DOI:10.1002/jqs.3390020205.  
869
- 870

- 
- 871 Tallavaara, Miikka, and Petro Pesonen  
872 2020 Human ecodynamics in the north-west coast of Finland 10,000–2000 years ago. *Quaternary Interna-*  
873 *tional* 549:26–35. DOI:10.1016/j.quaint.2018.06.032.
- 874 Wang, Ian  
875 2019 *topoDistance: Calculating topographic paths and distances. R package version 1.0.1. Https://CRAN.r-*  
876 *project.org/package=topoDistance.*
- 877 Wenn, Camilla Cecilie  
878 2012 *Bosetningsspor, produksjonsområde og dyrkningsspor fra Neolitikum til Folkevandringstid. Bratsberg,*  
879 *63/69, 244. Skien kommune, Telemark.* University of Oslo, Museum of Cultural History, Oslo.
- 880 Wikell, Roger, Fredrik Molin, and Mattias Pettersson  
881 2009 The Archipelago of Eastern Middle Sweden - Mesolithic Settlement in Comparison with  $^{14}\text{C}$  and  
Shoreline Dating. In *Chronology and Evolution within the Mesolithic of North-West Europe*, edited by Philippe Crombé, Mark van Strydonck, Joris Sergant, Mathieu Boudin, and Machteld Bats, pp.  
417–434. Cambridge Scholar Publishing, Brussels.
- 882 Yubero-Gómez, María, Xavier Rubio-Campillo, and Javier López-Cachero  
883 2016 The study of spatiotemporal patterns integrating temporal uncertainty in late prehistoric settlements  
in northeastern Spain. *Archaeological and Anthropological Sciences* 8(3):477–490. DOI:10.1007/s12520-  
884 015-0231-x.
- 885