
¹ A simulation-based assessment of the relation between Stone Age
² sites and relative sea-level change along the Norwegian Skagerrak
³ coast

⁴ Isak Roalkvam^{1,*}

⁵ 24 March, 2022

⁶ ¹ University of Oslo, Institute of Archaeology, Conservation and History

⁷ * Correspondence: Isak Roalkvam <isak.roalkvam@iakh.uio.no>

⁸ **1 Introduction**

⁹ The post-glacial relative sea-level fall that characterises large areas of Fennoscandia is fundamental to
¹⁰ its archaeology. This follows not only from the dramatic changes to the landscape that this process has
¹¹ represented throughout prehistory, but also from the fact that if archaeological phenomena were situated
¹² close to the contemporary shoreline when they were in use, a reconstruction of the trajectory of shoreline
¹³ displacement can be used to date these phenomena based on their altitude relative to the present day sea-level.
¹⁴ This method, also called shoreline dating, has long history of use in the region and is frequently applied
¹⁵ to assign an approximate date to diverse archaeological phenomena such as rock art, grave cairns, various
¹⁶ harbour and sea-side constructions and, as is the focus of this study, Stone Age sites (e.g. Åkerlund 1996;
¹⁷ Bjerck 2005; Nordqvist 1995; Schmitt et al. 2009; Sognnes 2003; Tallavaara and Pesonen 2020; Wikell et al.
¹⁸ 2009).

¹⁹ The close association between Stone Age settlements in the northern parts of Scandinavia and shifting
²⁰ prehistoric shorelines was established by the end of the 19th century (De Geer 1896), and was first applied
²¹ as a dating method at the turn of the century (Brøgger 1905; Hollender 1901). Shoreline dating has been
²² fundamental to Norwegian Stone Age archaeology ever since (e.g. Berg-Hansen 2009; Bjerck 1990, 2008a;
²³ Breivik 2014; Johansen 1963; Mikkelsen 1975; Nummedal 1923; Shetelig 1922; Solheim and Persson 2018).
²⁴ The method is used both independently, and to compliment other sources of temporal data such as typological
²⁵ indicators or radiometric dates. However, given the coarse and fuzzy resolution of established typological
²⁶ frameworks, the vast amount of surveyed sites that only contain generic lithic debitage that could hail
²⁷ from any part of the period, and as the conditions for the preservation of organic material is typically poor
²⁸ in Norway, dating with reference to shoreline displacement is often the only and most precise method by
²⁹ which one can hope to date the sites. Shoreline dating is consequently fundamental to our understanding
³⁰ of the Norwegian Stone Age. This is both because it is central to the temporal framework on which our
³¹ understanding of the period is based, but also because the method is only applicable so long as the societies
³² in question have continuously settled on or close to the contemporary shoreline. Consequently, adherence or
³³ deviation from this pattern also has major implications for the socio-economic foundations of the societies in
³⁴ question.

³⁵ Despite its important role for Norwegian Stone Age archaeology, the applicability of dating by reference to
³⁶ shoreline displacement has only been evaluated using relatively coarse methods. The aim of this paper is to
³⁷ provide a systematic and comprehensive review of the degree to which radiocarbon dates correspond with the
³⁸ dates informed by our current knowledge of shoreline displacement in a larger area of south-eastern Norway,
³⁹ using a more refined methodological approach. The goal is to quantify the degree to which the assumption

40 of shore-bound settlement holds through the Stone Age, and in turn have this inform a refined method for
41 shoreline dating. As presented in more detail below, this problem involves the combined evaluation of three
42 major analytical dimensions. One is the questions of when the sites were in use, the second pertains to the
43 reconstruction of the contemporaneous sea-level, and the third follows from the fact that the relation between
44 site and shoreline is inherently spatial. Taking inspiration from studies that have integrated various sources
45 of spatio-temporal uncertainty through Monte Carlo simulation (e.g. Bevan et al. 2013; Crema et al. 2010;
46 Crema 2012, 2015; Yubero-Gómez et al. 2016), a similar approach is adopted here.

47 2 Background

48 Relative sea-level (RSL) can be defined as the mean elevation of the surface of the sea relative to land, or,
49 more formally, the difference in elevation between the geoid and the surface of the Earth as measured from
50 the Earth's centre (Shennan 2015). Variation in this relative distance follow from a range of effects (e.g.
51 Milne et al. 2009). Of central importance here is eustasy and istostasy. The eustatic sea-level is understood
52 as the sea-level if the water has been evenly distributed across the Earth's surface without adjusting for
53 variation in the rigidity of the Earth, its rotation, or the self-gravitation inherent to the water body itself
54 (Shennan 2015). The eustatic sea-level is mainly impacted by glaciation and de-glaciation, which can bind
55 or release large amounts of water into the oceans (Mörner 1976). Istostasy, on the other hand, pertains to
56 adjustments in the crust to regain gravitational equilibrium relative to the underlying viscous mantle. This is
57 often the result of glacial istostasy, which follows from glaciation and de-glaciation and corresponding loading
58 and unloading of weight, as well as from erosion of the crust, which causes its weight to be redistributed.
59 These effects thus causes the lithosphere to either subside due to increased weight, or to rebound and lift
60 upwards due to lower weight (Milne 2015).

61 Following the end of the Weichselian and the final retreat of the Fennoscandian Ice Sheet (e.g. Hughes
62 et al. 2016; Stroeven et al. 2016, see Figure 1), the isostatic rebound has been so severe that most areas
63 of Norway have been subject to a continuous relative sea-level regression, despite corresponding eustatic
64 sea-level rise (e.g. Mörner 1979; Svendsen and Mangerud 1987). In other words, the RSL has been dropping
65 throughout prehistory. As this process is the result of glacial loading, the rate of uplift is more severe towards
66 the centre of the ice sheet. Thus, some areas on the outer coast have had a more stable RSL or been subject
67 to marine transgression (e.g. Romundset et al. 2015; Svendsen and Mangerud 1987). These conditions
68 are directly reflected in the archaeological record. In areas where the sea-level has been stable over longer
69 periods of time, people have often reused coastal site locations multiple times and over long time-spans,
70 creating a mix of settlement events that are difficult to disentangle (e.g. Hagen 1963; Reitan and Berg-Hansen
71 2009). Transgression phases, on the other hand, can lead to complete destruction of the sites, bury them
72 in marine sediments, or in the outermost periphery, still submerged today (Bjerck 2008a; Glørstad et al.
73 2020). This can lead to a hiatus in the archaeological record for certain sub-phases in the impacted areas.
74 Comparatively, given a continuous and still ongoing shoreline regression from as high as c. 220 m above
75 present sea-level in the inner Oslo fjord, any one location in south-eastern Norway has only been shore-bound
76 within a relatively limited time-span, and the sites have not been impacted by any transgressions (Hafsten
77 1957, 1983; Romundset et al. 2018; Sørensen 1979). This makes the region especially useful for evaluating
78 the assumption of a shore-bound settlement pattern over a long and continuous time-span.

79 The method of shoreline dating has been met with scepticism as related to the fundamental premise that
80 most sites would have been consistently shore-bound, been characterised as a relative dating method for
81 sites located at different elevations within a constrained geographical area, or been argued to offer no more
82 than a earliest possible date for when a site could have been in use (see review by Nordqvist 1999). The
83 most common application in Norway has arguably been to use the method to provide an approximate date
84 for the occupation of the sites, often in combination with other dating methods (see for example chapters
85 in Damlien et al. 2021; Jakslund 2014; Melvold and Persson 2014; Reitan and Persson 2014; Reitan and
86 Sundström 2018; Solheim 2017 and below). Recently the method has also been used independently to date a
87 larger number sites to get a general impression of site frequency over time. This is done by aggregating point
88 estimates of shoreline dates in 100, 200 or 500 year bins (Breivik 2014; Breivik and Bjerck 2018; Fossum




Figure 1: Deglaciation at 1000 year intervals from c. 17–8 kyr BCE. The study area defined later in the text is marked with a red outline (deglaciation data from Hughes et al. 2016, although see Romundset et al. 2019 for the study area).

⁸⁹ 2020; Nielsen 2021; Solheim and Persson 2018). In his review, Nordqvist (1999) argues that there can be
⁹⁰ little doubt concerning the general applicability of the method – what is less clear is the level of reliability
⁹¹ and chronological resolution that it can offer (see also Johansen 1963).

⁹² The shore-bound settlement location of prehistoric hunter-fisher-gatherers in Norway is generally believed to
⁹³ follow both from the exploitation of aquatic resources and from movement and communication, which would
⁹⁴ have been efficient on waterways (Bjerck 1990; Brøgger 1905:166; also discussed in Berg-Hansen 2009; Bergsvik
⁹⁵ 2009). The same logic has also been extended to the hinter- and inland regions, where sites are to be located
⁹⁶ along rivers and lakes (Brøgger 1905:166; Glørstad 2010:57–87; although see Gundersen 2013; Mjærum 2018;
⁹⁷ Schülke 2020). This is to take a dramatic turn at the transition to the Late Neolithic, around 2400 BCE, with
⁹⁸ the introduction of the Neolithic proper (Prescott 2020; Solheim 2021). The introduction of a comprehensive
⁹⁹ Neolithic cultural package, including a shift to agro-pastoralism and the introduction of the farm is to have
¹⁰⁰ led site locations to be more withdrawn from the shoreline (e.g. Bakka and Kaland 1971; Østmo 2008:223;
¹⁰¹ prescott2012?). That is not to say that waterways and aquatic resources were no longer exploited, but
¹⁰² rather that these activities would not have been as tightly integrated with settlement and tool-production
¹⁰³ areas as in preceding periods (Glørstad 2012). At an earlier stage, at the transition to the Early Neolithic
¹⁰⁴ (c. 3900 BCE), pottery is introduced to the sites, and there are some indications of an initial uptake of
¹⁰⁵ agriculture at some sites in the Oslo fjord region. However, this appears to be small in scale and is believed
¹⁰⁶ to be combined with a continued and predominantly hunter-gatherer life-way, possibly followed by a complete
¹⁰⁷ de-Neolithisation in the Middle Neolithic (Erik 1955; Nielsen et al. 2019; Østmo 1988:225–227). Nielsen
¹⁰⁸ (2021) has recently argued that the initial uptake of agriculture in Early Neolithic south-eastern Norway is
¹⁰⁹ combined with a more complex settlement pattern, and that a simple foraging/agricultural dichotomy would
¹¹⁰ underplay the variation present in the Early and Middle Neolithic settlement data (see also e.g. Amundsen
¹¹¹ et al. 2006; Østmo 1988; Solheim 2012:74). Seen in relation to the question of interest here, the empirical
¹¹² expectation for the above outlined development would thus be a predominantly shore-bound settlement in the
¹¹³ Mesolithic, possibly followed by a more varied association between sites and the shore-line with the transition
¹¹⁴ to the Early Neolithic around 3900 BCE, and finally a decisive shift with the Late Neolithic c. 2400 BCE.

¹¹⁵ Based on the generally accepted premise that most pre-Late Neolithic sites in south-eastern Norway located
¹¹⁶ lower than the marine limit were situated on or close to the contemporaneous shoreline, it is common to err
¹¹⁷ on the side of a shore-bound site location unless there is strong evidence to suggest otherwise. This is for
¹¹⁸ example reflected in survey projects, which are often guided by both a digital and mental reconstruction
¹¹⁹ of past sea-levels (e.g. Berg-Hansen 2009; Eskeland 2017). Similarly, following an excavation, if typological
¹²⁰ indicators in the assemblages correspond with available shoreline displacement curves, a shore-bound site
¹²¹ location is often assumed, even if the typologically informed date-span is too wide to decisively verify this.
¹²² It is also common to combine this with a qualitative consideration of the landscape surrounding the sites,
¹²³ and an evaluation of the degree to which the site location would have been sensible if the site was not shore
¹²⁴ bound (e.g. Jakslund 2014; Johansen 1963; Nummedal 1923). This can for example pertain to accessibility. If
¹²⁵ the site is situated on a ledge in a steep and jagged area of the present day landscape it would make intuitive
¹²⁶ sense that the site was in use when the ocean reached closer to its elevation, as the site would have been
¹²⁷ accessible by means of watercraft. Although it appears that the arguments for such site locations are sensible
¹²⁸ and can for the most part be assumed to hold, comprehensive evaluations and attempts at quantification of
¹²⁹ this tendency are relatively few (see also Ilves and Darmark 2011).

¹³⁰ One of the more extensive evaluations of the relationship between archaeological radiocarbon dates and
¹³¹ RSL-change was done by Solheim and colleagues (Breivik et al. 2018; Solheim 2020), who compared 102
¹³² radiocarbon dates from 33 Mesolithic sites on the western side of the Oslo fjord to the displacement curve
¹³³ for the Larvik area. They found an overlap between the probability density of the radiocarbon dates with
¹³⁴ the shoreline displacement curve for 86.5% of the sites. However, where there was a discrepancy, the main
¹³⁵ occupation of the sites are still believed to have been shore-bound rather than associated with the deviating
¹³⁶ ¹⁴C-dates. This is based on typological and technological characteristics of the assemblages. Whether these
¹³⁷ mismatches represent later shorter visits that are responsible for the younger radiocarbon dates, or whether
¹³⁸ these dates are entirely erroneous can be difficult to evaluate (e.g. Persson 2008; Schülke 2020). However,
¹³⁹ this distinction is not deemed critical here, as what is of interest is settlements and tool-production areas as
¹⁴⁰ evidenced by artefact inventories or multiple site features. Not remnants of stays as ephemeral to only be

discernible by individual features or dubious ^{14}C -dates. The evaluation of the relevance of radiocarbon dates to settlement activity will here therefore be entirely dependent on, and follow the discretion of the original excavation reports.

Other previous evaluations of the correspondence between radiocarbon- and RSL-informed dates have typically followed the same structure as that of Breivik et al. (2018), involving a visual inspection of radiocarbon probability density functions plotted against local shoreline displacement curves based on the elevation of the site (e.g. Åkerlund et al. 1995; Åstveit 2018; Solheim 2020; see also Bjerck 2008b; Kleppe 1985; Ramstad 2009). This approach has a couple of limitations. First of all, the displacement curves are sometimes applied directly to larger study areas, with only some studies having taken the variable uplift-rates into account when performing this comparison (e.g. Åstveit 2018; Fossum 2020; Møller 1987; Persson 2008). Secondly, with this method, the wider the uncertainty range associated with either radiocarbon date or displacement curve, the higher the probability that the confidence intervals overlap, and the higher the probability that we conclude in favour of our hypothesis. This thus leads to an inferential framework that favours uncertainty, which is hardly desirable. In statistical terms this follows from the fact that while one cannot conclude that two dates are different if their confidence intervals overlap, this does not necessarily mean that they are the same. The question thus necessitates a flip from a null-hypothesis of no significant difference, to one of equivalence (e.g. Lakens et al. 2018), as the question of interest is effectively one of synchronicity between events (cf. Parnell et al. 2008). Another limitation of this often-employed method is that it only takes into account the vertical distance between the sites and the sea-level. While this is the main parameter of interest for shoreline dating, the practical implications of a vertical difference in RSL will be highly dependent on local topography and bathymetry. RSL-change can have more dramatic consequences in a landscape characterised by a low relief, as the horizontal displacement of the shoreline will be greater. Taking the spatial nature of the relationship between site and shoreline into account will consequently help get more directly at the behavioural dimension of this relation, and help move the analysis beyond a purely instrumental consideration of the applicability of shoreline dating.

3 Data

To get at the relationship between sites and the contemporaneous shoreline this analysis was dependent on a study area with good control of the trajectory of prehistoric shoreline displacement. While there is displacement data available for other areas of south-eastern Norway (Creel et al. 2022; e.g. Hafsten 1957; Sørensen 1979, 1999), considerable methodological developments in recent years means that the most well-established displacement curves are from the region stretching from Horten county in the north-east, to Arendal in the south-west. This area has newly compiled displacement curves for Horten (**romundset2021?**), Larvik (Sørensen et al. in prep; Sørensen, Henningsmoen, et al. 2014; Sørensen, Høeg, et al. 2014), Tvedstrand (Romundset 2018; Romundset et al. 2018), and Arendal (Romundset 2018).

The employed shoreline displacement data is based on the so-called isolation basin method (e.g. Kjemperud 1986; Romundset et al. 2011), which involves extracting cores from a series of basins situated on bedrock at different elevations beneath the marine limit, and dating the transition from marine to lacustrine sediments. Each basin thus represent a high precision sea-level index point (SLIP) which are combined using what has been termed the isobase method to a continuous time series for RSL-change, projected to a common isobase (see Creel et al. 2022:5). Furthermore, to minimise the impact of variable uplift rates, the basins are located in a as constrained area of the landscape as possible. Following from the morphology of the retreating ice sheet, the uplift is more severe towards the north-east, meaning that this needs to be adjusted for in the case that any basins are located any significant distance from the common isobase perpendicular to this gradient (Figure 2). The SLIPs indicate the isolation of the basin from the highest astronomical tide, which is adjusted to mean sea-level in the compilation of the displacement curves based on the present day tidal range. This is assumed to have been the same throughout the Holocene (Sørensen, Henningsmoen, et al. 2014:44). Furthermore, the confidence bands of the displacement curves and their trajectory are quite complex constructs, and are the integrated result of both expert knowledge and more objectively quantifiable parameters. The reason for this is in part that the curves do not only contain uncertainty as related to

190 radiometric dates, which are well defined, but also hold potential error as related to the interpretation and
191 analysis of sediment cores, the nature and condition of the basin outlets and the adjustment to a common
192 isobase, to name but a few Creel et al. (2022). For more details and evaluations done for the compilation of
193 each curve, the reader is therefore referred to the individual publications.

194 The archaeological data compiled for the analysis consists of excavated Stone Age sites with available spatial
195 data from the coastal region between Horten county in the north-east, to Arendal in the south-west (Figure
196 2). These number 155 sites. Of these, 91 sites are associated with a total of 547 radiocarbon dates. Of these,
197 in turn, 67 sites are related to the 259 radiocarbon dates that fall within the Stone Age (9500–1700 BCE),
198 with 95% probability. These sites and ^{14}C -dates form the basis for the analysis. Spatial data in the form of
199 site limits and features, as defined by the excavating archaeologists, were retrieved from local databases at the
200 Museum of Cultural History—the institution responsible for archaeological excavations in the region. In the
201 compiled dataset, each radiocarbon date has been associated with the site features or excavation unit from
202 where they originate, or, where these weren't available, the spatial limit of the entire site. Due to somewhat
203 variable practices between excavations, what available spatial geometry best represents the site limit was
204 decided based on an evaluation of the excavation reports. This means that the limits are variably given as
205 that defined during initial survey, area de-turfed before excavation, area stripped with excavator following
206 the excavation, manually excavated area, or convex hull polygons generated around the site features.

207 Three of the sites have been associated with agriculture, either directly or in the form building structures.
208 The first is Nordby 1 at which the ^{14}C -dates are associated with a Late Neolithic long-house (Gjerpe and
209 Bukkemoen 2008). The Middle Neolithic phase at Kvastad A2 (Stokke and Reitan 2018) and Late Neolithic
210 phase at Nauen A (Persson 2008) are both directly related to farming activities. Both of these sites also
211 have radiocarbon dates and lithic inventory associated with Mesolithic forager activities. Following from the
212 expected deviance from the settlement patterns that are to characterise forager sites, these agricultural phases
213 are highlighted in the analysis below. Finally, Nielsen (2021) has recently suggested that Early and Middle
214 Neolithic features from the otherwise younger sites Bratsberg (Wenn 2012) and Larønningen (Røberg 2012)
215 could be related to early agricultural activity in the Oslo fjord region. Due to the uncertain and somewhat
216 speculative nature of this suggestion, these are omitted here.

217 The elevation data used for the analysis is a digital terrain model (DTM) freely available from the Norwegian
218 Mapping Authority (Norwegian Mapping Authority 2018, <https://hoydedata.no>). It was here opted for the
219 10m resolution DTM rather than the higher-resolution 1m version. In addition to resulting in considerably less
220 processing time, the higher resolution elevation model is more vulnerable to smaller-scale modern disturbances
221 that the 10m version is not impacted by. The 10m resolution DTM of the study area is a down-sampled
222 version of the 1m version and has a height accuracy with a systematic error of 0.1m (Norwegian Mapping
223 Authority 2018). All data and R programming code (R Core Team 2021) required to run the analyses, as
224 well as the derived data are freely available in an online repository at <https://osf.io/7f9su/>, organised as a
225 digital research compendium following Marwick et al. (2018).

226 4 Methods

227 The method of shoreline dating is based on the spatial relationship between two phenomena, occupation of
228 sites and shoreline displacement, each associated with their own range of temporal uncertainty. The first task
229 was therefore to ascribe likely date ranges and associated uncertainty to these dimensions. To take account
230 of the gradient in the isostatic rebound, the trajectory of shoreline displacement was first interpolated to
231 each site location based on the distance to the isobases of the displacement curves using inverse distance
232 weighting (e.g. Conolly 2020; Conolly and Lake 2006:94–97). This was done for each year along the entirety
233 of the curves, weighting the interpolation by the squared inverse of the distances. The result of this process is
234 shown for an example site in Figure 3. For the date ranges associated with the sites, all radiocarbon dates
235 were first individually calibrated using the IntCal20 calibration curve (Reimer et al. 2020) using OxCal v4.4.4
236 (Bronk Ramsey 2009) through the oxcAAR package for R (Hinz et al. 2021). Radiocarbon dates associated
237 with each site were then grouped if they overlapped with 99.7% probability, meaning these were effectively
238 taken to represent the same settlement phase. In the case where there are multiple dates believed to belong




Figure 2: A) Distribution of the 67 analysed sites relative to the isobases of the displacement curves. The isobases have a direction of 327° (Romundset et al. 2018), B) Displacement curves. Note the increasing⁷ steepness of the curves towards the north-east.

239 to a single phase, these were modelled using the Boundary function in OxCal and then summed. Multiple
240 phases at a single site were treated as independent of each other.

241 As the excavation of archaeological sites typically follow from residential and commercial development, as well
242 as the expansion of infrastructure, the area immediately surrounding the sites has sometimes been severely
243 impacted by modern disturbances. In addition to employing 10m resolution DTM to alleviate some of these
244 issues, this also necessitated some additional editing of the elevation raster. This involved manually defining
245 the extent of problem areas such as railways, highways, quarries and the like. The DTM values on these were
246 then set to missing, and new elevation values were interpolated from the surrounding terrain. This was done
247 using regularised spline interpolation with tension (e.g. Conolly 2020), using the default settings of r.fillnulls
248 from GRASS GIS (GRASS Development Team 2017) in R through the package rgrass7 (Bivand 2021). In
249 addition to code and original spatial data being available in the digital research compendium for this paper,
250 the analysis of each individual site is presented in the supplementary material where it has been noted when
251 the area surrounding a site has been edited in this manner.

252 Armed with a likely date range for the occupation(s) of each site, an estimated trajectory of relative sea-level
253 change at that location, and a DTM edited to remove substantial modern disturbances, the simulations were
254 performed. A single simulation run involved first drawing a single year from the posterior density estimate of
255 a given occupation phase of a site (Figure 4). This year then has a corresponding likely elevation range for
256 the contemporaneous shoreline from which an elevation value was drawn uniformly, using intervals of 5cm.
257 The sea-level was then raised to this elevation on the DTM by defining all elevation values at or below this
258 altitude as missing. Polygons were then created from the resulting areas with missing values. The horizontal
259 distance was then found by measuring the shortest distance between site and sea polygons, and the vertical
260 distance by subtracting the elevation of the sea-level from the lowest elevation of the site polygon. The
261 topographic distance between site and sea was also found by measuring the distance while taking into account
262 the slope of the terrain on the DTM. This was done using the topoDistance package for R (Wang 2019).
263 The topographic distance was measured between the site polygon and the horizontally closest point on the
264 shoreline. This means that the distance is not necessarily measured as the closest topographic distance to the
265 shoreline, but rather as the shortest topographic path to the horizontally closest point on the shoreline. Not
266 finding the topographically closest point significantly reduced the computational cost of the analysis, and is
267 deemed unlikely to have a considerable impact on the results given the distances considered. The shortest
268 topographic path was found using the Moore neighbourhood of eight cells (e.g. Conolly and Lake 2006:253;
269 Herzog 2013). In the case where the sea-polygons intersects the site polygon, all distance measures were
270 set to zero. In the case that the sea-polygons completely contain the site, the horizontal and topographic
271 distance measures were made negative, and the vertical distance was instead measured to the highest point
272 on the site polygon. While it is safe to assume that an archaeological site was not occupied when it was
273 located beneath sea-level, a negative result can reflect the inherent uncertainty in this procedure, and might
274 also help identify discrepancies in displacement data or radiocarbon dates. Negative values were therefore
275 retained with the exception of for the sites Gunnarsrød 5 and Pjonkerød R1, where the negative values are
276 believed to result from modern disturbances in the DTM rather than the ^{14}C -dates or displacement curves
277 (see supplementary material for more details).

278 This process was repeated 1000 times for each phase for each site. The choice of 1000 simulation runs follows
279 from an evaluation of when the mean distances between site and shoreline converged when running 5000
280 iterations of the simulation on the site Hovland 5, available in the supplementary material (cf. Crema et al.
281 2010:1125). Hovland 5 was chosen for this evaluation as it has a fairly imprecise date and is located in area
282 of quite complex topography.

283 5 Simulation results

284 Overall, as is indicated by the measures for central tendency and the almost solid line along the 0m mark on
285 the y-axes, the simulations show that the sites tend to have been situated close to the shoreline when they
286 were in use (Figure 6). Some of the sites are situated considerable distances from the shoreline when the dates
287 believed to be erroneous in the original reports are included (Figure 6A), but if one accepts the interpretation




Figure 3: Example site Hegna vest 1 (Fossum 2017). A) Location of the site in the present day landscape. The red outline is the site limit. B) Radiocarbon dates associated with the site. Fill colour indicates what dates are assumed to belong to the same settlement phase. Multiple dates are modelled using the Boundary function in OxCal and then summed. The red outline indicates that the date does not match the typological indicators in the artefact assemblage of the site. C) The location of the site within the study area relative to isobases of the employed displacement curves. D) Displacement curve interpolated to the site location.




Figure 4: Example of a single simulation run on the site Hegna vest 1. A) The simulation starts by drawing a single year from the posterior density estimate. B) This then corresponds to an elevation range on the interpolated displacement curve. A single elevation is drawn uniformly from this range using 5cm intervals. C) The sea-level is then adjusted on the DTM to this elevation and the various distance measures are found. D) The numerical result of the simulation run.




Figure 5: The result of 1000 simulation runs for each of the two groups of dates on the site Hegna vest 1. The first column of plots shows the radiocarbon probability density function from where dates were drawn during simulation. The second column displays the result of simulating the raised sea-level 1000 times. The more opaque the colour, the more times the sea-level was simulated to that location. The third column shows violin plots of the different distance measures across all simulations.

288 that these do not date the main occupation of the sites, as is indicated by the artefact inventories, Figure 6B
289 gives considerable support to the notion that the sites were in use when they were situated on or close to the
290 contemporaneous shoreline. The distances for some of the earliest sites appears somewhat high, but this
291 can likely be explained as the result of the steepness of the displacement curves for the earliest part of the
292 Holocene (Figure 2B), which leads the uncertainty of the ^{14}C -dates to give a wider possible elevation range
293 for the sea-level. Another immediately striking result is the apparent deviation from the shoreline towards
294 the end of the Stone Age, corresponding with the literature. From around 2500 BCE several sites are situated
295 a considerable distance from the shoreline, and while a couple remain horizontally and topographically close,
296 most appear to be elevated a considerable distance from the sea-level, as indicated on the plot for vertical
297 distance. There are also a couple of sites located some distance from the shoreline just after 4000 BCE. While
298 the sample size is limited, this would thus be in line with a development that sees an increase in settlements
299 located in the immediate inland around this time.

300 The negative values around 8000 BCE originate from the sites Løvås 1, 2 and 3. These are recently excavated,
301 well-dated sites situated in a relatively undisturbed area of the landscape. While there would be a danger of
302 circularity of having archaeological sites inform a reconstruction RSL-change, and in turn use these to evaluate
303 the degree of shore-bound settlement, the sites do clearly represent a upper limit for the sea-level, as they
304 would not have been in use when located under water. It could therefore seem that the Løvås sites represent
305 a case where the archaeological material indicates a slight discrepancy in the geological reconstruction of
306 shoreline displacement in the area.

307 Accepting that shoreline dating appears to loose utility around the transition to the Late Neolithic, as
308 indicated by the clear deviation in site location from the shoreline after this, the results for from Figure 6B is
309 given again in Figure 7A, excluding all simulation results younger than 2500 BCE. Furthermore, all negative
310 values have here been set to zero, under the assumption that these result from uncertainty or errors in the
311 data, and not actual site locations. The resulting best point estimate for the vertical distance between sites
312 and shoreline for the pre-Late Neolithic is given by the median at 4m, while 95% of the values fall within the
313 range 0–18m. That is, for 95% of the cases, the shoreline was simulated to be situated on or down to 18m
314 below the site location. While these values remain the same when only the Mesolithic dates are included
315 (Figure 7B), the mean and standard deviation is slightly constrained. Furthermore, while the median for
316 horizontal and topographic distance is only 10m across all plots in Figure 7, the variation in the statistics for
317 dispersion is greater, illustrating the point that minor variations in vertical distance can have substantial
318 consequences for these distance measures, depending on the surrounding topography.

319 An exponential function has been fit to the distributions for vertical distance using maximum likelihood
320 estimation (Figure 7). While it makes theoretical sense that a process of exponential decay explains this
321 relationship, it is also clear that this does not perfectly match the data. However, this can at least in part be
322 related to methodological factors, where the accumulation of distance-values on the 0m mark likely follow
323 from forcing negative values to zero, from the resolution of the spatial data, and from defining intersecting
324 sea- and site polygon as having a distance of zero. If one accepts this, the exponential decay function used
325 for describing the vertical distance between sites and shoreline can be combined with the displacement data
326 to provide a method for shoreline dating that takes this distance into account:

$$p(t) = \binom{n}{k} p^k (1-p)^{n-k} \quad (1)$$

327 Where x is [...] site_elevation, shoreline_displacement (upper and lower bound?), relationship between site
328 and shoreline. For each offset, subtract

329 (1) is used to shoreline date the same sites from where this relationship was derived, using the decay ratio of
330 0.173 (Figure 8) which was found when including all pre-Late Neolithic results (Figure 7A). For illustrative
331 purposes the Late Neolithic sites were also included. Following from having defined the distance between
332 intersecting sea- and site polygons as zero during simulations, the sites were dated using the mean elevation
333 of the site polygons to allow for some variation in elevation over the site limits. The synchronicity between
334 radiocarbon and shoreline dates was then evaluated using the method presented by Parnell et al. (2008,
335 Figure 9). Here, 100,000 age samples drawn from the probability density function of each shoreline date

were subtracted from 100,000 age samples drawn from the corresponding modelled ^{14}C -dates. The resulting range of the 95% highest density region (HDR, Hyndman 1996) was then checked to see if it crosses zero, in which case the dates are considered to be in agreement. When excluding the earliest phase at Gunnarsrød 5, the deviation of which is to be expected based on the simulation results (see above), the shoreline date correspond to the radiocarbon dates in 58 out of 68 cases (84%). Only including dates modelled to be older than 2500 BCE with 95% probability, i.e. older than the Late Neolithic, improves this to 56 out of 61 cases (92%). When only including dates older than 4000 BCE with 95% probability, i.e. only Mesolithic, the success rate is further increased to 46/49 (94%). The three failed Mesolithic shoreline dates are from the early sites Langemyr and Kvastad A2, with the possible implication that a lower decay ratio than what is used for characterising the distance between site and shoreline for all sites in aggregate should be used for sites known to be from the earliest part of the Mesolithic (see also Figure 6).




Figure 6: The result of running the analysis across all sites. Each data point is plotted with some transparency, meaning that the more intense the colour, the more often those values occurred. Results associated with agricultural activities are plotted in grey. The first row A) shows the result of including all dates to the Stone Age, including those seen as otherwise unrelated to the main occupation of the sites. The second row B) shows the result of excluding these. The table under each plot lists some corresponding statistics for central tendency and dispersion.



A**B**

Figure 7: Histograms showing the simulated distance from the shoreline using dates corresponding to the site inventory. Negative values have been set to zero. A) Simulated results older than 2500 BCE, and B) simulated results older than 4000 BCE.




Figure 8: The result of backwards shoreline dating the sites with radiocarbon dates corresponding to the artefact inventory using the method proposed here. The shoreline dates are plotted in grey and underlined¹⁵ with the 95% HDR in black. These are plotted against the modelled radiocarbon dates, which are given colour from oldest to youngest phase for each site, defined by non-overlapping dates at 99.7% probability.




Figure 9: Evaluation of the agreement between the shoreline dates and radiocarbon dates given in Figure 9. When the range of the 95% HDR for age difference crosses zero, the shoreline and radiocarbon dates are considered synchronous. Line segments with vertical bars indicate that this does not cross zero and that the dates are not in agreement. The division and colour coding at the top of the plots reflect the division of site phases given in Figure 9.

347 6 Re-dating previously shoreline dated sites

348 To further explore the implementation for shoreline dating outlined above, excavated and shoreline dated
349 Stone Age sites within the study area where ^{14}C -dates are not available or are not believed to date the main
350 occupation of the sites have been subjected to the approach (Figure 10). The resulting dates are compared
351 to those originally proposed in the excavation reports for the sites (numerical results are available in the
352 supplementary material). To avoid issues with recent disturbances on the DTM, the sites have been dated
353 based on the mean of the altitudes provided in the report for each site.

354 The comparison with previously reported dates is an illustrative, but unfair exercise for a few reasons. First
355 of all the dates provided in the reports are typically stated to be a very rough estimate, and are sometimes
356 given as a point estimate with an undefined, but implied or explicit uncertainty range. Secondly, seeing
357 as these reports are from various dates in time, many are based on now outdated data on RSL-change.
358 Finally, they are sometimes only meant to indicate a lower bound for when the sites could have been in use.
359 Overall, the results suggest that shoreline dating has generally been applied with a fairly reasonable degree
360 of success, seeing as these dates have typically been interpreted and informed research in an approximate
361 manner (although see e.g. Roalkvam 2022). That being said, the results do also indicate that shoreline dating
362 has at times been applied with an exaggerated degree of precision. While the implications of a more stable
363 RSL-change for the duration of use and re-use of site locations, and consequently the precision of shoreline
364 dating, is well known, it appears somewhat under-appreciated. The results also highlight the spatial and
365 temporal contingency of the method, illustrated by the variation in the range of the 95% HDRs for the dates.
366 In some cases the method provides a very precise date range and in others it offers little more than a *terminus*
367 *post quem*. This is dependent on the steepness of the displacement curves, leading to the general pattern of
368 older sites situated towards the north-east getting more precise dates (cf. Figure 2B). Further, as some of the
369 date ranges extend well beyond major chronological divisions, even into the Iron Age, they could be severely
370 and securely constrained with only cursory reference to typology. While this would be trivial in some cases,
371 the nature and uncertainty inherent to the method still means that this is arguably a required exercise that
372 should be explicitly performed. This final point also points to the possibility of drawing on other temporal
373 data, for example within a Bayesian framework, to further improve the precision of the dates that can be
374 achieved with shoreline dating.

375 Not least following from the fact that relatively few Preboreal ^{14}C -dates associated with anthropogenic
376 activity have been achieved in Norway (Åstveit 2018; Damlien and Solheim 2018; Kleppe 2018), the shoreline
377 dating of the earliest sites is essential for understanding the pioneer settlement and the initial colonisation
378 of the Scandinavian peninsula (e.g. Bang-Andersen 2012; Berg-Hansen 2018; Breivik 2014; Fuglestvedt
379 2012; Glørstad 2016). The shoreline dated Preboreal sites from the Brunlanes-project are among the earliest
380 known sites in Norway (Jaksland 2012a, 2012b; Jaksland and Persson 2014). These have a distinct Early
381 Mesolithic artefact inventory and are situated in a steep area of the landscape where it would be difficult to
382 envision use of the sites after the sea retreated any significant distance from their location. In the original
383 publication of the sites, Jaksland (2014) provides a thorough discussion of shoreline dating in general, and as
384 used for the dating of the Brunlanes sites specifically. A comparison of his results and the ones achieved
385 using the above-outlined approach are given in Figure 11A. The sites have been dated using what Jaksland
386 (2014) gives as the lowest elevation of finds at each site, and by employing a exponential decay ratio of
387 `rround(expfit3$estimate, 3)`, to allow for more deviance in the distance between site and shoreline. This
388 corresponds to the decay ratio for sites older than 7000 BCE in Figure 7.

389 The small discrepancies between the achieved results mainly follow from the fact that a slightly updated
390 version of the local displacement curve is applied here (cf. Sørensen et al. in prep). Jaksland's dates are
391 given a flat 200 and 50 year uncertainty range starting from what he gives as the earliest possible date. The
392 200 year uncertainty range is given if the sites were to be considered in isolation, while the argument for
393 the uncertainty range of only 50 years is based on the location of the sites relative to each other. Since they
394 are located in such a constrained and steep area of the landscape, the difference in elevation between the
395 sites is argued to establish their relative date and thus constrain the uncertainty ranges so that they don't
396 overlap. This information is not integrated in the approach outlined here, but could justify further reducing
397 the uncertainty ranges. Although their accuracy is of course ultimately dependent on the veracity of the




Figure 10: Re-dating previously shoreline dated sites in the study area without radiocarbon dates or with radiocarbon dates that do not correspond to the artefact inventories. The range of the 95% HDRs in grey are compared to the dates originally proposed by the excavation reports in red.

geological reconstruction, the high rate of RSL-change in this period does nonetheless result in very precise dates. Above it was shown how additional temporal data could be combined with the method to improve its accuracy and precision. This example, on the other hand, highlights the fact that the spatial nature of the method means that a consideration of the surrounding terrain and other sites can also help in increasing the precision of the method if this can be used to exclude certain sea-levels as unlikely for when a site was in use. One approach could also be to assess the spatial implication of a proposed shoreline date by simulating the adjusted sea-levels, as is done for Paurer 1 in Figure 11B, followed for example by a visual evaluation of the topography or by evaluating the distance and steepness of the slope to the shoreline. Based on this, it could conceivably be possible to exclude certain elevations as unlikely for the position of the shoreline when the site was in use. Such approaches would make less of an impact in this setting, where the 95% HDR is already quite constrained, but could considerably improve the precision of the method in cases where RSL-change has been less severe (cf. Figure 10).




Figure 11: Shoreline dating of the Brunlanes sites using site altitudes provided by Jaksland (2014:tab.4). A) The result of applying the approach to shoreline dating outlined above. The shoreline date in grey is underlined with the 95% HDR in black. Dates provided by Jaksland (2014) are plotted in red. The box indicates a 50 year uncertainty range which in combination with the red line extends 200 years. B) Map showing the centroids of the Paurer sites and Sky 1. The sea-level has been simulated using the probability density associated with the shoreline date for Paurer 1 (see also map in Jaksland 2014:fig.12a). Paurer 1 is the red point.

410 7 Concluding remarks

411 The most immediate contribution of this paper is what must be considered a confirmation of previous research
412 into the relation between coastal Norwegian Stone Age sites and the prehistoric shoreline. This is indicated
413 by the close relationship between sites and the shoreline up until the transition to the Neolithic at c. 4000
414 BCE, after which a couple of sites become situated some distance from the sea, followed by a more decisive
415 break at the transition to the Late Neolithic at c. 2500 BCE. This development is in agreement with the
416 literature. Furthermore, based on the quantitative nature of these findings, an initial formulation of a refined
417 method for the shoreline dating of pre-Late Neolithic Stone Age sites has been proposed. In addition to
418 taking the distance between sites and the isobases of the displacement curves into consideration when dating
419 the sites, this involves implementing equation (1) to account for the distance between the sites and the
420 shoreline. When no other information is available, it can at present be recommended to use the empirically
421 derived exponential decay ratio of 0.173 (Figure 10A) to characterise this relationship. Furthermore, while
422 this remains to be formalised and explored further, it was also showed how the accuracy of the method can
423 be improved by including more information, both with reference to the topographic location of the sites and
424 other temporal data. As the precision of the method is both geographically and temporally contingent due
425 to the trajectory of RSL-change, where older sites situated towards the north-east in the study area will
426 get a more precise date than younger sites located towards the south-west, the impact of such additional
427 information will also vary.

428 Future investigations and radiocarbon dates from Stone Age sites in the region can not only be used to
429 further evaluate and adjust the findings reported here, but a larger sample size could also lay the foundations
430 for refining the method by identifying subsets of sites for which the application of the method could be
431 adjusted. Given it's behavioural nature, it would for example seem likely that dimensions such as the nature
432 and purpose of visits to the sites will have implications for how close to the shoreline they were located.
433 Furthermore, other dimensions related to the topographic location of the sites could be similarly explored.
434 This for example pertains to the exposure of sites to wave action, which is likely to have been of concern
435 (e.g. Roalkvam 2020), and which presumably has implications for how close to the shoreline people settled
436 (Blankholm 2020; Helskog 1978). This is also related to the fact that while the mean sea-level is used for
437 dating the sites, a consideration of the tidal range could possibly also have implications for the site location
438 relative to the shoreline, depending on the topography (Helskog 1978). The potential of dimensions such as
439 these was also hinted at here with the estimation and cursory treatment of the horizontal and topographic
440 distance to the shoreline. If patterns related to such locational patterns can be discerned and unpicked, this
441 will not least be useful for improving the shoreline dating of sites which have only been surveyed and where
442 little information on the site beyond its location is available.

443 Some limitations and sources of likely variation and uncertainty that have not been considered should also be
444 mentioned. First of all the sample size is quite strained and the future addition more sites might alter the
445 picture considerably. Secondly, the DTM has only been corrected for major modern disturbances. This means
446 that erosion, although likely not that prevalent, has not been taken into account. Thirdly, the DTM has a
447 vertical error which could also benefit from being integrated in the analysis (cf. Lewis 2021). Fourthly, the
448 displacement curves were here interpolated to all site locations without accounting for increased uncertainty
449 as one moves further away from the isobases of the displacement curves. This is also related to the fact that
450 the RSL data can be handled in different ways than with the isobase method that has been used for the
451 compilation of the employed displacement curves (cf. Creel et al. 2022). Fifthly, neither the question of how
452 site limits are defined nor the elevation range over which these extend was given much consideration. Finally,
453 the division of phases at each site was here simply done by treating radiocarbon dates not overlapping at
454 99.7% as unrelated. This could also be done differently. While each of these factors will have variable impact
455 on the final results, they clearly represent dimensions which would all benefit from further consideration.

456 Finally, this analysis employed a simulation approach to integrate multiple sources of spatio-temporal
457 uncertainty. Here this was simply used to inform the question of the distance between sites and the shore-line.
458 However, this method and general framework can be extended to a wide range of use-cases where one needs
459 to visualise, and quantitatively or qualitatively evaluate the relationship between archaeological phenomena,
460 the prehistoric shoreline, and the uncertainty inherent in this reconstruction.

461 8 References

- 462 Åkerlund, Agneta
463 1996 *Human responses to shore displacement: Living by the sea in Eastern Middle Sweden during the Stone Age*. Riksantikvarieämbetet, Stockholm.
- 464
465 Åkerlund, Agneta, Jan Risberg, Urve Miller, and Per Gustafsson
466 1995 On the applicability of the ^{14}C method to interdisciplinary studies on shore displacement and settlement location. *PACT* 49:53–84.
- 467
468 Amundsen, Øystein, Stig Knutsen, Axel Mjærum, and Gaute Reitan
469 2006 Nøkleby i ski – en tidligeolittisk jordbruksboplass? *Primitive tider* 9:85–96.
- 470
471 Åstveit, Leif Inge
472 2018 The Early Mesolithic of Western Norway. In *Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 231–274. Equinox, Sheffield.
- 473
474 Bakka, Egil, and Peter Emil Kaland
475 1971 Early farming in Hordaland, western Norway. Problems and approaches in archaeology and pollen analysis. *Norwegian Archaeological Review* 4:1–17. DOI:10.1080/00293652.1971.9965136.
- 476
477 Bang-Andersen, Sveinung
478 2012 Colonizing Contrasting Landscapes. The Pioneer Coast Settlement and Inland Utilization in Southern Norway 10,000–9500 Years Before Present. *Oxford Journal of Archaeology* 31:103–120. DOI:10.1111/j.1468-0092.2012.00381.x.
- 479
480 Berg-Hansen, Inger Marie
481 2009 *Steinalderregistrering. Metodologi og forskningshistorie i Norge 1900–2000 med en feltstudie fra Lista i Vest-Agder*. Museum of Cultural History, University of Oslo, Oslo.
- 482
483 2018 Continuity and Change in Late- and Post-glacial Social Networks: Knowledge Transmission and Blade Production Methods in Ahrensburgian and Early Mesolithic North West Europe. In *The Early Settlement of Northern Europe. Transmission of Knowledge and Culture*, edited by Kjel Knutsson, Helena Knutsson, Jan Apel, and Håkon Glørstad, pp. 63–98. Equinox, Sheffield.
- 484
485 Bergsvik, Knut Andreas
486 2009 Caught in the middle: functional and ideological aspects of Mesolithic shores in Norway. In *Mesolithic horizons: Papers presented at the seventh international conference on the mesolithic in europe, belfast 2005*, edited by Sinéad B. McCartan, Rick Schulting, Graeme Warren, and Peter Woodman, pp. 602–609. Oxbow Books, Oxford.
- 487
488 Bevan, Andrew, Enrico R. Crema, Xiuzhen Li, and Alessio Palmisano
489 2013 Intensities, Interactions, and Uncertainties: Some New Approaches to Archaeological Distributions. In *Computational Approaches to Archaeological Spaces*, edited by Andrew Bevan and Mark Lake, pp. 27–52. Left Coast Press, Walnut Creek.
- 490
491 Bivand, Roger
492 2021 *rgrass7: Interface between GRASS 7 geographical information system and r*. R package version 0.2-6.
- 493
494 Bjerck, Hein Bjartmann
495 1990 Mesolithic site types and settlement patterns at Vega, Northern Norway. *Acta Archaeologica* 60:1–32.
- 496
497 2005 Strandlinjedatering. In *Norsk arkeologisk leksikon*, edited by Einar Østmo and Lotte Hedeager, pp. 363–364. Pax, Oslo.
- 498
499 2008a Norwegian mesolithic trends: A review. In *Mesolithic europe*, edited by Geoff Bailey and Penny Spikins, pp. 60–106. Cambridge University Press, Cambridge.
- 500
501 2008b Innledende betraktninger. In *NTNU vitenskapsmuseets arkeologiske undersøkelser ormen lange nye-hamna*, edited by Hein Bjartmann Bjerck, Leif Inge Åstveit, Trond Meling, Jostein Gundersen, Guro Jørgensen, and Staale Normann, pp. 548–551. Tapir Akademisk Forlag, Trondheim.

- 503 Blankholm, Hans Peter
504 2020 In the wake of the wake. An investigation of the impact of the storegga tsunami on the human
settlement of inner varangerfjord, northern norway. *Quaternary International* 549:65–73. DOI:<https://doi.org/10.1016/j.quaint.2018.05.050>.
- 505
506 Breivik, Heidi Mjelva
507 2014 Palaeo-oceanographic development and human adaptive strategies in the Pleistocene–Holocene transition:
508 A study from the Norwegian coast. *The Holocene* 24:1478–1490. DOI:10.1177/0959683614544061.
- 509 Breivik, Heidi Mjelva, Guro Fossum, and Steinar Solheim
510 2018 Exploring human responses to climatic fluctuations and environmental diversity: Two stories
from Mesolithic Norway. *Quaternary International* 465. Impacts of gradual and abrupt environmental changes on Late glacial to Middle Holocene cultural changes in Europe:258–275.
DOI:10.1016/j.quaint.2016.12.019.
- 511
512 Breivik, Heidi, and Hein Bjartmann Bjerck
513 2018 Early mesolithic central norway: A review of research history, settlements, and tool tradition. In *Early
economy and settlement in northern europe. Pioneering, resource use, coping with change*, edited by
Hans Peter Blankholm, pp. 169–206. Equinox, Sheffield.
- 514
515 Brøgger, Waldemar Christofer
516 1905 *Strandliniens Beliggenhed under Stenalderen i Det Sydøstlige Norge*. Norges geologiske undersøkelse,
Kristiania.
- 517
518 Bronk Ramsey, Christopher
519 2009 Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* 51(1):337–360.
DOI:10.1017/S0033822200033865.
- 520
521 Conolly, James
522 2020 Spatial interpolation. In, edited by Mark Gillings, Piraye Hacigüzeller, and Gary Lock, pp. 118–134.
Routledge, London & New York.
- 523
524 Conolly, James, and Mark Lake
525 2006 *Geographical information systems in archaeology*. Cambridge University Press, Cambridge.
- 526
527 Creel, Roger C., Jacqueline Austermann, Nicole S. Khan, William J. D’Andrea, Nicholas Balascio, Blake
528 Dyer, Erica Ashe, and William Menke
529 2022 Postglacial relative sea level change in Norway. *Quaternary Science Reviews* 282:107422.
DOI:10.1016/j.quascirev.2022.107422.
- 530
531 Crema, Enrico R.
532 2012 Modelling Temporal Uncertainty in Archaeological Analysis. *Journal of Archaeological Method and
Theory* 19(3):440–461. DOI:10.1007/s10816-011-9122-3.
- 533
534 2015 Time and probabilistic reasoning in settlement analysis. In *Mathematics and archaeology*, edited by
Juan A. Barcelo and Igor Bogdanovic, pp. 314–334. CRC Press, Boca Raton.
- 535
536 Crema, Enrico R., Andrew Bevan, and Mark W. Lake
537 2010 A probabilistic framework for assessing spatio-temporal point patterns in the archaeological record.
Journal of Archaeological Science 37(5):1118–1130. DOI:10.1016/j.jas.2009.12.012.
- 538
539 Damlien, Hege, Inger Marie Berg-Hansen, Lene Melheim, Mjærum Axel, Per Persson, Almut Schülke, and
540 Steinar Solheim (editors)
541 2021 *Steinalderen i Sørøst-Norge: Faglig program for steinalderundersøkelser ved Kulturhistorisk museum*.
Cappelen Damm Akademisk, Oslo.
- 542
543 Damlien, Hege, and Steinar Solheim
544 2018 The Pioneer Settlement of Eastern Norway. In *Early Economy and Settlement in Northern Europe.
Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 335–367.
Equinox, Sheffield.
- 545
546 De Geer, Gerard
547 1896 *Om Skandinaviens geografiska utveckling efter Istiden*. P. A. Norstedt & Söner, Stockholm.
- 548

-
- 549 Erik, Hinsch
550 1955 Traktbegerkultur – Megalitkultur. En studie av Øst-Norges eldste neolitiske gruppe. *Universitetets*
551 *Oldsaksamling Årbok* 1951–1953:10–177.
- 552 Eskeland, Knut Fossdal
553 2017 *Rapport, arkeologisk registrering. E18 Langangen Rugtvedt, 16/06999, Porsgrunn og Bamble kommune.*
554 Skien.
- 555 Fossum, Guro
556 2020 Specialists facing climate change. The 8200 cal BP event and its impact on the coastal settlement in
the inner oslo fjord, southeast norway. In *Coastal landscapes of the mesolithic: Human engagement*
with the coast from the atlantic to the baltic sea, edited by Almut Schülke, pp. 179–201. Routledge,
London & New York.
- 557 Fuglestvedt, Ingrid
558 2012 The Pioneer Condition on the Scandinavian Peninsula: the Last Frontier of a ‘Palaeolithic Way’ in
Europe. *Norwegian Archaeological Review* 45(1):1–29. DOI:10.1080/00293652.2012.669998.
- 559 Gjerpe, Lars Erik, and Grethe Bjørkan Bukkemoen
560 2008 Nordby 1 – Toskipede hus fra neolitikum-bronsealder og boplasspor fra jernalder. In *E18-prosjektet*
561 *Vestfold. Bind 3. Hus, boplass- og dyrkningspor*, edited by Lars Erik Gjerpe, pp. 7–38. University of
Oslo, Museum of Cultural History, Oslo.
- 562 Glørstad, Håkon
563 2010 *The Structure and History of the Late Mesolithic Societies in the Oslo Fjord Area 6300–3800 BC.*
Bricoleur Press, Lindome.
- 564 2012 Historical ideal types and the transition to the Late Neolithic in South Norway. In *Becoming European.*
565 *The transformation of third millennium Northern and Western Europe*, edited by Christopher Prescott
and Håkon Glørstad, pp. 82–99. Oxbow Books, Oxford & Oakville.
- 566 2016 Deglaciation, sea-level change and the Holocene colonization of Norway. *Geological Society, London,*
567 *Special Publications* 411:9–25. DOI:10.1144/SP411.7.
- 568 Glørstad, Håkon, Jostein Gundersen, Frode Kvalø, Pål Nymoen, David Simpson, and Birgitte Skar
569 2020 Submerged Stone Age from a Norwegian Perspective. In *The Archaeology of Europe’s Drowned*
570 *Landscapes*, edited by Geoff Bailey, Nena Galanidou, Hans Peeters, Hauke Jöns, and Moritz Mennenga,
pp. 125–140. Springer, Cham.
- 571 GRASS Development Team
572 2017 *Geographic Resources Analysis Support System (GRASS) Software, Version 7.2.* Open Source Geospatial
573 Foundation.
- 574 Gundersen, Jostein
575 2013 Verken fjord eller fjell – steinalderen i det kystnære innlandet. Gamle og nye funn fra Notodden i
576 Telemark. *Viking* 76:35–62.
- 577 Hafsten, Ulf
578 1957 De senkvartære strandlinje-forskyvningene i Oslotrakten belyst ved pollenanalytiske undersøkelser.
579 *Norwegian Journal of Geography* 16(1–8):74–99. DOI:10.1080/00291955708622137.
- 580 1983 Shore-level changes in South Norway during the last 13,000 years, traced by biostrati-
581 graphical methods and radiometric datings. *Norwegian Journal of Geography* 37(2):63–79.
582 DOI:10.1080/00291958308552089.
- 583 Hagen, Anders
584 1963 Problemkompleks Fosna. Opphav – kontakt med kontinentale grupper – forholdet til Komsa. In
585 *Boplatsproblem vid Kattegat och Skagerack*, pp. 53–59. Göteborg och Bohusläns forminnesförening &
Institutionen för nordisk fornkunskap, Gothenburg University, Gothenburg.
- 586 Helskog, Knut
587 1978 Late Holocene sea-level changes seen from prehistoric settlements. *Norwegian Journal of Geography*
32:111–119. DOI:10.1080/00291957808552032.
- 588 Herzog, Irmela

- 592 2013 The Potential and Limits of Optimal Path Analysis. In *Computational approaches to archaeological*
593 *spaces*, edited by Andrew Bevan and Mark Lake, pp. 179–211. Left Coast Press, Walnut Creek.
- 594 Hinz, Martin, Clemens Schmid, Daniel Knitter, and Carolin Tietze
- 595 2021 *oxcAAR: Interface to 'OxCal' radiocarbon calibration. R package version 1.1.0.*
- 596
- 597 Hollender, Artur
- 598 1901 Om sveriges nivåförändringar efter människans invandring. *Geologiska Föreningen i Stockholm Förhan-*
599 *dlingar* 23(4):1118–1130. DOI:10.1080/00293652.1975.9965220.
- 600 Hughes, Anna L. C., Richard Gyllencreutz, Øystein S. Lohne, Jan Mangerud, and John Inge Svendsen
601 2016 The last Eurasian ice sheets – a chronological database and time-slice reconstruction, DATED-1.
602 *Boreas* 45(1):1–45. DOI:https://doi.org/10.1111/bor.12142.
- 603 Hyndman, Rob J
- 604 1996 Computing and Graphing Highest Density Regions. *The American Statistician* 50(2):120–126.
- 605
- 606 Ilves, Kristin, and Kim Darmark
- 607 2011 Some Critical and Methodological Aspects of Shoreline Determination: Examples from the Baltic Sea
608 Region. *Journal of Archaeological Method and Theory* 18:147–165. DOI:10.1007/s10816-010-9084-x.
- 609 Jaksland, Lasse (editor)
- 610 2012a *E18 Brunlanesprosjektet. Bind II. Undersøkte lokaliteter fra tidligmesolitikum.* University of Oslo,
611 Museum of Cultural History, Oslo.
- 612 (editor)
- 613 2012b *E18 Brunlanesprosjektet. Bind III. Undersøkte lokaliteter fra tidligmesolitikum og senere.* University
614 of Oslo, Museum of Cultural History, Oslo.
- 615 2014 Kulturhistorisk sammenstilling. In, edited by Lasse Jaksland and Per Persson, pp. 11–46. University
616 of Oslo, Museum of Cultural History, Oslo.
- 617 Jaksland, Lasse, and Per Persson (editors)
- 618 2014 *E18 Brunlanesprosjektet. Bind I. Forutsetninger og kulturhistorisk sammenstilling.* University of Oslo,
619 Museum of Cultural History, Oslo.
- 620 Johansen, Erling
- 621 1963 Kyst(fangst)boplassenes strandbundenhet og strandlinjekronologien. In *Boplatsproblem vid Kattegat*
622 *och Skagerack*, pp. 90–92. Göteborg och Bohusläns fornminnesförening & Institutionen för nordisk
fornkunskap, Gothenburg University, Gothenburg.
- 623 Kjemperud, Alfred
- 624 1986 Late Weichselian and Holocene shoreline displacement in the Trondheimsfjord area, central Norway.
625 *Boreas* 15(1):61–82. DOI:10.1111/j.1502-3885.1986.tb00744.x.
- 626 Kleppe, Else Johansen
- 627 1985 *Archaeological Data on Shore Displacements in Norway.* Norges geografiske oppmåling, Hønefoss.
- 628
- 629 Kleppe, Jan Ingolf
- 630 2018 The Pioneer Colonization of Northern Norway. In *Early Economy and Settlement in Northern Europe.*
631 *Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 13–57. Equinox,
Sheffield.
- 632 Lakens, Daniël, Anne M. Scheel, and Peder M. Isager
- 633 2018 Equivalence Testing for Psychological Research: A Tutorial. *Advances in Methods and Practices in*
634 *Psychological Science* 1(2):259–269. DOI:10.1177/2515245918770963.
- 635 Lewis, Joseph
- 636 2021 Probabilistic Modelling for Incorporating Uncertainty in Least Cost Path Results: a Postdictive Roman
637 Road Case Study. *Journal of Archaeological Method and Theory* 28(3):911–924. DOI:10.1007/s10816-
021-09522-w.
- 638 Marwick, Ben, Carl Boettiger, and Lincoln Mullen

- 639 2018 Packaging Data Analytical Work Reproducibly Using R (and Friends). *The American Statistician*
640 72(1):80–88. DOI:10.1080/00031305.2017.1375986.
- 641 Melvold, Stine, and Per Persson (editors)
- 642 2014 *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og Porsgrunn. Bind 1. Tidlig- Og mellommesolittiske lokaliteter i Vestfold og Telemark*. Portal forlag, Kristiansand.
- 643 Mikkelsen, Egil
- 644 1975 Mesolithic in south-eastern norway. *Norwegian Archaeological Review* 8(1):1118–1130.
645 DOI:10.1080/11035890109445866.
- 646 Milne, Glenn A
- 647 2015 Glacial isostatic adjustment. In, pp. 421–437. Wiley, Chichester.
- 648 Milne, Glenn A., W. Roland Gehrels, Chris W. Hughes, and Mark E. Tamisiea
649 2009 Identifying the causes of sea-level change. *Nature Geoscience* 2(7):471–478. DOI:10.1038/ngeo544.
- 650 Mjærum, Axel
- 651 2018 Hinterland discoveries: Middle Mesolithic woodland utilization and the case of the site Eidsberg, eastern Norway. *Current Swedish Archaeology* 26(1):159–188. DOI:10.37718/CSA.2018.11.
- 652 Møller, Jakob J
- 653 1987 Shoreline relation and prehistoric settlement in northern norway. *Norwegian Journal of Geography* 41:45–60. DOI:<http://dx.doi.org/10.1080/00291958708552171>.
- 654 Mörner, Nils-Axel
- 655 1976 Eustasy and geoid changes. *The Journal of Geology* 84(2):123–151. DOI:10.1086/628184.
- 656 Nielsen, Svein Vatnåg
- 657 2021 Early farming in Southeastern Norway: New evidence and interpretations. *Journal of Neolithic Archaeology* 23:83–113. DOI:10.12766/jna.2021.4.
- 658 Nielsen, Svein Vatnåg, Per Persson, and Steinar Solheim
659 2019 De-Neolithisation in southern Norway inferred from statistical modelling of radiocarbon dates. *Journal of Anthropological Archaeology* 53:82–91. DOI:10.1016/j.jaa.2018.11.004.
- 660 Nordqvist, Bengt
- 661 1995 The Mesolithic settlement of the west coast of Sweden - with special emphasis on chronology and topography of coastal settlements. In *Man and Sea in the Mesolithic. Coastal settlement above and below present sea level*, edited by Anders Fischer, pp. 185–196. Oxbow Books, Oxford.
- 662 1999 The Chronology of the Western Swedish Mesolithic and Late Paleolithic: Old Answers in Spite of New Methods. In *The Mesolithic of Central Scandinavia*, edited by Joel Boaz, pp. 235–253. University of Oslo, Oslo.
- 663 Norwegian Mapping Authority
664 2018 *Produktspesifikasjon. Nasjonal modell for høydedata fra laserskanning (FKB-laser)*. FKB-laser_v30.
- 665 Nummedal, Anders
666 1923 Om flintpladsene. *Norwegian Journal of Geography* 7(2):89–141.
- 667 Østmo, Einar
668 1988 *Etableringen av jordbrukskultur i Østfold i steinalderen*. The University Collection of National Antiquities, University of Oslo, Oslo.
- 669 2008 *Auve. En fangstboplass fra yngre steinalder på Vesterøya i Sandefjord. I. Den arkeologiske del*. Museum of Cultural History, University of Oslo, Oslo.
- 670 Parnell, A. C., J. Haslett, J. R. M. Allen, C. E. Buck, and B. Huntley

- 687 2008 A flexible approach to assessing synchronicity of past events using Bayesian reconstructions of sedimentation history. *Quaternary Science Reviews* 27(19-20):1872–1885. DOI:10.1016/j.quascirev.2008.07.009.
- 688
- 689 Persson, Per
- 690 2008 Nauen 5.2 – Stenåldersboplater och fossil åkermark. In *E18-prosjektet Vestfold. Bind 2. Steinalderboplasser, boplasspor, graver og dyrkningsspor*, edited by Lars Erik Gjerpe, pp. 163–198. University of Oslo, Museum of Cultural History, Oslo.
- 691
- 692 Prescott, Christopher
- 693 2020 Interpreting Complex Diachronic "Neolithic"-Period Data in Norway. In *Farmers at the Frontier – A Pan European Perspective on Neolithisation*, edited by Kurt J. Gron, Lasse Sørensen, and Peter Rowley-Conwy, pp. 381–400. Oxbow Books, Oxford.
- 694
- 695 R Core Team
- 696 2021 *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- 697
- 698 Ramstad, Morten
- 699 2009 Eldre steinalder på Melkøya, representativitet, strandlinjer og transgresjon. In *Undersøkelsene på Melkøya. Melkøyaprosjektet – Kulturhistoriske registreringer og utgravninger 2001 og 2002*, edited by Anders Hesjedal, Morten Ramstad, and Anja R. Niemi, pp. 491–495. Tromsø museum, Universitetsmuseet, Tromsø.
- 700
- 701 Reimer, Paula J., William E. N. Austin, Edouard Bard, Alex Bayliss, Paul G. Blackwell, Christopher Bronk Ramsey, Martin Butzin, Hai Cheng, R. Lawrence Edwards, Michael Friedrich, Pieter M. Grootes, Thomas P. Guilderson, Irka Hajdas, Timothy J. Heaton, Alan G. Hogg, Konrad A. Hughen, Bernd Kromer, Sturt W. Manning, Raimund Muscheler, Jonathan G. Palmer, Charlotte Pearson, Johannes van der Plicht, Ron W. Reimer, David A. Richards, E. Marian Scott, John R. Southon, Christian S. M. Turney, Lukas Wacker, Florian Adolphi, Ulf Büntgen, Manuela Capano, Simon M. Fahrni, Alexandra Fogtmann-Schulz, Ronny Friedrich, Peter Köhler, Sabrina Kudsk, Fusa Miyake, Jesper Olsen, Frederick Reinig, Minoru Sakamoto, Adam Sookdeo, and Sahra Talamo
- 702
- 703
- 704
- 705
- 706
- 707
- 708
- 709
- 710
- 711
- 712
- 713
- 714
- 715
- 716
- 717
- 718
- 719
- 720
- 721
- 722
- 723
- 724
- 725
- 726
- 727
- 728
- 729
- 730
- 2020 The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* 62(4):725–757. DOI:10.1017/RDC.2020.41.
- Reitan, Gaute, and Inger Marie Berg-Hansen
- 2009 *Lundevågenprosjektet, delrapport 1. Sammenfattende rapport. Lunde, 6/1, 6/35 og Skjolnes 7/23, 7/27, Farsund kommune, Vest-Agder*. Oslo.
- Reitan, Gaute, and Per Persson (editors)
- 2014 *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og Porsgrunn. Bind 2. Seinmesolittiske, neolittiske og yngre lokaliteter i Vestfold og Telemark*. Portal forlag, Kristiansand.
- Reitan, Gaute, and Lars Sundström (editors)
- 2018 *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*. Cappelen Damm Akademisk, Oslo.
- Roalkvam, Isak
- 2020 Algorithmic classification and statistical modelling of coastal settlement patterns in Mesolithic south-eastern Norway. *Journal of Computer Applications in Archaeology* 3(1):288–307. DOI:<https://doi.org/10.5334/jcaa.60>.
- 2022 Exploring the composition of lithic assemblages in Mesolithic south-eastern Norway. *Journal of Archaeological Science: Reports* 42:103371. DOI:10.1016/j.jasrep.2022.103371.
- Røberg, Frank Halvar N.
- 2012 *Bosettings- og aktivitetsspor. Larønningen, 221/2138. Skien, Telemark*. University of Oslo, Museum of Cultural History, Oslo.
- Romundset, Anders
- 2018 Postglacial shoreline displacement in the Tvedstrand-Arendal area. In *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*, edited by Gaute Reitan and Lars Sundström, pp. 463–478. Cappelen Damm Akademisk, Oslo.

- 731 Romundset, Anders, Stein Bondevik, and Ole Bennike
732 2011 Postglacial uplift and relative sea level changes in Finnmark, northern Norway. *Quaternary Science Reviews* 30(19-20):2398–2421. DOI:10.1016/j.quascirev.2011.06.007.
733
- 734 Romundset, Anders, Ola Fredin, and Fredrik Høgaas
735 2015 A Holocene sea-level curve and revised isobase map based on isolation basins from near the southern tip of Norway. *Boreas* 44:383–400. DOI:10.1111/bor.12105.
736
- 737 Romundset, Anders, Thomas R. Lakeman, and Fredrik Høgaas
738 2018 Quantifying variable rates of postglacial relative sea level fall from a cluster of 24 isolation basins in southern Norway. *Quaternary Science Reviews* 197:175–192. DOI:10.1016/j.quascirev.2018.07.041.
739
- 740 2019 Coastal lake records add constraints to the age and magnitude of the Younger Dryas ice-front oscillation along the Skagerrak coastline in southern Norway. *Journal of Quaternary Science* 34(2):112–124. DOI:<https://doi.org/10.1002/jqs.3085>.
- 741
- 742 Schmitt, Lou, Stephan Larsson, Jan Burdukiewicz, John Ziker, Krister Svedhage, Jeanette Zamon, and Steffen Holger
743 2009 Chronological Insights, Cultural Change, and Resource Exploitation on the West Coast of Sweden During the Late Palaeolithic/Early Mesolithic Transition. *Oxford Journal of Archaeology* 28:1–27. DOI:10.1111/j.1468-0092.2008.00317.x.
- 744
- 745 Schülke, Almut
746 2020 First visit or revisit? Motivations of mobility and the use and reuse of sites in the changing coastal areas of Mesolithic southeastern Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement with the Coast from the Atlantic to the Baltic Sea*, edited by Almut Schülke, pp. 359–393. Routledge, London & New York.
- 747
- 748 Shennan, Ian
749 2015 Handbook of sea-level research: Framing research questions. In *Handbook of Sea-Level Research*, pp. 3–25. Wiley, Chichester.
- 750
- 751 Shetelig, Haakon
752 1922 *Primitive Tider i Norge – En oversikt over stenalderen*. John Griegs Forlag, Bergen.
- 753
- 754
- 755 Sognnes, Kalle
756 2003 On shoreline dating of rock art. *Acta Archaeologica* 74:189–209.
- 757
- 758 Solheim, Steinar
759 2012 Lokal praksis og fremmed opphav. Arbeidsdeling, sosiale relasjoner og differensiering i østnorsk tidligeolitikum. Unpublished PhD thesis, Oslo.
- 760
- 761 (editor)
762 2017 *E18 Rugsvedt-Dørdal. Arkeologiske undersøkelser av lokaliteter fra steinalder og jernalder i Bamble kommune, Telemark fylke*. Portal forlag, Kristiansand.
- 763
- 764 2020 Mesolithic coastal landscapes. Demography, settlement patterns and subsistence economy in southeastern Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement with the Coast from the Atlantic to the Baltic Sea*, edited by Almut Schülke. Routledge, London & New York.
- 765
- 766 2021 Timing the Emergence and Development of Arable Farming in Southeastern Norway by Using Summed Probability Distribution of Radiocarbon Dates and a Bayesian Age Model. *Radiocarbon*:1–22. DOI:10.1017/RDC.2021.80.
- 767
- 768 Solheim, Steinar, and Per Persson
769 2018 Early and mid-Holocene coastal settlement and demography in southeastern Norway: Comparing distribution of radiocarbon dates and shoreline-dated sites, 8500–2000 cal. BCE. *Journal of Archaeological Science: Reports*:334–343. DOI:10.1016/j.jasrep.2018.03.007.
- 770
- 771 Sørensen, Rolf
772 1979 Late Weichselian deglaciation in the Oslofjord area, south Norway. *Boreas* 8(2):241–246. DOI:<https://doi.org/10.1111/j.1502-3885.1979.tb00806.x>.
- 773

- 774 1999 En ^{14}C datert og dendrokronologisk kalibrert strandforksyvningskurve for søndre Østfold. Sørøst-Norge.
In *Museumslandskap. Artikkelsamling til Kerstin Griffin på 60-årsdagen. Bind A*, edited by Lotte
Selsing and Grete Lillehammer, pp. 227–242. AmS-rapport 12A. Museum of Archaeology, Stavanger.
- 775
- 776 Sørensen, Rolf, Kari E. Henningsmoen, Helge I. Høeg, and Veronika Gälman
- 777 2014 Holocene landhevningsstudier i søndre Vestfold og sørøstre Telemark – Revidert kurve. In *Vestfold-
baneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og Porsgrunn.
Bind 1*, edited by Stine Melvold and Per Persson, pp. 36–47. Portal, Kristiansand.
- 778
- 779 Sørensen, Rolf, Kari E Henningsmoen, Helge I Høeg, and Veronika Gälman
- 780 in prep Holocen vegetasjonshistorie og landhevning i søndre Vestfold og sørøstre Telemark. In *The Stone
Age in Telemark. Archaeological results and scientific analysis from Vestfoldbaneprosjektet and E18
Rugtvedt-Dørdal*, edited by Per Persson and Steinar Solheim.
- 781
- 782 Sørensen, Rolf, Helge I. Høeg, Kari E. Henningsmoen, Göran Skog, Solveig F. Labowsky, and Bjørg Stabell
- 783 2014 Utviklingen av det senglasiale og tidlig preboreale landskapet og vegetasjonen omkring steinalder-
boplassene ved Pauler. In *E18 Brunlanesprosjektet. Bind I. Forutsetninger og kulturhistorisk sam-
menstilling*, edited by Lasse Jakslund and Per Persson, pp. 171–213. University of Oslo, Museum of
Cultural History, Oslo.
- 784
- 785 Stokke, Jo-Simon Frøshaug, and Gaute Reitan
- 786 2018 Kvastad A2. Lokalitet med funn fra tidlig- og mellommesolitikum og dyrkningsspor fra mellom- og
senneolitikum. In *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*, edited by
Gaute Reitan and Lars Sundström, pp. 375–407. Cappelen Damm Akademisk, Oslo.
- 787
- 788 Stroeven, Arjen P., Clas Hättestrand, Johan Kleman, Jakob Heyman, Derek Fabel, Ola Fredin, Bradley W.
789 Goodfellow, Jonathan M. Harbor, John D. Jansen, Lars Olsen, Marc W. Caffee, David Fink, Jan Lundqvist,
790 Gunhild C. Rosqvist, Bo Strömberg, and Krister N. Jansson
- 791 2016 Deglaciation of Fennoscandia. *Quaternary Science Reviews* 147:91–121.
DOI:10.1016/j.quascirev.2015.09.016.
- 792
- 793 Svendsen, John Inge, and Jan Mangerud
- 794 1987 Late Weichselian and Holocene sea-level history for a cross-section of western Norway. *Journal of
Quaternary Science* 2(2):113–132. DOI:10.1002/jqs.3390020205.
- 795
- 796 Tallavaara, Miikka, and Petro Pesonen
- 797 2020 Human ecodynamics in the north-west coast of Finland 10,000–2000 years ago. *Quaternary Interna-
tional* 549:26–35. DOI:10.1016/j.quaint.2018.06.032.
- 798
- 799 Wang, Ian
- 800 2019 *topoDistance: Calculating topographic paths and distances. R package version 1.0.1.* [Https://CRAN.R-
project.org/package=topoDistance](https://CRAN.R-project.org/package=topoDistance).
- 801
- 802 Wenn, Camilla Cecilie
- 803 2012 *Bosettingsspor, produksjonsområde og dyrkningsspor fra Neolitikum til Folkevandringstid. Bratsberg,
63/69, 244. Skien kommune, Telemark.* University of Oslo, Museum of Cultural History, Oslo.
- 804
- 805 Wikell, Roger, Fredrik Molin, and Mattias Pettersson
- 806 2009 The Archipelago of Eastern Middle Sweden - Mesolithic Settlement in Comparison with ^{14}C and
Shoreline Dating. In *Chronology and Evolution within the Mesolithic of North-West Europe*, edited
by Philippe Crombé, Mark van Strydonck, Joris Sergant, Mathieu Boudin, and Machteld Bats, pp.
417–434. Cambridge Scholar Publishing, Brussels.
- 807
- 808 Yubero-Gómez, María, Xavier Rubio-Campillo, and Javier López-Cachero
- 809 2016 The study of spatiotemporal patterns integrating temporal uncertainty in late prehistoric settlements
in northeastern Spain. *Archaeological and Anthropological Sciences* 8(3):477–490. DOI:10.1007/s12520-
015-0231-x.
- 810