
1 A simulation-based assessment of the relation between Stone Age
2 sites and relative sea-level change along the Norwegian Skagerrak
3 coast

4 Isak Roalkvam^{1,*}

5 11 April, 2022

6 ¹ University of Oslo, Institute of Archaeology, Conservation and History

7 * Correspondence: Isak Roalkvam <isak.roalkvam@iakh.uio.no>

8 **1 Introduction**

9 The post-glacial relative sea-level fall that characterises large areas of Fennoscandia is fundamental to
10 its archaeology. This follows not only from the dramatic changes to the landscape that this process has
11 represented throughout prehistory, but also from the fact that if archaeological phenomena were situated
12 close to the contemporary shoreline when they were in use, a reconstruction of the trajectory of shoreline
13 displacement can be used to date these phenomena based on their altitude relative to the present day sea-level.
14 This method, also called shoreline dating, has long history of use in the region and is frequently applied
15 to assign an approximate date to diverse archaeological phenomena such as rock art, grave cairns, various
16 harbour and sea-side constructions and, as is the focus of this study, Stone Age sites (e.g. Åkerlund 1996;
17 Bjerck 2005; Nordqvist 1995; Schmitt et al. 2009; Sognnes 2003; Tallavaara and Pesonen 2020; Wikell et al.
18 2009).

19 The close association between Stone Age settlements in the northern parts of Scandinavia and shifting
20 prehistoric shorelines was established by the end of the 19th century (De Geer 1896), and was first applied
21 as a dating method at the turn of the century (Brøgger 1905; Hollender 1901). Shoreline dating has been
22 fundamental to Norwegian Stone Age archaeology ever since (e.g. Berg-Hansen 2009; Bjerck 1990, 2008a;
23 Breivik 2014; Johansen 1963; Mikkelsen 1975; Mjærum 2022; Nummedal 1923; Shetelig 1922; Solheim and
24 Persson 2018). The method is used both independently, and to compliment other sources of temporal
25 data such as typological indicators or radiometric dates. However, given the coarse and fuzzy resolution
26 of established typological frameworks, the vast amount of surveyed sites that only contain generic lithic
27debitage that could hail from any part of the period, and as the conditions for the preservation of organic
28 material is typically poor in Norway, dating with reference to shoreline displacement is often the only and
29 most precise method by which one can hope to date the sites. Shoreline dating is consequently fundamental to
30 our understanding of the Norwegian Stone Age. This is both because it is central to the temporal framework
31 on which our understanding of the period is based, but also because the method is only applicable so long as
32 the societies in question have continuously settled on or close to the contemporary shoreline. Consequently,
33 adherence or deviation from this pattern also has major implications for the socio-economic foundations of
34 the societies in question.

35 Despite its important role for Norwegian Stone Age archaeology, the applicability of dating by reference to
36 shoreline displacement has only been evaluated using relatively coarse methods. The aim of this paper is to
37 provide a systematic and comprehensive review of the degree to which radiocarbon dates correspond with the
38 dates informed by our current knowledge of shoreline displacement in a larger area of south-eastern Norway,
39 using a more refined methodological approach. The goal is to quantify the degree to which the assumption

40 of shore-bound settlement holds through the Stone Age, and in turn have this inform an improved method
41 for shoreline dating. As presented in more detail below, this problem involves the combined evaluation of
42 three major analytical dimensions. One is the questions of when the sites were in use, the second pertains to
43 the reconstruction of the contemporaneous sea-level, and the third follows from the fact that the relation
44 between site and shoreline is inherently spatial. Taking inspiration from studies that have integrated various
45 sources of spatio-temporal uncertainty through Monte Carlo simulation (e.g. Bevan et al. 2013; Crema et al.
46 2010; Crema 2012, 2015; Yubero-Gómez et al. 2016), a similar approach is adopted here.

47 2 Background

48 Relative sea-level (RSL) can be defined as the mean elevation of the surface of the sea relative to land, or,
49 more formally, the difference in elevation between the geoid and the surface of the Earth as measured from the
50 Earth's centre (Shennan 2015). Variation in this relative distance follow from a range of effects (e.g. Milne et
51 al. 2009). Of central importance here is eustasy and istostasy. The eustatic sea-level is understood as the
52 sea-level if the water has been evenly distributed across the Earth's surface without adjusting for variation in
53 the rigidity of the Earth, its rotation, or the self-gravitation inherent to the water body itself (Shennan 2015).
54 The eustatic sea-level is mainly impacted by glaciation and de-glaciation, which can bind or release large
55 amounts of water into the oceans (Mörner 1976). Istostasy, on the other hand, pertains to adjustments in the
56 crust to regain gravitational equilibrium relative to the underlying viscous mantle. This can be the result of
57 glacial istostasy, which follows from glaciation and de-glaciation and corresponding loading and unloading of
58 weight, as well as from erosion of the crust, which causes its weight to be redistributed. These effects thus
59 causes the lithosphere to either subside due to increased weight, or to rebound and lift upwards due to lower
60 weight (Milne 2015).

61 Following the end of the Weichselian and the final retreat of the Fennoscandian Ice Sheet (e.g. Hughes et al.
62 2016; Stroeven et al. 2016, see Figure 1), the isostatic rebound has been so severe that most areas of Norway
63 have been subject to a continuous relative sea-level regression, despite corresponding eustatic sea-level rise
64 (e.g. Mörner 1979; Svendsen and Mangerud 1987). In other words, the RSL has been dropping throughout
65 prehistory. As this process is the result of glacial loading, the rate of uplift is more severe towards the centre
66 of the ice sheet. Thus, some areas on the outer coast have had a more stable RSL or been subject to marine
67 transgression (e.g. Romundset et al. 2015; Svendsen and Mangerud 1987). These conditions are directly
68 reflected in the archaeological record. In areas where the sea-level has been stable over longer periods of
69 time, people have often reused coastal site locations multiple times and over long time-spans, creating a
70 mix of settlement phases that are difficult to disentangle (e.g. Hagen 1963; Reitan and Berg-Hansen 2009).
71 Transgression phases, on the other hand, can lead to complete destruction of the sites, bury them in marine
72 sediments, or in the outermost periphery, leave them still submerged today (Bjerck 2008a; Glørstad et al.
73 2020). This can lead to a hiatus in the archaeological record for certain sub-phases in the impacted areas.
74 Comparatively, given a continuous and still ongoing shoreline regression from as high as c. 220 m above
75 present sea-level in the inner Oslo fjord, any one location in south-eastern Norway has only been shore-bound
76 within a relatively limited time-span, and the sites have not been impacted by any transgressions (Hafsten
77 1957, 1983; Romundset et al. 2018; Sørensen 1979). This makes the region especially useful for evaluating
78 the assumption of a shore-bound settlement pattern over a long and continuous time-span.

79 The method of shoreline dating has been met with scepticism as related to the fundamental premise that
80 most sites would have been consistently shore-bound, been characterised as a relative dating method for sites
81 located at different elevations within a constrained geographical area, or been argued to offer no more than
82 a earliest possible date for when a site could have been in use (see review by Nordqvist 1999). The most
83 common application in Norway has arguably been to use shoreline dating to provide an approximate date for
84 the occupation of the sites, often in combination with other dating methods (see for example chapters in
85 Jakobsland 2014; Melvold and Persson 2014; Reitan and Persson 2014; Reitan and Sundström 2018; Solheim
86 2017 and below). Recently the method has also been used independently to date a larger number sites to get
87 a general impression of site frequency over time. This is done by aggregating point estimates of shoreline
88 dates in 100, 200 or 500 year bins (Breivik 2014; Breivik and Bjerck 2018; Fossum 2020; Nielsen 2021; Solheim




Figure 1: Deglaciation at 1000 year intervals from c. 17–8 kyr BCE. The study area defined later in the text is marked with a red outline (deglaciation data from Hughes et al. 2016, but see also Romundset et al. 2019 in relation to the study area).

89 and Persson 2018; see also Jørgensen et al. 2020; Tallavaara and Pesonen 2020). In his review, Nordqvist
90 (1999) argues that there can be little doubt concerning the general applicability of the method – what is less
91 clear is the level of reliability and chronological resolution that it can offer (see also Johansen 1963).

92 The shore-bound settlement location of prehistoric hunter-fisher-gatherers in Norway is generally believed
93 to follow both from the exploitation of aquatic resources and from movement and communication, which
94 would have been efficient on waterways (Bjerck 1990; Brøgger 1905:166; also discussed in Berg-Hansen 2009;
95 Bergsvik 2009). The same logic has also been extended to the hinter- and inland regions, where sites are
96 to be predominantly located along rivers and lakes (Brøgger 1905:166; Glørstad 2010:57–87; but see also
97 Gundersen 2013; Mjærum 2018; Schülke 2020). This is to take a dramatic turn at the transition to the
98 Late Neolithic, around 2400 BCE, with the introduction of the Neolithic proper (Prescott 2020; Solheim
99 2021). The introduction of a comprehensive Neolithic cultural package, including a shift to agro-pastoralism
100 and the introduction of the farm is to have led site locations to be more withdrawn from the shoreline (e.g.
101 Bakka and Kaland 1971; Østmo 2008:223; **prescott2012?**). That is not to say that waterways and aquatic
102 resources were no longer exploited, but rather that these activities would not have been as tightly integrated
103 with settlement and tool-production areas as in preceding periods (Glørstad 2012). At an earlier stage, at
104 the transition to the Early Neolithic (c. 3900 BCE), pottery is introduced to the sites, and there are some
105 indications of an initial uptake of agriculture at some sites in the Oslo fjord region. However, this appears
106 to be small in scale and is believed to be combined with a continued and predominantly hunter-gatherer
107 life-way, possibly followed by a complete de-Neolithisation in the Middle Neolithic (Hinsch 1955; Nielsen et
108 al. 2019; Østmo 1988:225–227). Nielsen (2021) has recently argued that the initial uptake of agriculture in
109 Early Neolithic south-eastern Norway is combined with a more complex settlement pattern, and that a simple
110 foraging/agricultural dichotomy would underplay the variation present in the Early and Middle Neolithic
111 settlement data (see also e.g. Amundsen et al. 2006; Østmo 1988; Solheim 2012:74). Seen in relation to
112 the question of interest here, the empirical expectation for the above outlined development would thus be
113 a predominantly shore-bound settlement in the Mesolithic, possibly followed by a more varied association
114 between sites and the shore-line with the transition to the Early Neolithic around 3900 BCE, and finally a
115 decisive shift with the Late Neolithic c. 2400 BCE.

116 Based on the generally accepted premise that most pre-Late Neolithic sites in south-eastern Norway located
117 lower than the marine limit were situated on or close to the contemporaneous shoreline, it is common to err
118 on the side of a shore-bound site location unless there is strong evidence to suggest otherwise. This is for
119 example reflected in survey projects, which are often guided by both a digital and mental reconstruction
120 of past sea-levels (e.g. Berg-Hansen 2009; Eskeland 2017). Similarly, following an excavation, if typological
121 indicators in the assemblages correspond with available shoreline displacement curves, a shore-bound site
122 location is often assumed, even if the typologically informed date-span is too wide to decisively verify this.
123 It is also common to combine this with a qualitative consideration of the landscape surrounding the sites,
124 and an evaluation of the degree to which the site location would have been sensible if the site was not shore
125 bound (e.g. Jakslund 2014; Johansen 1963; Nummedal 1923). This can for example pertain to accessibility. If
126 the site is situated on a ledge in a steep and jagged area of the present day landscape it would make intuitive
127 sense that the site was in use when the ocean reached closer to its elevation, as the site would have been
128 accessible by means of watercraft. Although it appears that the arguments for such site locations are sensible
129 and can for the most part be assumed to hold, comprehensive evaluations and attempts at quantification of
130 this tendency are relatively few (see also Ilves and Darmark 2011).

131 One of the more extensive evaluations of the relationship between archaeological radiocarbon dates and
132 RSL-change was done by Solheim and colleagues (Breivik et al. 2018; Solheim 2020), who compared 102
133 radiocarbon dates from 33 Mesolithic sites on the western side of the Oslo fjord to the displacement curve
134 for the Larvik area. They found an overlap between the probability density of the radiocarbon dates with
135 the shoreline displacement curve for 86.5% of the sites. However, where there was a discrepancy, the main
136 occupation of the sites are still believed to have been shore-bound rather than associated with the deviating
137 ^{14}C -dates. This is based on typological and technological characteristics of the assemblages. Whether these
138 mismatches represent later shorter visits that are responsible for the younger radiocarbon dates, or whether
139 these dates are entirely erroneous can be difficult to evaluate (e.g. Persson 2008; Schülke 2020). However,
140 this distinction is not deemed critical here, as what is of interest is settlements and tool-production areas as

141 evidenced by artefact inventories or multiple site features. Not remnants of stays as ephemeral to only be
142 discernible by individual features or dubious ^{14}C -dates. The evaluation of the relevance of radiocarbon dates
143 to settlement activity will here therefore be entirely dependent on, and follow the discretion of the original
144 excavation reports.

145 Other previous evaluations of the correspondence between radiocarbon- and RSL-informed dates have typically
146 followed the same structure as that of Breivik et al. (2018), involving a visual inspection of radiocarbon
147 probability density functions plotted against local shoreline displacement curves based on the elevation of the
148 site (e.g. Åkerlund et al. 1995; Åstveit 2018; Solheim 2020; see also Bjerck 2008b; Kleppe 1985; Ramstad
149 2009). This approach has a couple of limitations. First of all, the displacement curves are sometimes applied
150 directly to larger study areas, with only some studies having taken the variable uplift-rates into account when
151 performing this comparison (e.g. Åstveit 2018; Fossum 2020; Møller 1987; Persson 2008). Secondly, with this
152 method, the wider the uncertainty range associated with either radiocarbon date or displacement curve, the
153 higher the probability that the confidence intervals overlap, and the higher the probability that we conclude
154 in favour of our hypothesis. This thus leads to an inferential framework that favours uncertainty, which is
155 hardly desirable. In statistical terms this follows from the fact that while one cannot conclude that two dates
156 are different if their confidence intervals overlap, this does not necessarily mean that they are the same. The
157 question thus necessitates a flip from a null-hypothesis of no significant difference, to one of equivalence (e.g.
158 Lakens et al. 2018), as the question of interest is effectively one of synchronicity between events (cf. Parnell
159 et al. 2008). Another limitation of this often-employed method is that it only takes into account the vertical
160 distance between the sites and the sea-level. While this is the main parameter of interest for shoreline dating,
161 the practical implications of a vertical difference in RSL will be highly dependent on local topography and
162 bathymetry. RSL-change can have more dramatic consequences in a landscape characterised by a low relief,
163 as the horizontal displacement of the shoreline will be greater. Taking the spatial nature of the relationship
164 between site and shoreline into account will consequently help get more directly at the behavioural dimension
165 of this relation, and help move the analysis beyond a purely instrumental consideration of the applicability of
166 shoreline dating.

167 3 Data

168 To get at the relationship between sites and the contemporaneous shoreline, this analysis was dependent
169 on a study area with good control of the trajectory of prehistoric shoreline displacement. While there is
170 displacement data available for other areas of south-eastern Norway (e.g. Hafsten 1957; Sørensen 1979, 1999;
171 and recent compilation by Creel et al. 2022), considerable methodological developments in recent years means
172 that the most well-established displacement curves are from the region stretching from Horten county in
173 the north-east, to Arendal in the south-west. This area has newly compiled displacement curves for Horten
174 (**romundset2021?**), Larvik (Sørensen et al. in prep; Sørensen, Henningsmoen, et al. 2014; Sørensen, Høeg,
175 et al. 2014), Tvedstrand (Romundset 2018; Romundset et al. 2018), and Arendal (Romundset 2018).

176 The employed shoreline displacement data is based on the so-called isolation basin method (e.g. Kjemperud
177 1986; Romundset et al. 2011), which involves extracting cores from a series of basins situated on bedrock at
178 different elevations beneath the marine limit, and dating the transition from marine to lacustrine sediments.
179 Each basin thus represent a high precision sea-level index point (SLIP) which are combined using what has
180 been termed the isobase method to devise a continuous time series for RSL-change, projected to a common
181 isobase (see Creel et al. 2022:5). Furthermore, to minimise the impact of variable uplift rates, the basins
182 are located in a as constrained area of the landscape as possible. Following from the morphology of the
183 retreating ice sheet, the uplift is more severe towards the north-east, meaning that this needs to be adjusted
184 for in the case that any basins are located any significant distance from the common isobase perpendicular to
185 this gradient (Figure 2). The SLIPs indicate the isolation of the basins from the highest astronomical tide,
186 which is adjusted to mean sea-level in the compilation of the displacement curves based on the present day
187 tidal range. This is assumed to have been the same throughout the Holocene (Sørensen, Henningsmoen, et
188 al. 2014:44). Furthermore, the confidence bands of the displacement curves and their trajectory are quite
189 complex constructs, and are the integrated result of both expert knowledge and more objectively quantifiable

parameters. The reason for this is in part that the curves do not only contain uncertainty as related to radiometric dates, which are well defined, but also hold potential error as related to the interpretation and analysis of sediment cores, the nature and condition of the basin outlets and the adjustment to a common isobase, to name but a few (e.g. Romundset et al. 2011, 2019; for an alternative approach see Creel et al. 2022). For more details and evaluations done for the compilation of each curve, the reader is therefore referred to the individual publications.

The archaeological data compiled for the analysis consists of excavated Stone Age sites with available spatial data from the coastal region between Horten county in the north-east, to Arendal in the south-west (Figure 2). These number 155 sites. Of these, 91 sites are associated with a total of 547 radiocarbon dates. Of these, in turn, 67 sites are related to the 259 radiocarbon dates that fall within the Stone Age (9500–1700 BCE), with 95% probability. These sites and ^{14}C -dates form the basis for the analysis. Spatial data in the form of site limits and features, as defined by the excavating archaeologists, were retrieved from local databases at the Museum of Cultural History—the institution responsible for archaeological excavations in the region. In the compiled dataset, each radiocarbon date has been associated with the site features or excavation unit from where they originate, or, where these weren't available, the spatial limit of the entire site. Due to somewhat variable practices between excavations, what available spatial geometry best represents the site limit was decided based on an evaluation of the excavation reports. This means that the limits are variably given as that defined during initial survey, area de-turfed before excavation, area stripped with excavator following the excavation, manually excavated area, or convex hull polygons generated around the site features.

Three of the sites have been associated with agriculture, either directly or in the form building structures. The first is Nordby 1 at which the ^{14}C -dates are associated with a Late Neolithic long-house (Gjerpe and Bukkemoen 2008). The Middle Neolithic phase at Kvastad A2 (Stokke and Reitan 2018) and Late Neolithic phase at Nauen A (Persson 2008) are both directly related to farming acitivities. Both of these sites also have radiocarbon dates and lithic inventory associated with Mesolithic forager activities. Following from the expected deviance from the settlement patterns that are to characterise forager sites, these agricultural phases are highlighted in the analysis below. Finally, Nielsen (2021) has recently suggested that Early and Middle Neolithic features from the otherwise younger sites Bratsberg (Wenn 2012) and Larønningen (Røberg 2012) could be related to early agricultural activity in the Oslo fjord region. Due to the uncertain and somewhat speculative nature of this suggestion, these are omitted here.

The elevation data used for the analysis is a digital terrain model (DTM) freely available from the Norwegian Mapping Authority (Norwegian Mapping Authority 2018, <https://hoydedata.no>). It was here opted for the 10m resolution DTM rather than the higher-resolution 1m version. In addition to resulting in considerably less processing time, the higher resolution elevation model is more vulnerable to smaller-scale modern disturbances that the 10m version is not impacted by. The 10m resolution DTM of the study area is a down-sampled version of the 1m version and has a height accuracy with a systematic error of 0.1m (Norwegian Mapping Authority 2018). All data and R programming code (R Core Team 2021) required to run the analyses, as well as the derived data are freely available in an online repository at <https://osf.io/7f9su/>, organised as a digital research compendium following Marwick et al. (2018).

4 Methods

The method of shoreline dating is based on the spatial relationship between two phenomena, occupation of sites and shoreline displacement, each associated with their own range of temporal uncertainty. The first task was therefore to ascribe likely date ranges and associated uncertainty to these dimensions. To take account of the gradient in the isostatic rebound, the trajectory of shoreline displacement was first interpolated to each site location based on the distance to the isobases of the displacement curves using inverse distance weighting (e.g. Conolly 2020; Conolly and Lake 2006:94–97). This was done for each year along the entirety of the curves, weighting the interpolation by the standard squared inverse of the distances. The result of this process is shown for an example site in Figure 3. For the date ranges associated with the sites, all radiocarbon dates were first individually calibrated using the IntCal20 calibration curve (Reimer et al. 2020) using OxCal v4.4.4 (Bronk Ramsey 2009) through the oxcAAR package for R (Hinz et al. 2021). Radiocarbon dates associated




Figure 2: A) Distribution of the 67 analysed sites relative to the isobases of the displacement curves. The isobases have a direction of 327° (Romundset et al. 2018), B) Displacement curves. Note the increasing⁷ steepness of the curves towards the north-east.

239 with each site were then grouped if they overlapped with 99.7% probability, meaning these were effectively
240 taken to represent the same event, here termed settlement phases. In the case where there are multiple dates
241 believed to belong to a single settlement phase, these were modelled using the Boundary function in OxCal
242 and then summed. Multiple phases at a single site were treated as independent of each other.

243 The excavation of archaeological sites typically follow from residential and commercial development, as well as
244 the expansion of infrastructure. As the data collection for the DTM was started by the Norwegian Mapping
245 Authority in 2016, the area of the DTM immediately surrounding the sites has therefore sometimes been
246 severely impacted by disturbances after the excavation of the site. In addition to employing 10m resolution
247 DTM to alleviate some of these issues, this also necessitated some additional editing of the elevation raster.
248 This involved manually defining the extent of problem areas such as railways, highways, quarries and the like.
249 The DTM values on these were then set to missing, and new elevation values were interpolated from the
250 surrounding terrain. This was done using regularised spline interpolation with tension (e.g. Conolly 2020),
251 using the default settings of r.fillnulls from GRASS GIS (GRASS Development Team 2017) in R through the
252 package rgrass7 (Bivand 2021). In addition to code and original spatial data being available in the digital
253 research compendium for this paper, the analysis of each individual site is presented in the supplementary
254 material where it has been noted when the area surrounding a site has been edited in this manner.

255 Armed with a likely date range for the occupation(s) of each site, an estimated trajectory of relative sea-level
256 change at that location, and a DTM edited to remove substantial modern disturbances, the simulations were
257 performed. A single simulation run involved first drawing a single year from the posterior density estimate of
258 a given occupation phase of a site (Figure 4). This year then has a corresponding likely elevation range for
259 the contemporaneous shoreline from which an elevation value was drawn uniformly, using intervals of 5cm.
260 The sea-level was then raised to this elevation on the DTM by defining all elevation values at or below this
261 altitude as missing. Polygons were then created from the resulting areas with missing values. The horizontal
262 distance was then found by measuring the shortest distance between site and sea polygons, and the vertical
263 distance by subtracting the elevation of the sea-level from the lowest elevation of the site polygon. The
264 topographic distance between site and sea was also found by measuring the distance while taking into account
265 the slope of the terrain on the DTM. This was done using the topoDistance package for R (Wang 2019).
266 The topographic distance was measured between the site polygon and the horizontally closest point on the
267 shoreline. This means that the distance is not necessarily measured as the closest topographic distance to the
268 shoreline, but rather as the shortest topographic path to the horizontally closest point on the shoreline. Not
269 finding the topographically closest point significantly reduced the computational cost of the analysis, and is
270 deemed unlikely to have a considerable impact on the results given the distances considered. The shortest
271 topographic path was found using the Moore neighbourhood of eight cells (e.g. Conolly and Lake 2006:253;
272 Herzog 2013). In the case where the sea-polygons intersects the site polygon, all distance measures were
273 set to zero. In the case that the sea-polygons completely contain the site, the horizontal and topographic
274 distance measures were made negative, and the vertical distance was instead measured to the highest point
275 on the site polygon. While it is safe to assume that an archaeological site was not occupied when it was
276 located beneath sea-level, a negative result can reflect the inherent uncertainty in this procedure, and might
277 also help identify discrepancies in displacement data or radiocarbon dates. Negative values were therefore
278 retained with the exception of for the sites Gunnarsrød 5 and Pjonkerød R1, where the negative values are
279 believed to result from modern disturbances in the DTM rather than the ^{14}C -dates or displacement curves
280 (see supplementary material for more details).

281 This process was repeated 1000 times for each phase for each site. The choice of 1000 simulation runs follows
282 from an evaluation of when the mean distances between site and shoreline converged when running 5000
283 iterations of the simulation on the site Hovland 5, available in the supplementary material (cf. Crema et al.
284 2010:1125). Hovland 5 was chosen for this evaluation as it has a fairly imprecise date and is located in area
285 of quite complex topography.




Figure 3: Example site Hegna vest 1 (Fossum 2017). A) Location of the site on the edited 10m resolution DTM. The red outline is the site limit. B) Radiocarbon dates associated with the site. Fill colour indicates what dates are assumed to belong to the same settlement phase. Multiple dates are modelled using the Boundary function in OxCal and then summed. The red outline indicates that the date does not match the typological indicators in the artefact assemblage of the site. C) The location of the site within the study area relative to isobases of the employed displacement curves. D) Displacement curve interpolated to the site location.




Figure 4: Example of a single simulation run on the site Hegna vest 1. A) The simulation starts by drawing a single year from the posterior density estimate. B) This then corresponds to an elevation range on the interpolated displacement curve. A single elevation is drawn uniformly from this range using 5cm intervals. C) The sea-level is then adjusted on the DTM to this elevation and the various distance measures are found. D) The numerical result of the simulation run.




Figure 5: The result of 1000 simulation runs for each of the two groups of dates on the site Hegna vest 1. The first column of plots shows the radiocarbon probability density function from where dates were drawn during simulation. The second column displays the result of simulating the raised sea-level 1000 times. The more opaque the colour, the more times the sea-level was simulated to that location. The third column shows violin plots of the different distance measures across all simulations.

286 5 Simulation results

- 287 Overall, as is indicated by the measures for central tendency and the almost solid line along the 0m mark on
288 the y-axes, the simulations show that the sites tend to have been situated close to the shoreline when they
289 were in use (Figure 6). Some of the sites are situated considerable distances from the shoreline when the dates
290 believed to be erroneous in the original reports are included (Figure 6A), but if one accepts the interpretation
291 that these do not date the main occupation of the sites, as is indicated by the artefact inventories, Figure
292 6B gives considerable support to the notion that the sites were in use when they were situated on or close
293 to the contemporaneous shoreline. The distances for some of the earliest sites appears somewhat high, but
294 this can likely be explained as the result of the steepness of the displacement curves for the earliest part of
295 the Holocene (Figure 2B), which leads the uncertainty of the ^{14}C -dates to give a wider possible elevation
296 range for the simulated sea-level. Another immediately striking result is the apparent deviation from the
297 shoreline towards the end of the Stone Age. From around 2500 BCE several sites are situated a considerable
298 distance from the shoreline, and while a couple remain horizontally and topographically close, most appear
299 to be elevated a considerable distance from the sea-level, as indicated on the plot for vertical distance. There
300 are also a couple of sites located some distance from the shoreline just after 4000 BCE. While the sample size
301 is limited, this would thus be in line with a development that sees an increase in settlements located in the
302 immediate inland around this time.
- 303 The negative values around 8000 BCE originate from the sites Løvås 1, 2 and 3. These are recently excavated,
304 well-dated sites situated in a relatively undisturbed area of the landscape. While there would be a danger of
305 circularity of having archaeological sites inform a reconstruction RSL-change, and in turn use these to evaluate
306 the degree of shore-bound settlement, the sites do clearly represent a upper limit for the sea-level, as they
307 would not have been in use when located under water. It could therefore seem that the Løvås sites represent
308 a case where the archaeological material indicates a slight discrepancy in the geological reconstruction of
309 shoreline displacement in the area.
- 310 Accepting that shoreline dating appears to loose utility around the transition to the Late Neolithic, as
311 indicated by the clear deviation in site location from the shoreline after this, the results for from Figure 6B is
312 given again in Figure 7A, excluding all simulation results younger than 2500 BCE. Furthermore, all negative
313 values have here been set to zero, under the assumption that these result from uncertainty or errors in the
314 data, and not actual site locations. The resulting best point estimate for the vertical distance between sites
315 and shoreline for the pre-Late Neolithic is given by the median at 4m, while 95% of the values fall within the
316 range 0–18m. That is, for 95% of the cases, the shoreline was simulated to be situated on or down to 18m
317 below the site location. While these values remain the same when only the Mesolithic dates are included
318 (Figure 7B), the mean and standard deviation is slightly constrained. Furthermore, while the median for
319 horizontal and topographic distance is only 10m across all plots in Figure 7, the variation in the statistics for
320 dispersion is greater, illustrating the point that minor variations in vertical distance can have substantial
321 consequences for these distance measures, depending on the surrounding topography.
- 322 An exponential function has been fit to the distributions for vertical distance using maximum likelihood
323 estimation (Figure 7). While it makes theoretical sense that a process of exponential decay explains this
324 relationship, it is also clear that this does not perfectly match the data. However, this can at least in part be
325 related to methodological factors, where the accumulation of distance-values on the 0m mark likely follow
326 from forcing negative values to zero, from the resolution of the spatial data, and from defining intersecting
327 sea- and site polygon as having a distance of zero. If one accepts this, the probability density function for
328 exponential decay can be used to characterise the vertical distance between sites and the shoreline:

$$f(x) = \lambda e^{-\lambda x} \quad (1)$$

- 329 Where λ is the decay ratio and x is the distance to the shoreline. This can then be combined with the
330 elevation of the site and the shoreline displacement data to provide a method for shoreline dating that takes
331 this distance into account:

332 Where x is [...] the site elevation, i is the trajectory for shoreline displacement, here given as a uniform
333 probability range between the upper and lower bound of the interpolated displacement curve.

334 Eq. (2) is used to shoreline date the same sites from where this relationship was derived (Figure 8), employing
335 $\lambda = 0.173$ for the decay ratio in (1). This is the ratio identified when considering all of the pre-Late Neolithic
336 simulation results (Figure 7A). For illustrative purposes the Late Neolithic sites were also subjected to the
337 method for shoreline dating. Following from having defined the distance between intersecting sea- and site
338 polygons as zero during simulations, the sites were dated using the mean elevation of the site polygons to
339 allow for some variation in elevation over the site limits. The synchronicity between radiocarbon and shoreline
340 dates was then evaluated using the method presented by Parnell et al. (2008, Figure 9). Here, 100,000 age
341 samples drawn from the probability density function of each shoreline date were subtracted from 100,000
342 age samples drawn from the corresponding modelled ^{14}C -dates. The resulting range of the 95% highest
343 density region (HDR, Hyndman 1996) was then checked to see if it crosses zero, in which case the dates are
344 considered to be in agreement. When excluding the earliest occupation phase at Gunnarsrød 5, the deviation
345 of which is to be expected based on the simulation results (see above), the shoreline date correspond to the
346 radiocarbon dates in 58 out of 68 cases (84%). Only including dates modelled to be older than 2500 BCE
347 with 95% probability, i.e. older than the Late Neolithic, improves this to 56 out of 61 cases (92%). When only
348 including dates older than 4000 BCE with 95% probability, i.e. only Mesolithic, the success rate is further
349 increased to 46/49 (94%). The three failed Mesolithic shoreline dates are from the early sites Langemyr and
350 Kvastad A2, with the likely implication that a lower decay ratio than what is used for characterising the
351 distance between site and shoreline for all sites in aggregate should be used for sites known to be from the
352 earliest part of the Mesolithic (see also Figure 6).

353 6 Re-dating previously shoreline dated sites

354 To further explore the implementation for shoreline dating presented above, excavated and shoreline dated
355 Stone Age sites within the study area where ^{14}C -dates are not available or are not believed to date the
356 main occupation of the sites have been subjected to the outlined approach (Figure 10). The resulting dates
357 are compared to those originally proposed in the excavation reports for the sites (the numerical results are
358 available in the supplementary material). To avoid issues with recent disturbances on the DTM, the sites
359 have been dated based on the mean of the altitudes provided in the report for each site.

360 The comparison with previously reported dates is an illustrative, but unfair exercise for a few reasons. First
361 of all the dates provided in the reports are typically stated to be a very rough estimate, and are sometimes
362 given as a point estimate with an undefined, but implied or explicit uncertainty range. Secondly, seeing
363 as these reports are from various dates in time, many are based on now outdated data on RSL-change.
364 Finally, they are sometimes only meant to indicate a lower bound for when the sites could have been in use.
365 Overall, the results suggest that shoreline dating has generally been applied with a fairly reasonable degree
366 of success, seeing as these dates have typically been interpreted and informed research in an approximate
367 manner (although see e.g. Roalkvam 2022). That being said, the results do also indicate that shoreline
368 dating has at times been applied with an exaggerated degree of precision. While the implications of a more
369 stable RSL-change for the duration of use and re-use of site locations are well known, this also appears to
370 be somewhat under-appreciated for the purposes of shoreline dating. The results also highlight the spatial
371 and temporal contingency of the method, illustrated by the variation in the range of the 95% HDRs for the
372 dates. In some cases the method provides a very precise date range and in others it offers little more than a
373 *terminus post quem*. This is dependent on the steepness of the displacement curves, leading to the general
374 pattern of older sites situated towards the north-east getting more precise dates (cf. Figure 2B). Further, as
375 some of the date ranges extend well beyond major chronological divisions, even into the Iron Age, they could
376 be severely and securely constrained with only cursory reference to typology. While this would be trivial in
377 some cases, the nature and uncertainty inherent to the method still means that this is arguably a required



A**B**

Figure 6: The result of running the analysis across all sites. Each data point is plotted with some transparency, meaning that the more intense the colour, the more often those values occurred. Results associated with agricultural activities are plotted in grey. The first row A) shows the result of including all dates to the Stone Age, including those seen as otherwise unrelated to the main occupation of the sites. The second row B) shows the result of excluding these. The table under each plot lists some corresponding statistics for central tendency and dispersion.



A**B**

Figure 7: Histograms showing the simulated distance from the shoreline using dates corresponding to the site inventories. Negative values have been set to zero. A) Simulated results older than 2500 BCE, and B) simulated results older than 4000 BCE.




Figure 8: The result of backwards shoreline dating the sites with radiocarbon dates corresponding to the artefact inventory using the method proposed here. The shoreline dates are plotted in grey and underlined¹⁶ with the 95% HDR in black. These are plotted against the modelled radiocarbon dates, which are given colour from oldest to youngest occupation phase for each site, defined by non-overlapping dates at 99.7% probability.




Figure 9: Evaluation of the agreement between the shoreline dates and radiocarbon dates given in Figure 8. When the range of the 95% HDR for age difference crosses zero, the shoreline and radiocarbon dates are considered synchronous. Line segments with vertical bars indicate that this does not cross zero and that the dates are not in agreement. The division and colour coding at the top of the plots reflect the division of site phases given in Figure 8.

378 exercise that should be explicitly performed. This also points to the possibility of drawing on other temporal
379 data, for example within a Bayesian framework, to further improve the precision of the dates that can be
380 achieved with shoreline dating.

381 Not least following from the fact that relatively few Preboreal ^{14}C -dates associated with anthropogenic
382 activity have been achieved in Norway (Åstveit 2018; Damlien and Solheim 2018; Kleppe 2018), the shoreline
383 dating of the earliest sites is essential for understanding the pioneer settlement and the initial colonisation of
384 the Scandinavian peninsula (e.g. Bang-Andersen 2012; Berg-Hansen 2018; Breivik 2014; Fuglestvedt 2012;
385 Glørstad 2016). The shoreline dated Preboreal sites from the Brunlanes-project are among the earliest known
386 sites in Norway (Jaksland 2012a, 2012b; Jaksland and Persson 2014). These have a distinct Early Mesolithic
387 artefact inventory and are situated in a steep area of the landscape where it would be difficult to envision use
388 of the sites after the sea retreated any significant distance from their location. In the original publication
389 of the sites, Jaksland (2014) provides a thorough discussion of shoreline dating in general, and as used for
390 the dating of the Brunlanes sites specifically. A comparison of his results and the ones achieved using the
391 above-outlined approach are given in Figure 11A. The sites have been dated using what Jaksland (2014) gives
392 as the lowest elevation of finds at each site, and by employing a exponential decay ratio of 0.13, to allow for
393 more deviance in the distance between site and shoreline. This corresponds to the decay ratio for sites older
394 than 7000 BCE in Figure 7.

395 The small discrepancies between the achieved results mainly follow from the fact that a slightly updated
396 version of the local displacement curve is applied here (cf. Sørensen et al. in prep). Jaksland's dates are
397 given a flat 200 and 50 year uncertainty range starting from what he gives as the earliest possible date. The
398 200 year uncertainty range is given if the sites were to be considered in isolation, while the argument for
399 the uncertainty range of only 50 years is based on the location of the sites relative to each other. Since
400 they are located in such a constrained and steep area of the landscape, the difference in elevation between
401 the sites is argued to establish their relative date and thus constrain the uncertainty ranges so that they
402 don't overlap. This information is not integrated in the approach outlined here, but could justify further
403 reducing the uncertainty ranges. Although their accuracy is of course ultimately dependent on the veracity of
404 the geological reconstruction, the high rate of RSL-change in this period does result in very precise dates.
405 Above it was suggested that additional temporal data could be combined with the method to improve its
406 accuracy and precision. This example, on the other hand, highlights the fact that the spatial nature of the
407 method means that a consideration of the surrounding terrain and other sites can also help in increasing the
408 precision of the method if this can be used to exclude certain sea-levels as unlikely for when a site was in use.
409 One approach could also be to assess the spatial implication of a proposed shoreline date by simulating the
410 adjusted sea-levels, as is done for Paurer 1 in Figure 11B, followed for example by a visual evaluation of the
411 topography or by evaluating the distance and steepness of the slope to the shoreline. If this is developed
412 further, it could conceivably be possible to exclude certain elevations as unlikely for the position of the
413 shoreline when the site was in use. Such approaches would make less of an impact in this setting, where the
414 95% HDR is already quite constrained, but could considerably improve the precision of the method in cases
415 where RSL-change has been less severe (cf. Figure 10).

416 7 Concluding remarks

417 The most immediate contribution of this paper is what must be considered a confirmation of previous research
418 into the relation between coastal Norwegian Stone Age sites and the prehistoric shoreline. This is indicated by
419 the close relationship between sites and the shoreline up until the transition to the Neolithic at c. 4000 BCE,
420 after which a couple of sites become situated some distance from the sea, followed by a more decisive break
421 at the transition to the Late Neolithic at c. 2500 BCE. This development is in agreement with the literature.
422 Furthermore, based on the quantitative nature of these findings, an initial formulation of a refined method
423 for the shoreline dating of pre-Late Neolithic Stone Age sites has been proposed. Apart from taking the
424 distance between sites and the isobases of the displacement curves into consideration when dating the sites,
425 this involves implementing Eq. (2) to account for the distance between the sites and the shoreline. When no
426 other information is available, it can at present be recommended to use the empirically derived exponential

$$\lambda = 0.173$$




Figure 10: Re-dating previously shoreline dated sites in the study area without radiocarbon dates or with radiocarbon dates that do not correspond to the artefact inventories. The 95% HDRs in grey are compared¹⁹ to the dates originally proposed by the excavation reports in red.




Figure 11: Shoreline dating of the Brunlanes sites using site altitudes provided by Jaksland (2014:tab.4). A) The result of applying the approach to shoreline dating outlined above. The shoreline date in grey is underlined with the 95% HDR in black. Dates provided by Jaksland (2014) are plotted in red. The box indicates a 50 year uncertainty range which in combination with the red line extends 200 years. B) Map showing the centroids of the Paurer sites and Sky 1. The sea-level has been simulated using the probability density associated with the shoreline date for Paurer 1 (see also map in Jaksland 2014:fig.12a). Paurer 1 is the red point.

decay ratio of 0.173 (Figure 10A) to characterise this relationship. Furthermore, while this remains to be formalised and explored further, it was also showed how the accuracy of the method can be improved by including more information, both with reference to the topographic location of the sites and other temporal data. As the precision of the method is both geographically and temporally contingent due to the trajectory of RSL-change, where older sites situated towards the north-east in the study area will get a more precise date than younger sites located towards the south-west, the impact of such additional information will also vary.

Future investigations and radiocarbon dates from Stone Age sites in the region can not only be used to further evaluate and adjust the findings reported here, but a larger sample size could also lay the foundations for refining the method by identifying subsets of sites for which the application of the method could be adjusted. Given it's behavioural nature, it would for example seem likely that dimensions such as the nature and purpose of visits to the sites will have implications for how close to the shoreline they were located. Furthermore, other dimensions related to the topographic location of the sites could be similarly explored. This for example pertains to the exposure of sites to wave action, which is likely to have been of concern (Roalkvam 2020), and which presumably has implications for how close to the shoreline people settled (Blankholm 2020; Helskog 1978). This is also related to the fact that while the mean sea-level is used for dating the sites, a consideration of the tidal range could possibly also have implications for the site location relative to the shoreline, depending on the topography (Helskog 1978). The potential of dimensions such as these was also hinted at here with the estimation and cursory treatment of the horizontal and topographic distance to the shoreline. If patterns related to such locational patterns can be discerned and unpicked, this will not least be useful for improving the shoreline dating of sites which have only been surveyed and where little information beyond their location is available.

Some limitations and sources of likely variation and uncertainty that have not been considered should also be mentioned. First of all the sample size is quite strained and the future addition more sites might alter the picture considerably. Secondly, the DTM has only been corrected for major modern disturbances. This means that other forms of erosion, although likely not that prevalent, has not been taken into account. Thirdly, the DTM has a vertical error which could also benefit from being integrated in the analysis (cf. Lewis 2021). Fourthly, the displacement curves were here interpolated to all site locations without accounting for increased uncertainty as one moves further away from the isobases of the displacement curves. This is also related to the fact that the RSL data can be handled in different ways than with the isobase method that has been used for the compilation of the employed displacement curves (cf. Creel et al. 2022). Fifthly, neither the question of how site limits are defined nor the elevation range over which these extend was given much consideration (cf. Mjærum 2022). Finally, the radiocarbon dates and division of settlement phases at each site was here simply done by treating radiocarbon dates not overlapping at 99.7% as representing unrelated occupations. This could also be handled differently (Bronk Ramsey 2009, 2015). While each of these factors will have variable impact on the final results, they clearly represent dimensions which would all benefit from further consideration.

Finally, this analysis employed a simulation approach to integrate multiple sources of spatio-temporal uncertainty. Here this was simply used to inform the question of the distance between sites and the shoreline. However, this method and general framework can be extended to a wide range of use-cases where one needs to visualise, and quantitatively or qualitatively evaluate the relationship between archaeological phenomena, the prehistoric shoreline, and the uncertainty inherent in this reconstruction.

469 8 References

- 470 Åkerlund, Agneta
471 1996 *Human responses to shore displacement: Living by the sea in Eastern Middle Sweden during the Stone*
472 *Age*. Riksantikvarieämbetet, Stockholm.
- 473 Åkerlund, Agneta, Jan Risberg, Urve Miller, and Per Gustafsson
474 1995 On the applicability of the ^{14}C method to interdisciplinary studies on shore displacement and settlement
475 location. *PACT* 49:53–84.
- 476 Amundsen, Øystein, Stig Knutsen, Axel Mjærum, and Gaute Reitan
477 2006 Nøkleby i Ski – en tidligeolittisk jordbruksboplass? *Primitive tider* 9:85–96.
- 478
- 479 Åstveit, Leif Inge
480 2018 The Early Mesolithic of Western Norway. In *Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 231–274.
481 Equinox, Sheffield.
- 482 Bakka, Egil, and Peter Emil Kaland
483 1971 Early farming in Hordaland, western Norway. Problems and approaches in archaeology and pollen
484 analysis. *Norwegian Archaeological Review* 4:1–17. DOI:10.1080/00293652.1971.9965136.
- 485 Bang-Andersen, Sveinung
486 2012 Colonizing Contrasting Landscapes. The Pioneer Coast Settlement and Inland Utilization in
487 Southern Norway 10,000–9500 Years Before Present. *Oxford Journal of Archaeology* 31:103–120.
488 DOI:10.1111/j.1468-0092.2012.00381.x.
- 489 Berg-Hansen, Inger Marie
490 2009 *Steinalderregistrering. Metodologi og forskningshistorie i Norge 1900–2000 med en feltstudie fra Lista i Vest-Agder*. Museum of Cultural History, University of Oslo, Oslo.
- 491 2018 Continuity and Change in Late- and Post-glacial Social Networks: Knowledge Transmission and
492 Blade Production Methods in Ahrensburgian and Early Mesolithic North West Europe. In *The Early Settlement of Northern Europe. Transmission of Knowledge and Culture*, edited by Kjel Knutsson, Helena Knutsson, Jan Apel, and Håkon Glørstad, pp. 63–98. Equinox, Sheffield.
- 493 Bergsvik, Knut Andreas
494 2009 Caught in the middle: functional and ideological aspects of Mesolithic shores in Norway. In *Mesolithic Horizons: Papers presented at the Seventh International Conference on the Mesolithic in Europe, Belfast 2005*, edited by Sinéad B. McCartan, Rick Schulting, Graeme Warren, and Peter Woodman, pp. 602–609. Oxbow Books, Oxford.
- 495 Bevan, Andrew, Enrico R. Crema, Xiuzhen Li, and Alessio Palmisano
496 2013 Intensities, Interactions, and Uncertainties: Some New Approaches to Archaeological Distributions.
497 In *Computational Approaches to Archaeological Spaces*, edited by Andrew Bevan and Mark Lake, pp.
498 27–52. Left Coast Press, Walnut Creek.
- 499 Bivand, Roger
500 2021 *rgrass7: Interface Between GRASS 7 Geographical Information System and R*. R package version
501 0.2-6.
- 502 Bjerck, Hein Bjartmann
503 1990 Mesolithic site types and settlement patterns at Vega, Northern Norway. *Acta Archaeologica* 60:1–32.
- 504
- 505 2005 Strandlinjedatering. In *Norsk arkeologisk leksikon*, edited by Einar Østmo and Lotte Hedeager, pp.
506 363–364. Pax, Oslo.
- 507 2008a Norwegian Mesolithic Trends: A Review. In *Mesolithic Europe*, edited by Geoff Bailey and Penny
508 Spikins, pp. 60–106. Cambridge University Press, Cambridge.
- 509 2008b Innledende betraktninger. In *NTNU Vitenskapsmuseets arkeologiske undersøkelser Ormen Lange Nyhamna*, edited by Hein Bjartmann Bjerck, Leif Inge Åstveit, Trond Meling, Jostein Gundersen, Guro Jørgensen, and Staale Normann, pp. 548–551. Tapir Akademisk Forlag, Trondheim.

- 511 Blankholm, Hans Peter
512 2020 In the wake of the wake. An investigation of the impact of the Storegga tsunami on the human
settlement of inner Varangerfjord, northern Norway. *Quaternary International* 549:65–73. DOI:<https://doi.org/10.1016/j.quaint.2018.05.050>.
- 513
514 Breivik, Heidi Mjelva
515 2014 Palaeo-oceanographic development and human adaptive strategies in the Pleistocene–Holocene transition:
516 A study from the Norwegian coast. *The Holocene* 24:1478–1490. DOI:10.1177/0959683614544061.
- 517 Breivik, Heidi Mjelva, Guro Fossum, and Steinar Solheim
518 2018 Exploring human responses to climatic fluctuations and environmental diversity: Two stories
from Mesolithic Norway. *Quaternary International* 465. Impacts of gradual and abrupt environmental changes on Late glacial to Middle Holocene cultural changes in Europe:258–275.
519 DOI:10.1016/j.quaint.2016.12.019.
- 520 Breivik, Heidi, and Hein Bjartmann Bjerck
521 2018 Early Mesolithic Central Norway: A Review of Research History, Settlements, and Tool Tradition. In
Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change,
522 edited by Hans Peter Blankholm, pp. 169–206. Equinox, Sheffield.
- 523 Brøgger, Waldemar Christofer
524 1905 *Strandliniens Beliggenhed under Stenalderen i Det Sydøstlige Norge*. Norges geologiske undersøkelse,
525 Kristiania.
- 526 Bronk Ramsey, Christopher
527 2009 Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* 51(1):337–360.
528 DOI:10.1017/S0033822200033865.
- 529 2015 Bayesian Approaches to the Building of Archaeological Chronologies. In *Mathematics and Archaeology*,
530 edited by Juan A. Barcelo and Igor Bogdanovic, pp. 272–292. CRC Press, Boca Raton.
- 531 Conolly, James
532 2020 Spatial interpolation. In *Archaeological Spatial Analysis: A Methodological Guide*, edited by Mark
Gillings, Piraye Hacıgüzeller, and Gary Lock, pp. 118–134. Routledge, London & New York.
- 533 Conolly, James, and Mark Lake
534 2006 *Geographical Information Systems in Archaeology*. Cambridge University Press, Cambridge.
- 535
536 Creel, Roger C., Jacqueline Austermann, Nicole S. Khan, William J. D’Andrea, Nicholas Balascio, Blake
537 Dyer, Erica Ashe, and William Menke
538 2022 Postglacial relative sea level change in Norway. *Quaternary Science Reviews* 282:107422.
539 DOI:10.1016/j.quascirev.2022.107422.
- 540 Crema, Enrico R.
541 2012 Modelling Temporal Uncertainty in Archaeological Analysis. *Journal of Archaeological Method and*
542 *Theory* 19(3):440–461. DOI:10.1007/s10816-011-9122-3.
- 543 2015 Time and Probabilistic Reasoning in Settlement Analysis. In *Mathematics and Archaeology*, edited by
544 Juan A. Barcelo and Igor Bogdanovic, pp. 314–334. CRC Press, Boca Raton.
- 545 Crema, Enrico R., Andrew Bevan, and Mark W. Lake
546 2010 A probabilistic framework for assessing spatio-temporal point patterns in the archaeological record.
547 *Journal of Archaeological Science* 37(5):1118–1130. DOI:10.1016/j.jas.2009.12.012.
- 548 Damlien, Hege, and Steinar Solheim
549 2018 The Pioneer Settlement of Eastern Norway. In *Early Economy and Settlement in Northern Europe. Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 335–367.
550 Equinox, Sheffield.
- 551 De Geer, Gerard
552 1896 *Om Skandinaviens geografiska utveckling efter Istiden*. P. A. Norstedt & Söner, Stockholm.
- 553
554 Eskeland, Knut Fossdal

- 556 2017 *Rapport, arkeologisk registrering. E18 Langangen Rugtvedt, 16/06999, Porsgrunn og Bamble kommune.*
557 Skien.
- 558 Fossum, Guro
- 559 2020 Specialists facing climate change. The 8200 cal BP event and its impact on the coastal settlement in
the inner Oslo fjord, southeast Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement*
with the Coast from the Atlantic to the Baltic Sea, edited by Almut Schülke, pp. 179–201. Routledge,
London & New York.
- 560 Fuglestvedt, Ingrid
- 561 2012 The Pioneer Condition on the Scandinavian Peninsula: the Last Frontier of a ‘Palaeolithic Way’ in
Europe. *Norwegian Archaeological Review* 45(1):1–29. DOI:10.1080/00293652.2012.669998.
- 562 Gjerpe, Lars Erik, and Grethe Bjørkan Bukkemoen
- 563 2008 Nordby 1 – Toskipede hus fra neolitikum-bronsealder og boplasspor fra jernalder. In *E18-prosjektet*
Vestfold. Bind 3. Hus, boplass- og dyrkningspor, edited by Lars Erik Gjerpe, pp. 7–38. University of
Oslo, Museum of Cultural History, Oslo.
- 564 Glørstad, Håkon
- 565 2010 *The Structure and History of the Late Mesolithic Societies in the Oslo Fjord Area 6300-3800 BC.*
Bricoleur Press, Lindome.
- 566 2012 Historical ideal types and the transition to the Late Neolithic in South Norway. In *Becoming European.*
The transformation of third millennium Northern and Western Europe, edited by Christopher Prescott
and Håkon Glørstad, pp. 82–99. Oxbow Books, Oxford & Oakville.
- 567 2016 Deglaciation, sea-level change and the Holocene colonization of Norway. *Geological Society, London,*
Special Publications 411:9–25. DOI:10.1144/SP411.7.
- 568 Glørstad, Håkon, Jostein Gundersen, Frode Kvalø, Pål Nymoen, David Simpson, and Birgitte Skar
- 569 2020 Submerged Stone Age from a Norwegian Perspective. In *The Archaeology of Europe’s Drowned*
Landscapes, edited by Geoff Bailey, Nena Galanidou, Hans Peeters, Hauke Jöns, and Moritz Mennenga,
pp. 125–140. Springer, Cham.
- 570 GRASS Development Team
- 571 2017 *Geographic Resources Analysis Support System (GRASS) Software, Version 7.2.* Open Source Geospatial
Foundation.
- 572 Gundersen, Jostein
- 573 2013 Verken fjord eller fjell – steinalderen i det kystnære innlandet. Gamle og nye funn fra Notodden i
Telemark. *Viking* 76:35–62.
- 574 Hafsten, Ulf
- 575 1957 De senkvartære strandlinje-forskyvningene i Oslotrakten belyst ved pollenanalytiske undersøkelser.
Norwegian Journal of Geography 16(1-8):74–99. DOI:10.1080/00291955708622137.
- 576 1983 Shore-level changes in South Norway during the last 13,000 years, traced by biostrati-
graphical methods and radiometric datings. *Norwegian Journal of Geography* 37(2):63–79.
DOI:10.1080/00291958308552089.
- 577 Hagen, Anders
- 578 1963 Problemkompleks Fosna. Opphav – kontakt med kontinentale grupper – forholdet til Komsa. In
Boplatsproblem vid Kattegat och Skagerack, pp. 53–59. Göteborg och Bohusläns forminnesförening &
Institutionen för nordisk fornkunskap, Gothenburg University, Gothenburg.
- 579 Helskog, Knut
- 580 1978 Late Holocene sea-level changes seen from prehistoric settlements. *Norwegian Journal of Geography*
32:111–119. DOI:10.1080/00291957808552032.
- 581 Herzog, Irmela
- 582 2013 The Potential and Limits of Optimal Path Analysis. In *Computational Approaches to Archaeological*
Spaces, edited by Andrew Bevan and Mark Lake, pp. 179–211. Left Coast Press, Walnut Creek.
- 583 Hinsch, Erik
- 584 1955 Traktbegerkultur – Megalitkultur. En studie av Øst-Norges eldste neolitiske gruppe. *Universitetets*
Oldsaksamling Årbok 1951–1953:10–177.

- 599
- 600 Hinz, Martin, Clemens Schmid, Daniel Knitter, and Carolin Tietze
601 2021 *oxcAAR: Interface to 'OxCal' radiocarbon calibration. R package version 1.1.0.*
- 602
- 603 Hollender, Artur
604 1901 Om sveriges nivåförändringar efter människans invandring. *Geologiska Föreningen i Stockholm Förhandlingar* 23(4):1118–1130. DOI:10.1080/00293652.1975.9965220.
- 605
- 606 Hughes, Anna L. C., Richard Gyllencreutz, Øystein S. Lohne, Jan Mangerud, and John Inge Svendsen
607 2016 The last Eurasian ice sheets – a chronological database and time-slice reconstruction, DATED-1.
608 *Boreas* 45(1):1–45. DOI:<https://doi.org/10.1111/bor.12142>.
- 609 Hyndman, Rob J
610 1996 Computing and Graphing Highest Density Regions. *The American Statistician* 50(2):120–126.
- 611
- 612 Ilves, Kristin, and Kim Darmark
613 2011 Some Critical and Methodological Aspects of Shoreline Determination: Examples from the Baltic Sea
614 Region. *Journal of Archaeological Method and Theory* 18:147–165. DOI:10.1007/s10816-010-9084-x.
- 615 Jaksland, Lasse (editor)
616 2012a *E18 Brunlanesprosjektet. Bind II. Undersøkte lokaliteter fra tidligmesolitikum.* University of Oslo,
617 Museum of Cultural History, Oslo.
- 618 (editor)
619 2012b *E18 Brunlanesprosjektet. Bind III. Undersøkte lokaliteter fra tidligmesolitikum og senere.* University
620 of Oslo, Museum of Cultural History, Oslo.
- 621 2014 Kulturhistorisk sammenstilling. In *E18 brunlanesprosjektet. Bind i. Forutsetninger og kulturhistorisk
622 sammenstilling*, edited by Lasse Jaksland and Per Persson, pp. 11–46. University of Oslo, Museum of
Cultural History, Oslo.
- 623 Jaksland, Lasse, and Per Persson (editors)
624 2014 *E18 Brunlanesprosjektet. Bind I. Forutsetninger og kulturhistorisk sammenstilling.* University of Oslo,
625 Museum of Cultural History, Oslo.
- 626 Johansen, Erling
627 1963 Kyst(fangst)boplassenes strandbundenhet og strandlinjekronologien. In *Boplatssproblem vid Kattegat
628 och Skagerack*, pp. 90–92. Göteborg och Bohusläns fornminnesförening & Institutionen för nordisk
fornkunskap, Gothenburg University, Gothenburg.
- 629 Jørgensen, Erlend Kirkeng, Petro Pesonen, and Miikka Tallavaara
630 2020 Climatic changes cause synchronous population dynamics and adaptive strategies among coastal
631 hunter-gatherers in Holocene northern Europe. *Quaternary Research*:1–16. DOI:10.1017/qua.2019.86.
- 632 Kjemperud, Alfred
633 1986 Late Weichselian and Holocene shoreline displacement in the Trondheimsfjord area, central Norway.
634 *Boreas* 15(1):61–82. DOI:10.1111/j.1502-3885.1986.tb00744.x.
- 635 Kleppe, Else Johansen
636 1985 *Archaeological Data on Shore Displacements in Norway.* Norges geografiske oppmåling, Hønefoss.
- 637
- 638 Kleppe, Jan Ingolf
639 2018 The Pioneer Colonization of Northern Norway. In *Early Economy and Settlement in Northern Europe.
Pioneering, Resource Use, Coping with Change*, edited by Hans Peter Blankholm, pp. 13–57. Equinox,
640 Sheffield.
- 641 Lakens, Daniël, Anne M. Scheel, and Peder M. Isager
642 2018 Equivalence Testing for Psychological Research: A Tutorial. *Advances in Methods and Practices in
643 Psychological Science* 1(2):259–269. DOI:10.1177/2515245918770963.
- 644 Lewis, Joseph

- 645 2021 Probabilistic Modelling for Incorporating Uncertainty in Least Cost Path Results: a Postdictive Roman
Road Case Study. *Journal of Archaeological Method and Theory* 28(3):911–924. DOI:10.1007/s10816-
021-09522-w.
- 646 Marwick, Ben, Carl Boettiger, and Lincoln Mullen
- 647 2018 Packaging Data Analytical Work Reproducibly Using R (and Friends). *The American Statistician*
72(1):80–88. DOI:10.1080/00031305.2017.1375986.
- 648 Melvold, Stine, and Per Persson (editors)
- 649 2014 *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og*
Porsgrunn. Bind 1. Tidlig- Og mellommesolittiske lokaliteter i Vestfold og Telemark. Portal forlag,
Kristiansand.
- 650 Mikkelsen, Egil
- 651 1975 Mesolithic in South-Eastern Norway. *Norwegian Archaeological Review* 8(1):1118–1130.
DOI:10.1080/11035890109445866.
- 652 Milne, Glenn A
- 653 2015 Glacial isostatic adjustment. In *Handbook of sea-level research*, edited by Ian Shennan, Antony J Long,
and Benjamin P Horton, pp. 421–437. Wiley, Chichester.
- 654 Milne, Glenn A., W. Roland Gehrels, Chris W. Hughes, and Mark E. Tamisiea
- 655 2009 Identifying the causes of sea-level change. *Nature Geoscience* 2(7):471–478. DOI:10.1038/ngeo544.
- 656 Mjærum, Axel
- 657 2018 Hinterland discoveries: Middle Mesolithic woodland utilization and the case of the site Eidsberg,
eastern Norway. *Current Swedish Archaeology* 26(1):159–188. DOI:10.37718/CSA.2018.11.
- 658 2022 A Matter of Scale: Responses to Landscape Changes in the Oslo Fjord, Norway, in the Mesolithic.
Open Archaeology 8(1):62–84. DOI:10.1515/opar-2022-0225.
- 659 Møller, Jakob J
- 660 1987 Shoreline relation and prehistoric settlement in northern norway. *Norwegian Journal of Geography*
41:45–60. DOI:<http://dx.doi.org/10.1080/00291958708552171>.
- 661 Mörner, Nils-Axel
- 662 1976 Eustasy and Geoid Changes. *The Journal of Geology* 84(2):123–151. DOI:10.1086/628184.
- 663 1979 The Fennoscandian Uplift and Late Cenozoic Geodynamics: Geological Evidence. *GeoJournal* 3(3):287–
318. DOI:10.1007/BF00177634.
- 664 Nielsen, Svein Vatsvåg
- 665 2021 Early farming in Southeastern Norway: New evidence and interpretations. *Journal of Neolithic*
Archaeology 23:83–113. DOI:10.12766/jna.2021.4.
- 666 Nielsen, Svein Vatsvåg, Per Persson, and Steinar Solheim
- 667 2019 De-Neolithisation in southern Norway inferred from statistical modelling of radiocarbon dates. *Journal*
of Anthropological Archaeology 53:82–91. DOI:10.1016/j.jaa.2018.11.004.
- 668 Nordqvist, Bengt
- 669 1995 The Mesolithic settlement of the west coast of Sweden - with special emphasis on chronology and
topography of coastal settlements. In *Man and Sea in the Mesolithic. Coastal settlement above and*
below present sea level, edited by Anders Fischer, pp. 185–196. Oxbow Books, Oxford.
- 670 1999 The Chronology of the Western Swedish Mesolithic and Late Paleolithic: Old Answers in Spite of New
Methods. In *The Mesolithic of Central Scandinavia*, edited by Joel Boaz, pp. 235–253. University of
Oslo, Oslo.
- 671 Norwegian Mapping Authority
- 672 2018 *Produktspesifikasjon. Nasjonal modell for høydedata fra laserskanning (FKB-laser).* FKB-laser_v30.
- 673 Nummedal, Anders
- 674 1923 Om flintpladsene. *Norwegian Journal of Geography* 7(2):89–141.

- 692 Østmo, Einar
693 1988 *Etableringen av jordbrukskultur i Østfold i steinalderen*. The University Collection of National
694 Antiquities, University of Oslo, Oslo.
695 2008 *Auve. En fangstboplass fra yngre steinalder på Vesterøya i Sandefjord. I. Den arkeologiske del*. Museum
696 of Cultural History, University of Oslo, Oslo.
697 Parnell, A. C., J. Haslett, J. R. M. Allen, C. E. Buck, and B. Huntley
698 2008 A flexible approach to assessing synchronicity of past events using Bayesian reconstructions of sedimenta-
699 tion history. *Quaternary Science Reviews* 27(19-20):1872–1885. DOI:10.1016/j.quascirev.2008.07.009.
700 Persson, Per
701 2008 Nauen 5.2 – Stenåldersboplatter och fossil åkermark. In *E18-prosjektet Vestfold. Bind 2. Steinalderbo-
702 plasser, boplasspor, graver og dyrkningspor*, edited by Lars Erik Gjerpe, pp. 163–198. University of
Oslo, Museum of Cultural History, Oslo.
703 Prescott, Christopher
704 2020 Interpreting Complex Diachronic "Neolithic"-Period Data in Norway. In *Farmers at the Frontier –
705 A Pan European Perspective on Neolithisation*, edited by Kurt J. Gron, Lasse Sørensen, and Peter
Rowley-Conwy, pp. 381–400. Oxbow Books, Oxford.
706 R Core Team
707 2021 *R: A language and environment for statistical computing*. R Foundation for Statistical Computing,
708 Vienna, Austria.
709 Ramstad, Morten
710 2009 Eldre steinalder på Melkøya, representativitet, strandlinjer og transgresjon. In *Undersøkelsene på
711 Melkøya. Melkøyaprosjektet – Kulturhistoriske registreringer og utgravninger 2001 og 2002*, edited by
Anders Hesjedal, Morten Ramstad, and Anja R. Niemi, pp. 491–495. Tromsø museum, Universitetes-
museet, Tromsø.
712 Reimer, Paula J., William E. N. Austin, Edouard Bard, Alex Bayliss, Paul G. Blackwell, Christopher Bronk
713 Ramsey, Martin Butzin, Hai Cheng, R. Lawrence Edwards, Michael Friedrich, Pieter M. Grootes, Thomas
714 P. Guilderson, Irka Hajdas, Timothy J. Heaton, Alan G. Hogg, Konrad A. Hughen, Bernd Kromer, Sturt
715 W. Manning, Raimund Muscheler, Jonathan G. Palmer, Charlotte Pearson, Johannes van der Plicht, Ron
716 W. Reimer, David A. Richards, E. Marian Scott, John R. Southon, Christian S. M. Turney, Lukas Wacker,
717 Florian Adolphi, Ulf Büntgen, Manuela Capoano, Simon M. Fahrni, Alexandra Fogtmann-Schulz, Ronny
718 Friedrich, Peter Köhler, Sabrina Kudsk, Fusa Miyake, Jesper Olsen, Frederick Reinig, Minoru Sakamoto,
719 Adam Sookdeo, and Sahra Talamo
720 2020 The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon*
721 62(4):725–757. DOI:10.1017/RDC.2020.41.
722 Reitan, Gaute, and Inger Marie Berg-Hansen
723 2009 *Lundevågenprosjektet, delrapport 1. Sammenfattende rapport. Lunde, 6/1, 6/35 og Skjolnes 7/23,
724 7/27, Farsund kommune, Vest-Agder*. Oslo.
725 Reitan, Gaute, and Per Persson (editors)
726 2014 *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og
727 Porsgrunn. Bind 2. Seinmesolittiske, neolittiske og yngre lokaliteter i Vestfold og Telemark*. Portal
forlag, Kristiansand.
728 Reitan, Gaute, and Lars Sundström (editors)
729 2018 *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*. Cappelen Damm Akademisk,
730 Oslo.
731 Roalkvam, Isak
732 2020 Algorithmic classification and statistical modelling of coastal settlement patterns in Mesolithic south-
eastern Norway. *Journal of Computer Applications in Archaeology* 3(1):288–307. DOI:<https://doi.org/10.5334/jcaa.60>.
733 2022 Exploring the composition of lithic assemblages in Mesolithic south-eastern Norway. *Journal of
734 Archaeological Science: Reports* 42:103371. DOI:10.1016/j.jasrep.2022.103371.
735 Rønberg, Frank Halvar N.

- 737 2012 *Bosettings- og aktivitetsspor. Larønningen*, 221/2138. Skien, Telemark. University of Oslo, Museum
738 of Cultural History, Oslo.
- 739 Romundset, Anders
- 740 2018 Postglacial shoreline displacement in the Tvedstrand-Arendal area. In *The Stone Age Coastal
Settlement in Aust-Agder, Southeast Norway*, edited by Gaute Reitan and Lars Sundström, pp.
741 463–478. Cappelen Damm Akademisk, Oslo.
- 742 Romundset, Anders, Stein Bondevik, and Ole Bennike
- 743 2011 Postglacial uplift and relative sea level changes in Finnmark, northern Norway. *Quaternary Science
Reviews* 30(19-20):2398–2421. DOI:10.1016/j.quascirev.2011.06.007.
- 744 Romundset, Anders, Ola Fredin, and Fredrik Høgaas
- 745 2015 A Holocene sea-level curve and revised isobase map based on isolation basins from near the southern
746 tip of Norway. *Boreas* 44:383–400. DOI:10.1111/bor.12105.
- 747 Romundset, Anders, Thomas R. Lakeman, and Fredrik Høgaas
- 748 2018 Quantifying variable rates of postglacial relative sea level fall from a cluster of 24 isolation basins in
749 southern Norway. *Quaternary Science Reviews* 197:175–192. DOI:10.1016/j.quascirev.2018.07.041.
- 750 2019 Coastal lake records add constraints to the age and magnitude of the Younger Dryas ice-front oscillation
751 along the Skagerrak coastline in southern Norway. *Journal of Quaternary Science* 34(2):112–124.
752 DOI:<https://doi.org/10.1002/jqs.3085>.
- 753 Schmitt, Lou, Stephan Larsson, Jan Burdukiewicz, John Ziker, Krister Svedhage, Jeanette Zamon, and
754 Steffen Holger
- 755 2009 Chronological Insights, Cultural Change, and Resource Exploitation on the West Coast of Sweden
During the Late Palaeolithic/Early Mesolithic Transition. *Oxford Journal of Archaeology* 28:1–27.
756 DOI:10.1111/j.1468-0092.2008.00317.x.
- 757 Schülke, Almut
- 758 2020 First visit or revisit? Motivations of mobility and the use and reuse of sites in the changing coastal
areas of Mesolithic southeastern Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement
with the Coast from the Atlantic to the Baltic Sea*, edited by Almut Schülke, pp. 359–393. Routledge,
759 London & New York.
- 760 Shennan, Ian
- 761 2015 Handbook of sea-level research: Framing research questions. In *Handbook of Sea-Level Research*, edited
762 by Ian Shennan, Antony J Long, and Benjamin P Horton, pp. 3–25. Wiley, Chichester.
- 763 Shetelig, Haakon
- 764 1922 *Primitive Tider i Norge – En oversikt over stenalderen*. John Griegs Forlag, Bergen.
- 765
- 766 Sognnes, Kalle
- 767 2003 On shoreline dating of rock art. *Acta Archaeologica* 74:189–209.
- 768
- 769 Solheim, Steinar
- 770 2012 Lokal praksis og fremmed opphav. Arbeidsdeling, sosiale relasjoner og differensiering i østnorsk
771 tidligeolitikum. Unpublished PhD thesis, Oslo.
- 772 (editor)
- 773 2017 *E18 Rugtvedt-Dørdal. Arkeologiske undersøkelser av lokaliteter fra steinalder og jernalder i Bamble
kommune, Telemark fylke*. Portal forlag, Kristiansand.
- 774
- 775 2020 Mesolithic coastal landscapes. Demography, settlement patterns and subsistence economy in south-
eastern Norway. In *Coastal Landscapes of the Mesolithic: Human Engagement with the Coast from the
Atlantic to the Baltic Sea*, edited by Almut Schülke. Routledge, London & New York.
- 776
- 777 2021 Timing the Emergence and Development of Arable Farming in Southeastern Norway by Using Summed
Probability Distribution of Radiocarbon Dates and a Bayesian Age Model. *Radiocarbon* 63(5):1–22.
778 DOI:10.1017/RDC.2021.80.
- 779 Solheim, Steinar, and Per Persson

- 780 2018 Early and mid-Holocene coastal settlement and demography in southeastern Norway: Comparing distribution of radiocarbon dates and shoreline-dated sites, 8500–2000 cal. BCE. *Journal of Archaeological Science: Reports* 19:334–343. DOI:10.1016/j.jasrep.2018.03.007.
- 781
- 782 Sørensen, Rolf
- 783 1979 Late Weichselian deglaciation in the Oslofjord area, south Norway. *Boreas* 8(2):241–246. DOI:<https://doi.org/10.1111/j.1502-3885.1979.tb00806.x>.
- 784
- 785 1999 En ^{14}C datert og dendrokronologisk kalibrert strandforksyvningskurve for sørøst Østfold. Sørøst-Norge. In *Museumslandskap. Artikkelsamling til Kerstin Griffin på 60-årsdagen. Bind A*, edited by Lotte Selsing and Grete Lillehammer, pp. 227–242. AmS-rapport 12A. Museum of Archaeology, Stavanger.
- 786
- 787 Sørensen, Rolf, Kari E. Henningsmoen, Helge I. Høeg, and Veronika Gälman
- 788 2014 Holocene landhevningsstudier i sørøst Vestfold og sørøstre Telemark – Revidert kurve. In *Vestfoldbaneprosjektet. Arkeologiske undersøkelser i forbindelse med ny jernbane mellom Larvik og Porsgrunn. Bind 1*, edited by Stine Melvold and Per Persson, pp. 36–47. Portal, Kristiansand.
- 789
- 790 Sørensen, Rolf, Kari E Henningsmoen, Helge I Høeg, and Veronika Gälman
- 791 in prep Holocen vegetasjons historie og landhevning i sørøst Vestfold og sørøstre Telemark. In *The Stone Age in Telemark. Archaeological results and scientific analysis from Vestfoldbaneprosjektet and E18 Rugtvedt-Dørdal*, edited by Per Persson and Steinar Solheim.
- 792
- 793 Sørensen, Rolf, Helge I. Høeg, Kari E. Henningsmoen, Göran Skog, Solveig F. Labowsky, and Bjørg Stabell
- 794 2014 Utviklingen av det senglasiale og tidlig preboreale landskapet og vegetasjonen omkring steinalderboplassene ved Pauler. In *E18 Brunlanesprosjektet. Bind I. Forutsetninger og kulturhistorisk sammenstilling*, edited by Lasse Jakobsen and Per Persson, pp. 171–213. University of Oslo, Museum of Cultural History, Oslo.
- 795
- 796 Stokke, Jo-Simon Frøshaug, and Gaute Reitan
- 797 2018 Kvastad A2. Lokalitet med funn fra tidlig- og mellommesolitikum og dyrkningsspor fra mellom- og senneolitikum. In *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*, edited by Gaute Reitan and Lars Sundström, pp. 375–407. Cappelen Damm Akademisk, Oslo.
- 798
- 799 Stroebe, Arjen P., Clas Hättestrand, Johan Kleman, Jakob Heyman, Derek Fabel, Ola Fredin, Bradley W. Goodfellow, Jonathan M. Harbor, John D. Jansen, Lars Olsen, Marc W. Caffee, David Fink, Jan Lundqvist, Gunhild C. Rosqvist, Bo Strömberg, and Krister N. Jansson
- 800 2016 Deglaciation of Fennoscandia. *Quaternary Science Reviews* 147:91–121.
- 801 DOI:10.1016/j.quascirev.2015.09.016.
- 802
- 803 Svendsen, John Inge, and Jan Mangerud
- 804 1987 Late Weichselian and Holocene sea-level history for a cross-section of western Norway. *Journal of Quaternary Science* 2(2):113–132. DOI:10.1002/jqs.3390020205.
- 805
- 806 Tallavaara, Miikka, and Petro Pesonen
- 807 2020 Human ecodynamics in the north-west coast of Finland 10,000–2000 years ago. *Quaternary International* 549:26–35. DOI:10.1016/j.quaint.2018.06.032.
- 808
- 809 Wang, Ian
- 810 2019 *topoDistance: Calculating topographic paths and distances. R package version 1.0.1.* [Https://CRAN.R-project.org/package=topoDistance](https://CRAN.R-project.org/package=topoDistance).
- 811
- 812 Wenn, Camilla Cecilie
- 813 2012 *Bosettingsspor, produksjonsområde og dyrkningsspor fra Neolitikum til Folkevandringstid. Bratsberg, 63/69, 244. Skien kommune, Telemark.* University of Oslo, Museum of Cultural History, Oslo.
- 814
- 815 Wikell, Roger, Fredrik Molin, and Mattias Pettersson
- 816 2009 The Archipelago of Eastern Middle Sweden - Mesolithic Settlement in Comparison with ^{14}C and Shoreline Dating. In *Chronology and Evolution within the Mesolithic of North-West Europe*, edited by Philippe Crombé, Mark van Strydonck, Joris Sergant, Mathieu Boudin, and Machteld Bats, pp. 417–434. Cambridge Scholar Publishing, Brussels.
- 817
- 818 Yubero-Gómez, María, Xavier Rubio-Campillo, and Javier López-Cachero
- 819

-
- 820 2016 The study of spatiotemporal patterns integrating temporal uncertainty in late prehistoric settlements
in northeastern Spain. *Archaeological and Anthropological Sciences* 8(3):477–490. DOI:10.1007/s12520-
821 015-0231-x.