
1 A simulation-based assessment of the relation between Stone Age
2 sites and relative sea-level change along the Norwegian Skagerrak
3 coast

4 Isak Roalkvam^{1,*}

5 10 March, 2022

6 ¹ University of Oslo, Institute of Archaeology, Conservation and History

7 * Correspondence: Isak Roalkvam <isak.roalkvam@iakh.uio.no>

8 **1 Introduction**

9 The post-glacial relative sea-level fall that characterises large areas of Fennoscandia is fundamental to
10 its archaeology. This follows not only from the dramatic changes to the landscape that this process has
11 represented throughout prehistory, but also from the fact that if archaeological phenomena were situated
12 close to the contemporary shoreline when they were in use, a reconstruction of the trajectory of shoreline
13 displacement can be used to date these phenomena based on their altitude relative to the present day sea-level.
14 This method, also called shoreline dating, has long history of use in the region and is frequently applied
15 to assign an approximate date to diverse archaeological phenomena such as rock art, grave cairns, various
16 harbour and sea-side constructions and, as is the focus of this study, Stone Age sites (e.g. Åkerlund 1996;
17 Bjerck 2005; Nordqvist 1995; Schmitt et al. 2009; Sognnes 2003; Tallavaara and Pesonen 2020; Wikell et al.
18 2009).

19 The close association between Stone Age settlements in the northern parts of Scandinavia and shifting
20 prehistoric shorelines was becoming apparent by the end of the 19th century (**degeer1896?**), and was firmly
21 established and first applied as a dating method in Sweden (Hollender 1901) and Norway (Brøgger 1905) at
22 the turn of the century. Shoreline dating has been fundamental to Stone Age archaeology in Norway ever
23 since (e.g. Berg-Hansen 2009; Bjerck 1990, 2008a; Breivik 2014; Johansen 1963; Mikkelsen 1975; Nummedal
24 1923; Solheim and Persson 2018). The method is used both independently, and to compliment other sources
25 of temporal data such as typological indicators or radiometric dates. However, given the coarse and fuzzy
26 resolution of established typological frameworks, the vast amount of surveyed sites that only contain generic
27 lithicdebitage that could hail from any part of the period, and as the conditions for the preservation of organic
28 material is typically poor in Norway, dating with reference to shoreline displacement is often the only and most
29 accurate method by which one can hope to date the sites. Shoreline dating is consequently fundamental to
30 our understanding of the Norwegian Stone Age. This is both because it is central to the temporal framework
31 on which our understanding of the period is based, but also because the method is only applicable so long as
32 the societies in question have continuously settled on or close to the contemporary shoreline. Consequently,
33 adherence or deviation from this pattern also has major implications for the socio-economic foundations of
34 the societies in question.

35 Despite its important role for Norwegian Stone Age archaeology, the applicability of dating by reference to
36 shoreline displacement has only been evaluated using relatively coarse methods. The aim of this paper is to
37 provide a systematic and comprehensive review of the degree to which radiocarbon dates correspond with the
38 dates informed by our current knowledge of shoreline displacement in a larger area of south-eastern Norway,
39 using a more refined methodological approach. The goal is to quantify the degree to which the assumption

40 of shore-bound settlement holds through the Stone Age, and in turn have this inform a refined method for
41 shoreline dating. As presented in more detail below, this problem involves the combined evaluation of three
42 major analytical dimensions. One is the questions of when the sites were in use, the second pertains to the
43 reconstruction of the contemporaneous sea-level, and the third follows from the fact that the relation between
44 site and shoreline is inherently spatial. Taking inspiration from studies that have integrated various sources
45 of spatio-temporal uncertainty through Monte Carlo simulation (e.g. Bevan et al. 2013; Crema et al. 2010;
46 Crema 2012, 2015; Yubero-Gómez et al. 2016), a similar approach is adopted here.

47 2 Background

48 Relative sea-level (RSL) can be defined as the mean elevation of the surface of the sea relative to land, or,
49 more formally, the difference in elevation between the geoid and the surface of the Earth as measured from
50 the Earth's centre (Shennan 2015). Variation in this relative distance follow from a range of effects (e.g.
51 Milne et al. 2009). Of central importance here is eustasy and istostasy. The eustatic sea-level is understood
52 as the sea-level if the water has been evenly distributed across the Earth's surface without adjusting for
53 variation in the rigidity of the Earth, its rotation, or the self-gravitation inherent to the water body itself
54 (Shennan 2015). The eustatic sea-level is mainly impacted by glaciation and de-glaciation, which can bind
55 or release large amounts of water into the oceans (Mörner 1976). Istostasy, on the other hand, pertains to
56 adjustments in the crust to regain gravitational equilibrium relative to the underlying viscous mantle. This is
57 often the result of glacial istostasy, which follows from glaciation and de-glaciation and corresponding loading
58 and unloading of weight, as well as from erosion of the crust, which causes its weight to be redistributed.
59 These effects thus causes the lithosphere to either subside due to increased weight, or to rebound and lift
60 upwards due to lower weight (Milne 2015).

61 Following the end of the Weichselian and the final retreat of the Fennoscandian Ice Sheet (e.g. Hughes et al.
62 2016; Stroeven et al. 2016, see Figure 1), the isostatic rebound has been so severe that most areas of Norway
63 have been subject to a continuous relative sea-level regression, despite corresponding eustatic sea-level rise
64 (e.g. Mörner 1979; Svendsen and Mangerud 1987). In other words, the RSL has been dropping throughout
65 prehistory. As this process is the result of glacial loading, the rate of uplift is more severe towards the centre
66 of the ice sheet. Thus, some areas on the outer coast have had a more stable RSL or been subject to marine
67 transgression (e.g. Romundset et al. 2015; Svendsen and Mangerud 1987). These conditions are directly
68 reflected in the archaeological record. In areas where the sea-level has been stable over longer periods of
69 time, people have often reused coastal site locations multiple times and over long time-spans, creating a
70 mix of settlement events that are difficult to disentangle (e.g. Hagen 1963; Reitan and Berg-Hansen 2009).
71 Transgression phases, on the other hand, can lead to complete destruction of the sites, or bury them in marine
72 sediments, leading to a hiatus in the archaeological record for certain sub-phases in the impacted areas (Bjerck
73 2008a; Glørstad et al. 2020). Comparatively, given a continuous and still ongoing shoreline regression from as
74 high as c. 220 m above present sea-level in the inner Oslo fjord, any one location in south-eastern Norway has
75 only been shore-bound within a relatively limited time-span, and the sites have not been impacted by any
76 transgressions (Hafsten 1957, 1983; Romundset et al. 2018; Sørensen 1979). This makes the region especially
77 useful for evaluating the assumption of a shore-bound settlement over a long and continuous time-span.

78 The method of shoreline dating has been met with scepticism as related to the degree sites would have
79 been consistently shore-bound, been characterised as a relative dating method for sites located at different
80 elevations within a constrained geographical area, or been argued to offer no more than a earliest possible
81 date for when a site could have been in use (Breivik and Bjerck 2018:174). The most common application in
82 Norway has arguably been to use the method to provide an approximate date for the occupation of the sites,
83 typically in combination with other dating methods [see for example chapters in @; ; and below]. Recently
84 the method has also been used independently to date a larger number sites to get a general impression of site
85 frequency over time, typically by aggregating point estimates of shoreline dates in 100, 200 or 500 year bins
86 (e.g. Breivik 2014; Breivik and Bjerck 2018; Fossum 2020; Nielsen 2021; Solheim and Persson 2018). In his
87 discussion of the method, Nordqvist (1999), in part drawing on Johansen (1963), argues that there can be
88 little doubt concerning its general applicability – what is less clear is the level of reliability and chronological




Figure 1: Deglaciation at 1000 year intervals from c. 17–8 kyr BCE. The study area defined later in the text is marked with a red outline (deglaciation data from Hughes et al. 2016, although see Romundset et al. 2019 for the study area).

89 resolution that the method can offer.

90 The shore-bound settlement location of prehistoric hunter-fisher-gatherers in Norway is generally believed
91 to follow both from the exploitation of aquatic resources and from movement and communication, which
92 would have been efficient on waterways (discussed in Berg-Hansen 2009; Bergsvik 2009). The same logic
93 has also been extended to the hinter- and inland regions, where sites are to be located along rivers and
94 lakes (Brøgger 1905; Glørstad 2010:57–87; but see Gundersen 2013; Mjærum 2018; Schülke 2020). This is to
95 take a dramatic turn at the transition to the Late Neolithic, around 2400 BCE, with the introduction of
96 the Neolithic proper (Prescott 2020; Solheim 2021). The introduction of a comprehensive Neolithic cultural
97 package, including a shift to agro-pastoralism and the introduction of the farm is to have led site locations
98 to be more withdrawn from the shoreline (e.g. Bakka and Kaland 1971; Østmo 2008:223; **prescott2012?**).
99 That is not to say that waterways and aquatic resources were no longer exploited, but rather that these
100 activities would not have been as tightly integrated with settlement and tool-production areas as in preceding
101 periods (Glørstad 2011). At an earlier stage, at the transition to the Early Neolithic (c. 3900 BCE), pottery
102 is introduced to the sites, and there are some indications of an initial uptake of agriculture at some sites
103 in the Oslo fjord region. However, this appears to be small in scale and is believed to be combined with a
104 continued and predominantly hunter-gatherer life-way, possibly followed by a complete de-Neolithisation in
105 the Middle Neolithic (Nielsen et al. 2019; Østmo 1988:225–227; **hinsch1955?**). Nielsen (2021) has recently
106 argued that the initial uptake of agriculture in Early Neolithic south-eastern Norway is combined with a more
107 complex settlement pattern, and that a simple forager/farming dichotomy would underplay the variation
108 present in the Early and Middle Neolithic settlement data (see also e.g. Amundsen et al. 2006; Østmo 1988;
109 Solheim 2012:74). Seen in relation to the question of interest here, the empirical expectation for the above
110 outlined development would thus be a predominantly shore-bound settlement in the Mesolithic, possibly
111 followed by a more varied association between sites and the shore-line from around 3900 BCE, and a decisive
112 shift around 2400 BCE.

113 Based on the generally accepted premise that most pre-Late Neolithic sites in south-eastern Norway located
114 lower than the marine limit were situated on or close to the contemporaneous shoreline, it is common to err
115 on the side of a shore-bound site location unless there is strong evidence to suggest otherwise. This is for
116 example common in survey projects, which are often guided by both a digital and mental reconstruction
117 of past sea-levels (e.g. Berg-Hansen 2009; Eskeland 2017). Similarly, following an excavation, if typological
118 indicators in the assemblages correspond with available shoreline displacement curves it is common to assume
119 a shore-bound site location, even if the typologically informed date-span is too wide to decisively verify this.
120 It is also common to combine this with a qualitative consideration of the landscape surrounding the sites,
121 and an evaluation of the degree to which the site location would have been sensible if the site was not shore
122 bound (e.g. Jakslund 2014; Johansen 1963; Nummedal 1923). This can for example pertain to accessibility. If
123 the site is situated on a ledge in a steep and jagged area of the present day landscape it would make intuitive
124 sense that the site was in use when the ocean reached closer to its elevation, as the site would have been
125 accessible by means of watercraft. Although it appears that the arguments for such site locations are sensible
126 and can for the most part be assumed to hold, comprehensive evaluations and attempts at quantification of
127 this tendency are relatively few (see also Ilves and Darmark 2011).

128 One of the more comprehensive evaluations of the relationship between radiocarbon dates and RSL-change
129 was done by Solheim and colleagues (Breivik et al. 2018; Solheim 2020), who compared 102 radiocarbon
130 dates from 33 Mesolithic sites on the western side of the Oslo fjord to the displacement curve for the Larvik
131 area. They found an overlap between the probability density of the radiocarbon dates with the shoreline
132 displacement curve for 86.5% of the sites. However, where there was a discrepancy, the main occupation of
133 the sites are still believed to have been shore-bound rather than associated with the deviating ^{14}C -dates.
134 This is based on typological and technological characteristics of the assemblages. Whether these mismatches
135 represent later shorter visits that are responsible for the younger radiocarbon dates, or whether these are
136 entirely erroneous results can be difficult to evaluate (e.g. Persson 2008; Schülke 2020). However, this
137 distinction is not deemed critical here, as what is of interest is the relationship between the prehistoric
138 shoreline and settlements and tool-production areas evidenced by artefact inventories or multiple site features.
139 Not remnants of stays as ephemeral to only be discernible by individual features or dubious ^{14}C -dates. The
140 evaluation of the relevance of radiocarbon dates to settlement activity will here therefore be entirely dependent

on, and follow the discretion of the original excavation reports.

Other previous evaluations of the correspondence between radiocarbon- and RSL-informed dates have typically followed the same structure as that of Breivik et al. (2018), involving a visual inspection of radiocarbon probability density functions plotted against local shoreline displacement curves based on the elevation of the site (e.g. Åkerlund et al. 1995; Åstveit 2018; Solheim 2020; see also Bjerck 2008b; Kleppe 1985). This approach has a couple of limitations. First of all, the displacement curves are sometimes applied directly to larger study areas, with only some studies having taken the variable uplift-rates into account when performing this comparison (e.g. Åstveit 2018; Fossum 2020; Møller 1987; Persson 2008). Secondly, with this method, the wider the uncertainty range associated with either radiocarbon date or displacement curve, the higher the probability that the confidence intervals overlap, and the higher the probability that we conclude in favour of our hypothesis. This thus leads to an inferential framework that favours uncertainty, which is hardly desirable. In statistical terms this follows from the fact that while one cannot conclude that two dates are different if their confidence intervals overlap, this does not necessarily mean that they are the same. The question thus necessitates a flip from a null-hypothesis of no significant difference, to one of equivalence (e.g. Lakens et al. 2018), as the question of interest is effectively one of synchronicity between events (cf. Parnell et al. 2008). Another limitation of this often-employed method is that it only takes into account the vertical distance between the sites and the sea-level. While this is the main parameter of interest for shoreline dating, the practical implications of a vertical difference in RSL will be highly dependent on local topography and bathymetry. RSL-change can have more dramatic consequences in a landscape characterised by a low relief, as the horizontal displacement of the shoreline will be greater. Taking the spatial nature of the relationship between site and shoreline into account will consequently help get more directly at the behavioural dimension of this relation, and move the analysis beyond a purely instrumental consideration of the applicability of shoreline dating. Suggested ways to help mitigate and integrate the issues presented above into to the analysis are presented in the methods section.

3 Data

To get at the relationship between sites and the contemporaneous shoreline, this analysis was dependent on a study area with good control of the trajectory of prehistoric shoreline displacement. While there is displacement data available for other areas of south-eastern Norway (e.g. Hafsten 1957; Sørensen 1979, 1999), considerable methodological developments in recent years means that the most well-established displacement curves are from the region stretching from Horten county in the north-east, to Arendal in the south-west. This area has newly compiled displacement curves for Horten (**romundset2021?**), Larvik (Sørensen et al. in press, 2014; Sørensen, Høeg, et al. 2014), Tvedstrand (Romundset 2018; Romundset et al. 2018), and Arendal (Romundset 2018).

The employed shoreline displacement curves are all based on the so-called isolation basin method (see e.g. Kjemperud 1986; Romundset et al. 2011). This involves extracting cores from a series of basins situated on bedrock at different elevations beneath the marine limit, and dating the transition from marine to lacustrine sediments. Each curve is thus construed from a series of cored basins located at different elevations, each representing a high precision sea-level index point (SLIP). Furthermore, to minimise the impact of variable uplift rates, the basins are located in a as constrained area of the landscape as possible. Following from the morphology of the retreating ice sheet, the uplift is more severe towards the north-east, meaning that this needs to be adjusted for in the case that any basins are located any significant distance from a common isobase perpendicular to this gradient (Figure 2). The resulting SLIPs indicate the isolation of the basin from the highest astronomical tide. This is adjusted to mean sea-level for the compilation of the displacement curve based on the present day tidal range, which is assumed to have been the same throughout the Holocene (Sørensen, Henningsmoen, et al. 2014:44). Furthermore, the confidence bands of the displacement curves and their trajectory are quite complex constructs, and are the integrated result of both expert knowledge and more objectively quantifiable parameters. The reason for this is in part that the curves do not only contain uncertainty as related to radiometric dates, which are well-defined, but also hold potential error as related to the interpretation and analysis of sediment cores, the nature and condition of the basin outlets and the

190 adjustment to a common isobase, to name but a few (Romundset et al. 2011, e.g. 2019). For more details
191 and evaluations done for the compilation of each curve, the reader is therefore referred to the individual
192 publications.

193 The archaeological data compiled for the analysis consists of excavated Stone Age sites with available spatial
194 data from the coastal region between Horten county in the north-east, to Arendal in the south-west (Figure
195 2). These number 155 sites. Of these, 91 sites are associated with a total of 547 radiocarbon dates. Of these,
196 in turn, 69 sites are related to the 266 radiocarbon dates that fall within the Stone Age (9500–1700 BCE),
197 with 95% probability. These sites and ^{14}C -dates form the basis for the analysis. Spatial data in the form of
198 site limits and features, as defined by the excavating archaeologists, were retrieved from local databases at the
199 Museum of Cultural History—the institution responsible for archaeological excavations in the region. In the
200 compiled dataset, each radiocarbon date has been associated with the site features or excavation unit from
201 where they originate, or, where these weren't available, the spatial limit of the entire site. Due to somewhat
202 variable practices between excavations, what available spatial geometry best represents the site limit was
203 decided based on an evaluation of the excavation reports. This means that the limits are variably given as
204 that defined during initial survey, area de-turfed before excavation, area stripped with excavator following
205 the excavation, manually excavated area, or convex hull polygons generated around the site features.

206 Three of the sites have been associated with farming activities, either directly or in the form building
207 structures. The first is Nordby 1, at which the ^{14}C -dates are associated with a Late Neolithic long-house
208 (Gjerpe and Bukkemoen 2008). The Middle Neolithic phase at Kvastad A2 (Stokke and Reitan 2018) and
209 Late Neolithic phase at Nauen A (Persson 2008) are both directly related to farming. Both of these sites also
210 have radiocarbon dates and lithic inventory associated with Mesolithic forager activities. Following from the
211 expected deviance from the settlement patterns that are to characterise forager sites, these agricultural phases
212 are thus treated separately below. Finally, Nielsen (2021) has recently suggested that Early and Middle
213 Neolithic features from the otherwise younger sites Bratsberg (Wenn 2012) and Larønningen (Røberg 2012)
214 could be related to early agricultural activity in the Oslo fjord region. Due to the uncertain and somewhat
215 speculative nature of this suggestion, these are omitted here.

216 [Table with sites]

217 The elevation data used for the analysis is a digital terrain model (DTM) freely available from the Norwegian
218 Mapping Authority (**mapping2019?**). It was here opted for the 10m resolution DTM rather than the higher-
219 resolution 1m version. In addition to resulting in considerably less processing time, the higher resolution
220 elevation model is more vulnerable to smaller-scale modern disturbances that the 10m version is not impacted
221 by. The 10m DTM is a down-sampled version of the 1m resolution DTM, which is based on aerial laser
222 scanning using a minimum of two and up to five points per m² and has a vertical error of (**mapping2019?**).
223 All data and R programming code (R Core Team 2021) required to run the analyses, as well as the derived
224 data are freely available in an online repository at <https://osf.io/7f9su/>, organised as a digital research
225 compendium following Marwick et al. (2018).

226 4 Methods

227 The method of shoreline dating is based on the spatial relationship between two phenomena, occupation of
228 sites and shoreline displacement, each associated with their own range of temporal uncertainty. The first task
229 was therefore to ascribe likely date ranges and associated uncertainty to these dimensions. To take account of
230 the gradient in the isostatic rebound, the trajectory of shoreline displacement was first interpolated to each
231 site location from the displacement curves using inverse distance weighting based on the distance between the
232 site and the isobases of the displacement curves (e.g. Conolly 2020; Conolly and Lake 2006:94–97). This was
233 done for each year along the entirety of the curves, weighting the interpolation by the squared inverse of the
234 distances. The result of this process is shown for an example site in Figure 3. For the date ranges associated
235 with the sites, all radiocarbon dates were first individually calibrated using the IntCal20 calibration curve
236 (Reimer et al. 2020) using OxCal v4.4.4 (Bronk Ramsey 2009) through the oxcAAR package for R (Hinz et
237 al. 2021). Radiocarbon dates associated with each site were then grouped if they overlapped with 99.7%




Figure 2: A) Distribution of the 69 analysed sites relative to the isobases of the displacement curves. The isobases have a direction of 327° (Romundset et al. 2018), B) Displacement curves. Note the increasing⁷ steepness of the curves towards the north-east.

238 probability, meaning these were effectively taken to represent the same settlement phase. In the case where
239 there are multiple dates believed to belong to a single phase, these were subjected to Bayesian modelling
240 using the Boundary function in OxCal and then summed. Multiple phases at a single site were treated as
241 independent of each other.

242 As the excavation of archaeological sites typically follow from residential and commercial development, as well
243 as the expansion of infrastructure, the area immediately surrounding the sites has sometimes been severely
244 impacted by modern disturbances. In addition to employing 10m resolution DTM to alleviate some of these
245 issues, this also necessitated some additional editing of the elevation raster. This involved manually defining
246 the extent of problem areas such as railways, highways, quarries and the like. The DTM values on these
247 were then set to missing, and new elevation values were interpolated from the surrounding terrain. This was
248 done using regularised spline interpolation with tension (e.g. Conolly 2020), using the default settings of
249 r.fillnulls from GRASS GIS (GRASS Development Team 2017) in R through the package rgrass7 (Bivand
250 2021). In addition to code and original spatial data being available in the research compendium for this
251 paper, it has also been noted in the supplementary material when the area surrounding a site has been edited
252 in this manner.

253 Armed with a likely date range for the occupation(s) of each site, an estimated trajectory of relative sea-level
254 change at that location, and a DTM edited to remove substantial modern disturbances, the simulations were
255 performed. A single simulation run involved first drawing a single year from the posterior density estimate of
256 a given occupation phase of a site (Figure 4). This year then has a corresponding likely elevation range for
257 the contemporaneous shoreline from which an elevation value was drawn uniformly, using intervals of 5cm.
258 The sea-level was then raised to this elevation on the DTM by defining all elevation values at or below this
259 altitude as missing. Polygons were then created from the resulting areas with missing values. The horizontal
260 distance was then found by measuring the shortest distance between site and sea polygons, and the vertical
261 distance by subtracting the elevation of the sea-level from the lowest elevation of the site polygon. The
262 topographic distance between site and sea was also found by measuring the shortest distance while taking into
263 account the slope of the terrain on the DTM. This was done using the topoDistance package for R (Wang
264 2019). The topographic distance was measured between site polygon and the horizontally closest point on the
265 shoreline. This means that the distance is not necessarily measured as the closest topographic distance to the
266 shoreline, but rather as the shortest topographic path to the horizontally closest point on the shoreline. Not
267 finding the topographically closest point significantly reduced the computational cost of the analysis, and is
268 deemed unlikely to have a considerable impact on the results, given the distances considered. The shortest
269 topographic path was found using the Moore neighbourhood of eight cells (e.g. Conolly and Lake 2006:253;
270 Herzog 2013). In the case where the sea-polygons intersects the site polygon, all distance measures were set
271 to zero. In the case that the sea-polygon completely contain the site, the horizontal and topographic distance
272 measures were made negative, and the vertical distance was instead measured to the highest point on the site
273 polygon. While it is safe to assume that an archaeological site was not occupied when it was located beneath
274 sea-level, a negative result can reflect the inherent uncertainty in this procedure, and might also help identify
275 discrepancies in displacement data or radiocarbon dates. Negative values were therefore retained, with the
276 exception of for the sites Gunnarsrød 5 and Pjonkerød R1, where the negative values are believed to result
277 from modern disturbances in the DTM rather than the ^{14}C -dates or displacement curves (see supplementary
278 material for more details).

279 This process was repeated 1000 times for each phase for each site. The choice of 1000 simulation runs follows
280 from an evaluation of when the mean distances between site and shoreline converged when running 5000
281 iterations of the simulation on the site Hovland 5, available in the supplementary material (cf. Crema et al.
282 2010:1125). Hovland 5 was chosen for this evaluation as it has a fairly uncertain date, and is located in area
283 of quite complex topography.

284 5 Simulation results

285 Overall, as is indicated by the measures for central tendency and the almost solid line along the 0m mark
286 on the y-axes, the simulations show that the sites tend to have been situated close to the shoreline when




Figure 3: Example site Hegna vest 1 (Fossum 2017). A) Location of the site in the present day landscape. The red outline is the site limit. B) Radiocarbon dates associated with the site. Fill colour indicates what dates are assumed to belong to the same settlement phase. Multiple dates are modelled using the Boundary function in OxCal and then summed. The red outline indicates that the date does not match the typological indicators in the artefact assemblage of the site. C) The location of the site within the study area relative to isobases of the employed displacement curves. D) Displacement curve interpolated to the site location.




Figure 4: Example of a single simulation run on the site Hegna vest 1. A) The simulation starts by drawing a single year from the posterior density estimate. B) This then corresponds to an elevation range on the interpolated displacement curve. A single elevation is drawn uniformly from this range using 5cm intervals. C) The sea-level is then adjusted on the DTM to this elevation and the various distance measures are found. D) The numerical result of the simulation run.




Figure 5: The result of 1000 simulation runs for each of the two groups of dates on the site Hegna vest 1. The first column of plots shows the radiocarbon probability density function from where dates were drawn during simulation. The second column displays the result of simulating the raised sea-level 1000 times. The more opaque the colour, the more times the sea-level was simulated to that location. The third column shows violin plots of the different distance measures across all simulations.

they were in use (Figure 6). Some of the sites are situated considerable distances from the shoreline when the dates believed to be erroneous in the original reports are included, but if one accepts the interpretation that these do not date the main occupation of the sites, as is indicated by the artefact inventories, the second row of Figure 6 gives considerable support to the notion that the sites were in use when they were situated on or close to the contemporaneous shoreline. The distances for some of the earliest sites appears somewhat high, but this can likely be explained as the result of the steepness of the displacement curves for the earliest part of the Holocene (Figure 2B), which leads the uncertainty of the ^{14}C -dates to give a wider possible elevation range for the sea-level. Another immediately striking result is the apparent deviation from the shoreline towards the end of the Stone Age, corresponding with the literature. From around 2500 BCE several sites are situated a considerable distance from the shoreline, and while a couple remain horizontally and topographically close, most appear to be elevated a considerable distance from the sea-level, as indicated on the plot for vertical distance. There are also a couple of sites located some distance from the shoreline just after 4000 BCE. While the sample size is limited, this would thus be in line with a development that sees an increase in settlements located in the immediate inland around this time.

The negative values around 8000 BCE originate from the sites Løvås 1, 2 and 3. These are recently excavated, well-dated sites situated in a relatively undisturbed area of the landscape. While there would be a danger of circularity of having archaeological sites inform a reconstruction RSL-change, and in turn use these to evaluate the degree of shore-bound settlement, the sites do clearly represent a upper limit for the sea-level, as they would not have been in use when located under water. It could therefore seem that the Løvås sites represent a case where the archaeological material indicates a slight discrepancy in the geological reconstruction of shoreline displacement in the area.

Accepting that shoreline dating appears to loose utility around the transition to the Late Neolithic, as indicated by the clear deviation in site location from the shoreline after this, the results for the second row of Figure 6 is given again in Figure 7, excluding all simulation results younger than 2500 BCE. Furthermore, all negative values have here been set to zero, under the assumption that these result from uncertainty or errors in the data, and not actual site locations. The resulting best point estimate for the vertical distance between sites and shoreline for the pre-Late Neolithic is given by the median at 4m, while 95% of the values fall within the range 0–17m. That is, for 95% of the cases, the shoreline was simulated to be situated on or down to 17m below the site location. While these values remain the same, the mean and standard deviation is constrained slightly when only considering the Mesolithic dates in the secone row of Figure 7. Furthermore, while the median for horizontal and topographic distances is only 10m across all plots in Figure 7, it is also evident that their range is substantially wider than that for vertical distance, illustrating the point that minor variations in vertical distance can have substantial consequences for these distance measures, depending on the surrounding topography.

An exponential function has been fit to the distributions for vertical distance using maximum likelihood estimation (Figure 7). While it makes theoretical sense that a process of exponential decay explains this relationship, it is also clear that this does not perfectly match the data. However, this can at least in part be related to methodological factors where the accumulation of distance-values on the 0m mark likely follow from forcing negative values to zero, from the resolution of the spatial data, and from defining intersecting sea- and site polygon as having a distance of zero. If one accepts this, having derived an exponential decay function for describing the vertical distance between sites and shoreline can be combined with the displacement data to provide a method for shoreline dating that takes this distance into account:

Where x is [...] In Figure 8 this formula is used to shoreline date the same sites from where this relationship was derived. Following from having defined the distance between intersecting sea- and site polygons as zero during simulations, the sites were dated using the mean elevation of the site polygons to allow for some deviance in elevation over the site limits. The result of comparing the radiocarbon and shoreline dates using the method presented by Parnell et al. (2008) for assessing the synchronicity between dates is given in Figure 8B. Here, 100,000 age samples drawn from the probability density function of each shoreline date was subtracted from 100,000 age samples drawn from the corresponding ^{14}C -date. The resulting range of the 95% highest density region (HDR, Hyndman 1996) was then checked to see if it crosses zero, in which case the dates are considered to be in agreement. The result of this procedure indicates that the shoreline date correspond to the radiocarbon dates for 46 out of 49 sites (94%). The deviation from the shoreline

339 date at Gunnarsrød 5 is to be expected based on the simulation results (see above), but not at Langemyr
 340 and Kvastad A2. For these two sites the zero age difference can be seen to fall just outside the 95% interval
 341 (Figure 8B). If one accepts that sensitivity in the method leads to the failure to accurately date these two
 342 sites, and that the failed dating of Gunnarsrød 5 is instead due to extraneous factors related the DTM and
 343 not the method itself, one could alternatively see the success rate as defined by 44 out of 46 sites (96%)
 344 having been correctly dated. However, another possible implication of the failed dating of these two sites
 345 could be that a lower decay ratio than what is used for characterising the distance between site and shoreline
 346 for all sites in aggregate should be used for sites from the earliest part of the Mesolithic (cf. Figure 6).




Figure 6: The result of running the analysis across all sites. Each data point is plotted with some transparency, meaning that the more intense the colour, the more often those values occurred. Results associated with agricultural activities are plotted in grey. The first row shows the result of including all dates to the Stone Age, including those seen as otherwise unrelated to the main occupation of the sites. The second row shows the result of excluding these. The table under each plot lists some corresponding statistics for central tendency and dispersion.




Figure 7: Histograms showing the simulated distance from the shoreline using dates corresponding to the site inventory. Negative values have been set to zero. The first row only includes simulated results older than 2500 BCE and the second row only results older than 4000 BCE.




Figure 8: The result of backwards shoreline dating the sites with radiocarbon dates corresponding to the artefact inventory using the method proposed here. A) The shoreline dates are plotted in grey and underlined¹⁵ with the 95% HDR in black. These are plotted against the oldest radiocarbon-informed phase for each site in red. B) Using the method described by Parnell et al. (2008), the synchronicity of the dates is evaluated. When the range of the 95% HDR for age difference crosses zero the dates are considered synchronous.

347 6 Shoreline dating

348 To further explore the implementation for shoreline dating outlined above, excavated and shoreline-dated
349 Stone Age sites within the study area where ^{14}C -dates are not available or are not believed to date the main
350 occupation of the sites have been subjected to the method (Figure 9). The resulting dates are compared to
351 those originally proposed in the excavation reports for the sites (the numerical results are available in the
352 supplementary material). To avoid issues with recent disturbances on the DTM, the sites have been dated
353 based on the mean of the altitudes provided in the report for each site. The results highlight the spatial and
354 temporal contingency of the method, illustrated by the variation in the range of the 95% intervals for the
355 dates. In some cases the method provides a very precise date range and in others it offers little more than a
356 *terminus post quem*. This is dependent on the steepness of the displacement curves, leading to the general
357 pattern of older sites situated towards the north-east getting more precise dates (cf. Figure 2B).

358 The comparison with previously reported dates is an illustrative, but unfair exercise for a few reasons. First of
359 all the dates provided in the reports are typically stated to be a very rough estimate, and are sometimes given
360 as a point estimate with an implied or explicit, but undefined uncertainty range. Secondly, seeing as these
361 reports are from various dates in time, many are based on now outdated data on RSL-change. Finally, they
362 are sometimes only meant to indicate a lower bound for when the sites could have been in use. Nonetheless,
363 the results indicate that shoreline dating, in general, has been applied with a fairly reasonable degree of
364 success, seeing as these dates have typically been interpreted and informed research in an approximate
365 manner (although see e.g. **roalkvam2022?**). That being said, the results do also indicate that shoreline
366 dating has at times been applied with an exaggerated degree of precision, and while the implications of
367 a more stable RSL-change for the precision of shoreline dating is well-known, it appears to be somewhat
368 under-appreciated. Some of the date ranges resulting from the method outlined here clearly extend well
369 beyond major chronological divisions, even into the Iron Age, and could be severely and securely constrained
370 with only cursory reference to typology. However, while this is obvious in some cases, the nature and
371 uncertainty inherent to the method still means that this is arguably a required exercise that should be made
372 explicit. This final point also points to the possibility of drawing on other temporal data, for example within
373 a Bayesian framework, to further improve the precision of the dates that can be achieved with shoreline
374 dating.

375 Not least following from the fact that few Preboreal ^{14}C -dates associated with anthropogenic activity have
376 been achieved in Norway, the shoreline dating of the earliest sites is essential for understanding the pioneer
377 settlement and the initial colonisation of the Scandinavian peninsula (e.g. Bang-Andersen 2012; Breivik 2014;
378 Glørstad 2016). The shoreline-dated Preboreal sites from the Brunlanes-project are among the earliest known
379 sites in Norway (Jaksland 2012a, 2012b; Jaksland and Persson 2014). These have a distinct Early Mesolithic
380 artefact inventory and are situated in a steep area of the landscape where it would be difficult to envision use
381 of the sites after the sea retreated any significant distance from their location. In the original publication
382 of the sites, Jaksland (2014) provides a thorough discussion of shoreline dating in general, and as used for
383 the dating of the Brunlanes sites specifically. A comparison of his results and the ones achieved using the
384 above-outlined approach are given in Figure 10A. The sites have been dated using what Jaksland (2014) gives
385 as the lowest elevation of finds at each site, and by employing a exponential decay ratio of , to allow for more
386 deviance in the distance between site and shoreline. This corresponds to the decay ratio for sites older than
387 7000 BCE in Figure 7.

388 The small discrepancies between the achieved results mainly follow from the fact that a slightly updated
389 version of the local displacement curve is applied here (cf. Sørensen et al. in press). Jaksland's dates are
390 given a flat 200 and 50 year uncertainty range starting from what he gives as the earliest possible date. The
391 200 year uncertainty range is given if the sites were to be considered in isolation, while the argument for
392 the uncertainty range of only 50 years is based on the location of the sites relative to each other. Since they
393 are located in such a constrained and steep area of the landscape, the difference in elevation between the
394 sites is argued to establish their relative date and thus constrain the uncertainty ranges so that they don't
395 overlap. This information is not integrated in the approach outlined here, but could justify further reducing
396 the uncertainty ranges. Although their accuracy is of course ultimately dependent on the veracity of the
397 geological reconstruction, the high rate of RSL-change in this period does nonetheless result in very precise

$$\lambda = 0.177$$




Figure 9: Re-dating previously shoreline dated sites in the study area without radiocarbon dates or with radiocarbon dates that do not correspond to the artefact inventories. The range of the 95% HDR in grey are compared to the dates originally proposed by the excavation reports in red.¹⁷

398 dates. Above it was shown how additional temporal data could be combined with the method to improve its
 399 accuracy and precision. This example, on the other hand, highlights the fact that the spatial nature of the
 400 method means that a consideration of the surrounding terrain and other sites can also help in increasing the
 401 precision of the method if this can be used to exclude certain sea-levels as unlikely for when a site was in use.
 402 One approach could also be to assess the spatial implication of a proposed shoreline date by simulating the
 403 adjusted sea-levels, as is done for Paurer 1 in Figure 10B, followed for example by a visual evaluation of the
 404 topography or by evaluating the distance and steepness of the slope to the shoreline. Based on this, it could
 405 conceivably be possible to exclude certain elevations as unlikely for the position of the shoreline when the site
 406 was in use. Such approaches would make less of an impact in this setting, where the 95% HDR is already
 407 quite constrained, but could considerably improve the precision of the method in cases where RSL-change
 408 has been less severe (cf. Figure 9).




Figure 10: Shoreline dating of the Brunlanes sites using site altitudes provided by Jakobsson (2014:tab.4). A) The result of applying the approach to shoreline dating outlined above. Dates provided by Jakobsson (2014) are plotted in red. The box indicates a 50 year uncertainty range which in combination with the red line extends 200 years. Following Jakobsson's approach these start from the earliest possible date. B) Map showing the centroids of the Paurer sites and Sky 1. The sea-level has been simulated using the probability density associated with the shoreline date for Paurer 1 (see also map in Jakobsson 2014:fig.12a). Paurer 1 is the red point.

409 7 Concluding remarks

- 410 It is worth highlighting some sources of likely variation and uncertainty that have not been considered here.
411 First of all the DTM has only been corrected for major modern disturbances. This means that erosion,
412 although likely not that prevalent, has not been taken into account. Secondly, the DTM has a vertical
413 error (cf. Lewis 2021). Thirdly, the displacement curves were here interpolated to all site locations without
414 accounting for increased uncertainty as one moves further away from the isobases of the displacement curves.
415 Finally, the question of how site limits are defined nor the elevation range over which these extend was given
416 much consideration. While each of these factors will have variable but likely not too large of an impact on
417 the results, they clearly represent dimensions which would all benefit from further exploration.
- 418 The most immediate contribution of this paper is what must be considered a confirmation of previous research
419 into the relation between coastal Norwegian Stone Age sites and the prehistoric shoreline. This is indicated by
420 the close relationship between sites and the shoreline up until the transition to the Neolithic at c. 3900 BCE,
421 after which a couple of sites become situated some distance from the shoreline. This is followed by a more
422 decisive break at the transition to the Late Neolithic at c. 2400 BCE – in clear agreement with the literature.
423 Furthermore, based on the quantitative nature of these findings, a refined method for the shoreline dating
424 of pre-Late Neolithic Stone Age sites has been proposed. This involves both taking the distance between
425 sites and the isobases of the displacement curves into consideration when dating the sites, and implementing
426 formula X to account for the distance between the sites and the shoreline. When no other information is
427 available, it can at present be recommended to use the empirically derived exponential decay ratio identified in
428 Figure 9 to characterise this relationship. However, the accuracy of the method can be improved by including
429 more information, both with reference to the topographic location of the sites and other temporal data such
430 as radiocarbon dates and typological indicators in the artefact inventories. The precision of the method is, as
431 shown above, both geographically and temporally contingent due to the trajectory of RSL-change, where
432 older sites situated towards the north-east in the study area will get a more precise date than younger sites
433 located towards the south-west. The impact of such additional information will therefore also vary.
- 434 Future investigations and radiocarbon dates from Stone Age sites in the region can not only be used to further
435 evaluate and adjust the findings reported here, but a larger sample size could also lay the foundations for
436 refining the method by identifying subsets of sites for which the application of the method could be adjusted
437 and refined. Given its behavioural nature, it would for example seem likely that dimensions such as the
438 nature and purpose of visits to the sites will have implications for how close to the shoreline they were located.
439 Furthermore, other dimensions related to the topographic location of the sites could be similarly explored.
440 This for example pertains to the exposure of sites to wave action, which is likely to have been of concern
441 (e.g. Roalkvam 2020), and which presumably has implications for how close to the shoreline people settled
442 (Blankholm 2020; Helskog 1978). This is also related to the fact that while the mean sea-level is used for
443 dating the sites, a consideration of the tidal range could possibly also have implications for the site location
444 relative to the shoreline, depending on the topography (Helskog 1978). Dimensions such as these were given
445 a cursory treatment here, with the estimation of the horizontal and topographic distance to the shoreline,
446 but these dimensions could also be investigated further. If patterns related to such locational patterns can
447 be discerned, this will not least be useful for improving the shoreline dating of sites which have only been
448 surveyed and where little information on the site beyond its location is available.
- 449 Finally, this analysis was based on a simulation approach for integrating multiple sources of spatio-temporal
450 uncertainty. Here this was simply used to inform the question of the distance between sites and the shore-line.
451 However, this method and general framework can be extended to a wide range of use-cases where one needs
452 to visualise and quantitatively or qualitatively evaluate the relationship between archaeological phenomena,
453 the prehistoric shore-line, and the uncertainty inherent in this reconstruction.

454 8 References

- 455 Åkerlund, Agneta
456 1996 *Human responses to shore displacement: Living by the sea in eastern middle sweden during the stone*
457 *age.* Riksantikvarieämbetet.
- 458 Åkerlund, Agneta, Jan Risberg, Urve Miller, and Per Gustafsson
459 1995 On the applicability of the 14C method to interdisciplinary studies on shore displacement and settlement
460 location. *PACT* 49:53–84.
- 461 Amundsen, Øystein, Stig Knutsen, Axel Mjærum, and Gaute Reitan
462 2006 Nøkleby i ski – en tidligeolittisk jordbruksbopllass? *Primitive tider* 9:85–96.
- 463
- 464 Åstveit, Leif Inge
465 2018 The early mesolithic of western norway. In *Early economy and settlement in northern europe. Pioneering, resource use, coping with change*, edited by Hans Peter Blankholm, pp. 231–274. Equinox,
466 Sheffield.
- 467 Bakka, Egil, and Peter Emil Kaland
468 1971 Early farming in Hordaland, western Norway. Problems and approaches in archaeology and pollen
469 analysis. *Norwegian Archaeological Review* 4:1–17. DOI:10.1080/00293652.1971.9965136.
- 470 Bang-Andersen, Sveinung
471 2012 Colonizing Contrasting Landscapes. The Pioneer Coast Settlement and Inland Utilization in
472 Southern Norway 10,000–9500 Years Before Present. *Oxford Journal of Archaeology* 31:103–120.
DOI:10.1111/j.1468-0092.2012.00381.x.
- 473 Berg-Hansen, Inger Marie
474 2009 *Steinalderregistrering. Metodologi og forskningshistorie i norge 1900-2000 med en feltstudie fra lista i*
475 *vest-agder.* Museum of Cultural History, University of Oslo.
- 476 Bergsvik, Knut Andreas
477 2009 Caught in the middle: functional and ideological aspects of Mesolithic shores in Norway. In *Mesolithic*
478 *horizons: Papers presented at the seventh international conference on the mesolithic in europe, belfast*
2005, edited by Sinéad B. McCartan, Rick Schulting, Graeme Warren, and Peter Woodman, pp.
602–609. Oxbow Books, Oxford.
- 479 Bevan, Andrew, Enrico R. Crema, Xiuzhen Li, and Alessio Palmisano
480 2013 Intensities, interactions, and uncertainties: Some new approaches to archaeological distributions. In
481 *Computational approaches to archaeological spaces*, edited by Andrew Bevan and Mark Lake, pp. 27–52.
Left Coast Press, Walnut Creek.
- 482 Bivand, Roger
483 2021 *rgrass7: Interface between GRASS 7 geographical information system and r.* R package version 0.2-6.
- 484
- 485 Bjerck, Hein Bjartmann
486 1990 Mesolithic site types and settlement patterns at vega, northern norway. *Acta Archaeologica* 60:1–32.
- 487
- 488 2005 Strandlinjedatering. In *Norsk arkeologisk leksikon*, edited by Einar Østmo and Lotte Hedeager, pp.
489 363–364. Pax, Oslo.
- 490 2008a Norwegian mesolithic trends: A review. In *Mesolithic europe*, edited by Geoff Bailey and Penny
491 Spikins, pp. 60–106. Cambridge University Press, Cambridge.
- 492 2008b Innledende betraktninger. In *NTNU vitenskapsmuseets arkeologiske undersøkelser ormen lange ny-*
493 *hamna*, edited by Hein Bjartmann Bjerck, Leif Inge Åstveit, Trond Meling, Jostein Gundersen, Guro
Jørgensen, and Staale Normann, pp. 548–551. Tapir Akademisk Forlag, Trondheim.
- 494 Blankholm, Hans Peter
495 2020 In the wake of the wake. An investigation of the impact of the storegga tsunami on the human
settlement of inner varangerfjord, northern norway. *Quaternary International* 549:65–73. DOI:<https://doi.org/10.1016/j.quaint.2018.05.050>.
- 496

- 497 Breivik, Heidi Mjelva
498 2014 Palaeo-oceanographic development and human adaptive strategies in the Pleistocene–Holocene transition: A study from the Norwegian coast. *The Holocene* 24:1478–1490. DOI:10.1177/0959683614544061.
499
- 500 Breivik, Heidi Mjelva, Guro Fossum, and Steinar Solheim
501 2018 Exploring human responses to climatic fluctuations and environmental diversity: Two stories from Mesolithic Norway. *Quaternary International* 465. Impacts of gradual and abrupt environmental changes on Late glacial to Middle Holocene cultural changes in Europe:258–275. DOI:10.1016/j.quaint.2016.12.019.
502
- 503 Breivik, Heidi, and Hein Bjartmann Bjerck
504 2018 Early mesolithic central norway: A review of research history, settlements, and tool tradition. In *Early economy and settlement in northern europe. Pioneering, resource use, coping with change*, edited by Hans Peter Blankholm, pp. 169–206. Equinox, Sheffield.
505
- 506 Brøgger, Waldemar Christofer
507 1905 *Strandliniens Beliggenhed under Stenalderen i Det Sydøstlige Norge*. Norges geologiske undersøkelse, Kristiania.
508
- 509 Bronk Ramsey, Christopher
510 2009 Bayesian Analysis of Radiocarbon Dates. *Radiocarbon* 51(1):337–360.
511 DOI:10.1017/S0033822200033865.
- 512 Conolly, James
513 2020 Spatial interpolation. In, edited by Mark Gillings, Piraye Hacıgüzeller, and Gary Lock, pp. 118–134. Routledge, London & New York.
514
- 515 Conolly, James, and Mark Lake
516 2006 *Geographical information systems in archaeology*. Cambridge University Press, Cambridge.
517
- 518 Crema, Enrico R.
519 2012 Modelling Temporal Uncertainty in Archaeological Analysis. *Journal of Archaeological Method and Theory* 19(3):440–461. DOI:10.1007/s10816-011-9122-3.
520
- 521 2015 Time and probabilistic reasoning in settlement analysis. In *Mathematics and archaeology*, edited by Juan A. Barcelo and Igor Bogdanovic, pp. 314–334. CRC Press, Boca Raton.
522
- 523 Crema, Enrico R., Andrew Bevan, and Mark W. Lake
524 2010 A probabilistic framework for assessing spatio-temporal point patterns in the archaeological record. *Journal of Archaeological Science* 37(5):1118–1130. DOI:10.1016/j.jas.2009.12.012.
525
- 526 Eskeland, Knut Fossdal
527 2017 *Rapport, arkeologisk registrering. E18 Langangen Rugtvedt, 16/06999, Porsgrunn og Bamble kommune. Skien*.
- 528
- 529 Fossum, Guro
530 2020 Specialists facing climate change. The 8200 cal BP event and its impact on the coastal settlement in the inner oslo fjord, southeast norway. In *Coastal landscapes of the mesolithic: Human engagement with the coast from the atlantic to the baltic sea*, edited by Almut Schülke, pp. 179–201. Routledge, London & New York.
531
- 532 Gjerpe, Lars Erik, and Grethe Bjørkan Bukkemoen
533 2008 Nordby 1 – toskipede hus fra neolitum-bronsealder og boplasspor fra jernalder. In *E18-prosjektet vestfold. Bind 3. Hus, boplass- og dyrkningsspor*, edited by Lars Erik Gjerpe, pp. 7–38. University of Oslo, Museum of Cultural History, Oslo.
534
- 535 Glørstad, Håkon
536 2010 *The Structure and History of the Late Mesolithic Societies in the Oslo Fjord Area 6300-3800 BC*. Bricoleur Press, Lindome.
537
- 538 2011 Historical ideal types and the transition to the late neolithic in south norway. In, edited by Christopher Prescott and Håkon Glørstad, pp. 82–99. Oxbow Books, Oxford & Oakville.
539
- 540 2016 Deglaciation, sea-level change and the Holocene colonization of Norway. *Geological Society, London, Special Publications* 411:9–25. DOI:10.1144/SP411.7.
541

-
- 542 Glørstad, Håkon, Jostein Gundersen, Frode Kvalø, Pål Nymoen, David Simpson, and Birgitte Skar
543 2020 Submerged stone age from a norwegian perspective. In, edited by Geoff Bailey, Nena Galanidou, Hans
544 Peeters, Hauke Jöns, and Moritz Mennenga, pp. 125–140. Springer, Cham.
- 545 GRASS Development Team
546 2017 *Geographic resources analysis support system (GRASS) software, version 7.2*. Open Source Geospatial
547 Foundation.
- 548 Gundersen, Jostein
549 2013 Verken fjord eller fjell – steinalderen i det kystnære innlandet. Gamle og nye funn fra notodden i
550 telemark. *Viking* 76:35–62.
- 551 Hafsten, Ulf
552 1957 De senkvartære strandlinje-forskyvningene i oslotrakten belyst ved pollenanalytiske undersøkelser.
553 *Norwegian Journal of Geography* 16(1-8):74–99. DOI:10.1080/00291955708622137.
- 554 1983 Shore-level changes in South Norway during the last 13,000 years, traced by biostrati-
555 graphical methods and radiometric datings. *Norwegian Journal of Geography* 37(2):63–79.
DOI:10.1080/00291958308552089.
- 556 Hagen, Anders
557 1963 Problemkompleks Fosna. Opphav – kontakt med kontinentale grupper – forholdet til Komsa. In
558 *Boplatsproblem vid Kattegat och Skagerack*, pp. 53–59. Göteborg och Bohusläns formminnesförening
och Institutionen för nordisk fornkunskap vid Göteborgs universitet, Gothenburg.
- 559 Helskog, Knut
560 1978 Late Holocene sea-level changes seen from prehistoric settlements. *Norwegian Journal of Geography*
561 32:111–119. DOI:10.1080/00291957808552032.
- 562 Herzog, Irmela
563 2013 The potential and limits of optimal path analysis. In *Computational approaches to archaeological
564 spaces*, edited by Andrew Bevan and Mark Lake, pp. 179–211. Left Coast Press, Walnut Creek.
- 565 Hinz, Martin, Clemens Schmid, Daniel Knitter, and Carolin Tietze
566 2021 *oxcAAR: Interface to 'OxCal' radiocarbon calibration. R package version 1.1.0*.
- 567 Hollender, Artur
568 1901 Om sveriges nivåförändringar efter människans invandring. *Geologiska Föreningen i Stockholm Förhan-
569 dlingar* 23(4):1118–1130. DOI:10.1080/00293652.1975.9965220.
- 570 Hughes, Anna L. C., Richard Gyllencreutz, Øystein S. Lohne, Jan Mangerud, and John Inge Svendsen
571 2016 The last Eurasian ice sheets – a chronological database and time-slice reconstruction, DATED-1.
572 *Boreas* 45(1):1–45. DOI:<https://doi.org/10.1111/bor.12142>.
- 573 Hyndman, Rob J
574 1996 Computing and Graphing Highest Density Regions. *The American Statistician* 50(2):120–126.
- 575 Ilves, Kristin, and Kim Darmark
576 2011 Some Critical and Methodological Aspects of Shoreline Determination: Examples from the Baltic Sea
577 Region. *Journal of Archaeological Method and Theory* 18:147–165. DOI:10.1007/s10816-010-9084-x.
- 578 Jaksland, Lasse (editor)
579 2012a *E18 Brunlanesprosjektet. Bind II. Undersøkte lokaliteter fra tidligmesolitikum*. University of Oslo,
580 Museum of Cultural History, Oslo.
- 581 (editor)
582 2012b *E18 Brunlanesprosjektet. Bind III. Undersøkte lokaliteter fra tidligmesolitikum og senere*. University
583 of Oslo, Museum of Cultural History, Oslo.
- 584 2014 Kulturhistorisk sammenstilling. In, edited by Lasse Jaksland and Per Persson, pp. 11–46. University
585 of Oslo, Museum of Cultural History, Oslo.
- 586 Jaksland, Lasse, and Per Persson (editors)
587 2014 *E18 Brunlanesprosjektet. Bind I. Forutsetninger og kulturhistorisk sammenstilling*. University of Oslo,
588 Museum of Cultural History, Oslo.

- 590
- 591 Johansen, Erling
- 592 1963 Kyst(fangst)boplassenes strandbundenhet og strandlinjekronologien. In *Boplatsproblem vid Kattegat och Skagerack*, pp. 90–92. Göteborg och Bohusläns fornminnesförening och Institutionen för nordisk fornkunskap vid Göteborgs universitet, Gothenburg.
- 593
- 594 Kjemperud, Alfred
- 595 1986 Late Weichselian and Holocene shoreline displacement in the Trondheimsfjord area, central Norway. *Boreas* 15(1):61–82. DOI:10.1111/j.1502-3885.1986.tb00744.x.
- 596
- 597 Kleppe, Else Johansen
- 598 1985 *Archaeological data on shore displacements in norway*. Norges geografiske oppmåling.
- 599
- 600 Lakens, Daniël, Anne M. Scheel, and Peder M. Isager
- 601 2018 Equivalence Testing for Psychological Research: A Tutorial. *Advances in Methods and Practices in Psychological Science* 1(2):259–269. DOI:10.1177/2515245918770963.
- 602
- 603 Lewis, Joseph
- 604 2021 Probabilistic Modelling for Incorporating Uncertainty in Least Cost Path Results: a Postdictive Roman Road Case Study. *Journal of Archaeological Method and Theory* 28(3):911–924. DOI:10.1007/s10816-021-09522-w.
- 605
- 606 Marwick, Ben, Carl Boettiger, and Lincoln Mullen
- 607 2018 Packaging data analytical work reproducibly using r (and friends). *The American Statistician* 72(1):80–88. DOI:10.1080/00031305.2017.1375986.
- 608
- 609 Mikkelsen, Egil
- 610 1975 Mesolithic in south-eastern norway. *Norwegian Archaeological Review* 8(1):1118–1130. DOI:10.1080/11035890109445866.
- 611
- 612 Milne, Glenn A
- 613 2015 Glacial isostatic adjustment. In, pp. 421–437. Wiley, Chichester.
- 614
- 615 Milne, Glenn A., W. Roland Gehrels, Chris W. Hughes, and Mark E. Tamisiea
- 616 2009 Identifying the causes of sea-level change. *Nature Geoscience* 2(7):471–478. DOI:10.1038/ngeo544.
- 617
- 618 Mjærum, Axel
- 619 2018 Hinterland discoveries: Middle Mesolithic woodland utilization and the case of the site Eidsberg, eastern Norway. *Current Swedish Archaeology* 26(1):159–188. DOI:10.37718/CSA.2018.11.
- 620
- 621 Møller, Jakob J
- 622 1987 Shoreline relation and prehistoric settlement in northern norway. *Norwegian Journal of Geography* 41:45–60. DOI:<http://dx.doi.org/10.1080/00291958708552171>.
- 623
- 624 Mörner, Nils-Axel
- 625 1976 Eustasy and geoid changes. *The Journal of Geology* 84(2):123–151. DOI:10.1086/628184.
- 626
- 627 1979 The fennoscandian uplift and late cenozoic geodynamics: Geological evidence. *GeoJournal* 3(3):287–318. DOI:10.1007/BF00177634.
- 628
- 629 Nielsen, Svein Vatsvåg
- 630 2021 Early farming in Southeastern Norway: New evidence and interpretations. *Journal of Neolithic Archaeology* 23:83–113. DOI:10.12766/jna.2021.4.
- 631
- 632 Nielsen, Svein Vatsvåg, Per Persson, and Steinar Solheim
- 633 2019 De-Neolithisation in southern Norway inferred from statistical modelling of radiocarbon dates. *Journal of Anthropological Archaeology* 53:82–91. DOI:10.1016/j.jaa.2018.11.004.
- 634
- 635 Nordqvist, Bengt
- 636 1995 The mesolithic settlement of the west coast of sweden - with special emphasis on chronology and topography of coastal settlements. In *Man and sea in the mesolithic. Coastal settlement above and below present sea level*, edited by Anders Fischer, pp. 185–196. Oxbow Books, Oxford.
- 637

- 638 1999 The Chronology of the Western Swedish Mesolithic and Late Paleolithic: Old Answers in Spite of New
639 Methods. In *The Mesolithic of Central Scandinavia*, edited by Joel Boaz, pp. 235–253. University of
Oslo, Oslo.
- 640 Nummedal, Anders
641 1923 Om flintpladsene. *Norwegian Journal of Geography* 7(2):89–141.
- 642
643 Østmo, Einar
644 1988 *Etableringen av jordbrukskultur i østfold i steinalderen*. The University Collection of National Antiquities, University of Oslo, Oslo.
645
646 2008 *Auve. En fangstboplass fra yngre steinalder på vesterøya i sandefjord. I. Den arkeologiske del*. Museum
647 of Cultural History, University of Oslo, Oslo.
648 Parnell, A. C., J. Haslett, J. R. M. Allen, C. E. Buck, and B. Huntley
649 2008 A flexible approach to assessing synchronicity of past events using Bayesian reconstructions of sedimentation history. *Quaternary Science Reviews* 27(19-20):1872–1885. DOI:10.1016/j.quascirev.2008.07.009.
650
651 Persson, Per
652 2008 Nauen 5.2 – Stenåldersboplatter och fossil åkermark. In *E18-prosjektet vestfold. Bind 2. Steinalderboplasser, boplasspor, graver og dyrkningsspor*, edited by Lars Erik Gjerpe, pp. 163–198. University of
653 Oslo, Museum of Cultural History, Oslo.
654 Prescott, Christopher
655 2020 Interpreting complex diachronic "neolithic" -period data in norway. In, edited by Kurt J Gron, Lasse
656 Sørensen, and Peter Rowley-Conwy, pp. 381–400. Oxbow Books, Oxford.
657 R Core Team
658 2021 *R: A language and environment for statistical computing*. R Foundation for Statistical Computing,
659 Vienna, Austria.
660 Reimer, Paula J., William E. N. Austin, Edouard Bard, Alex Bayliss, Paul G. Blackwell, Christopher Bronk
661 Ramsey, Martin Butzin, Hai Cheng, R. Lawrence Edwards, Michael Friedrich, Pieter M. Grootes, Thomas
662 P. Guilderson, Irka Hajdas, Timothy J. Heaton, Alan G. Hogg, Konrad A. Hughen, Bernd Kromer, Sturt
663 W. Manning, Raimund Muscheler, Jonathan G. Palmer, Charlotte Pearson, Johannes van der Plicht, Ron
664 W. Reimer, David A. Richards, E. Marian Scott, John R. Southon, Christian S. M. Turney, Lukas Wacker,
665 Florian Adolphi, Ulf Büntgen, Manuela Capano, Simon M. Fahrni, Alexandra Fogtmann-Schulz, Ronny
666 Friedrich, Peter Köhler, Sabrina Kudsk, Fusa Miyake, Jesper Olsen, Frederick Reinig, Minoru Sakamoto,
667 Adam Sookdeo, and Sahra Talamo
668 2020 The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon*
669 62(4):725–757. DOI:10.1017/RDC.2020.41.
670 Reitan, Gaute, and Inger Marie Berg-Hansen
671 2009 *Lundevågenprosjektet, delrapport 1. Sammenfattende rapport. Lunde, 6/1, 6/35 og skjolnes 7/23,*
672 *7/27, farsund kommune, vest-agder*. Oslo.
673 Roalkvam, Isak
674 2020 Algorithmic classification and statistical modelling of coastal settlement patterns in mesolithic south-
675 eastern norway. *Journal of Computer Applications in Archaeology* 3(1):288–307. DOI:<https://doi.org/10.5334/jcaa.60>.
676 Røberg, Frank Halvar N.
677 2012 *Bosettings- og aktivitetsspor. Larønningen, 221/2138. Skien, telemark*. University of Oslo, Museum of
678 Cultural History, Oslo.
679 Romundset, Anders
680 2018 Postglacial shoreline displacement in the Tvedstrand-Arendal area. In *The Stone Age Coastal
Settlement in Aust-Agder, Southeast Norway*, edited by Gaute Reitan and Lars Sundström, pp.
681 463–478. Cappelen Damm Akademisk, Oslo.
682 Romundset, Anders, Stein Bondevik, and Ole Bennike
683 2011 Postglacial uplift and relative sea level changes in Finnmark, northern Norway. *Quaternary Science
Reviews* 30(19-20):2398–2421. DOI:10.1016/j.quascirev.2011.06.007.

- 684
- 685 Romundset, Anders, Ola Fredin, and Fredrik Høgaas
- 686 2015 A Holocene sea-level curve and revised isobase map based on isolation basins from near the southern
- 687 tip of Norway:18.
- 688 Romundset, Anders, Thomas R. Lakeman, and Fredrik Høgaas
- 689 2018 Quantifying variable rates of postglacial relative sea level fall from a cluster of 24 isolation basins in
- 690 southern Norway. *Quaternary Science Reviews* 197:175–192. DOI:10.1016/j.quascirev.2018.07.041.
- 691 2019 Coastal lake records add constraints to the age and magnitude of the Younger Dryas ice-front oscillation
- 692 along the Skagerrak coastline in southern Norway. *Journal of Quaternary Science* 34(2):112–124.
- 693 DOI:<https://doi.org/10.1002/jqs.3085>.
- 694 Schmitt, Lou, Stephan Larsson, Jan Burdukiewicz, John Ziker, Krister Svedhage, Jeanette Zamon, and
- 695 Steffen Holger
- 696 2009 Chronological Insights, Cultural Change, and Resource Exploitation on the West Coast of Sweden
- During the Late Palaeolithic/Early Mesolithic Transition. *Oxford Journal of Archaeology* 28:1–27.
- DOI:10.1111/j.1468-0092.2008.00317.x.
- 697 Schülke, Almut
- 698 2020 First visit or revisit? Motivations of mobility and the use and reuse of sites in the changing coastal
- areas of mesolithic southeastern norway. In *Coastal landscapes of the mesolithic: Human engagement*
- with the coast from the atlantic to the baltic sea, edited by Almut Schülke, pp. 359–393. Routledge,
- London & New York.
- 699 Shennan, Ian
- 700 2015 Handbook of sea-level research: Framing research questions. In, pp. 3–25. Wiley, Chichester.
- 701
- 702
- 703 Sognnes, Kalle
- 704 2003 On shoreline dating of rock art. *Acta Archaeologica* 74:189–209.
- 705
- 706 Solheim, Steinar
- 707 2012 Lokal praksis og fremmed opphav. Arbeidsdeling, sosiale relasjoner og differensiering i østnorsk
- 708 tidligneolitikum. Unpublished PhD thesis, Oslo.
- 709 2020 Mesolithic coastal landscapes. Demography, settlement patterns and subsistence economy in south-
- 710 eastern norway. In, edited by Almut Schülke. Routledge, London & New York.
- 711 2021 Timing the Emergence and Development of Arable Farming in Southeastern Norway by Using
- Summed Probability Distribution of Radiocarbon Dates and a Bayesian Age Model. *Radiocarbon*:1–22.
- DOI:10.1017/RDC.2021.80.
- 712
- 713 Solheim, Steinar, and Per Persson
- 714 2018 Early and mid-Holocene coastal settlement and demography in southeastern Norway: Comparing dis-
- tribution of radiocarbon dates and shoreline-dated sites, 8500–2000 cal. BCE. *Journal of Archaeological*
- 715 *Science: Reports*:334–343. DOI:10.1016/j.jasrep.2018.03.007.
- 716 Sørensen, Rolf
- 717 1979 Late Weichselian deglaciation in the Oslofjord area, south Norway. *Boreas* 8(2):241–246. DOI:<https://doi.org/10.1111/j.1502-3885.1979.tb00806.x>.
- 718
- 719 1999 En 14C datert og dendrokronologisk kalibrert strandforksyvningskurve for sørøst-norge. Sørøst-norge.
- In, edited by Lotte Selsing and Grete Lillehammer, pp. 227–242. AmS-rapport 12A. Museum of
- Archaeology, Stavanger.
- 720
- 721 Sørensen, Rolf, Kari E Henningsmoen, Helge I Høeg, and Veronika Gålman
- 722 in Holocen vegetasjonshistorie og landhevning i sørøst-norge. In, edited by Per
- 723 press Persson and Steinar Solheim.
- 724 2014 Holocene landhevningsstudier i sørøst-norge - revidert kurve. In, edited by
- Stine Melvold and Per Persson, pp. 36–47. Portal, Kristiansand.
- 725
- 726 Sørensen, Rolf, Helge I Høeg, Kari E Henningsmoen, Göran Skog, Solveig F Labowsky, and Bjørg Stabell

-
- 727 2014 Utviklingen av det senglasiale og tidlig preboreale landskapet og vegetasjonen omkring steinalderboplassene ved pauler. In *E18 brunlanesprosjektet. Bind i. Forutsetninger og kulturhistorisk sammenstilling*, edited by Lasse Jakslund and Per Persson, pp. 171–213. University of Oslo, Museum of Cultural History, Oslo.
- 728
- 729 Stokke, Jo-Simon Frøshaug, and Gaute Reitan
- 730 2018 Kvastad A2. Lokalitet med funn fra tidlig- og mellommesolitikum og dyrkningsspor fra mellom- og senneolitikum. In, edited by Gaute Reitan and Lars Sundström, pp. 375–407. Cappelen Damm Akademisk, Oslo.
- 731
- 732 Stroeven, Arjen P., Clas Hättestrand, Johan Kleman, Jakob Heyman, Derek Fabel, Ola Fredin, Bradley W. Goodfellow, Jonathan M. Harbor, John D. Jansen, Lars Olsen, Marc W. Caffee, David Fink, Jan Lundqvist, Gunhild C. Rosqvist, Bo Strömberg, and Krister N. Jansson
- 733
- 734
- 735 2016 Deglaciation of Fennoscandia. *Quaternary Science Reviews* 147. Special Issue: PAST Gateways (Palaeo-Arctic Spatial and Temporal Gateways):91–121. DOI:10.1016/j.quascirev.2015.09.016.
- 736
- 737 Svendsen, John Inge, and Jan Mangerud
- 738 1987 Late Weichselian and holocene sea-level history for a cross-section of western Norway. *Journal of Quaternary Science* 2(2):113–132. DOI:10.1002/jqs.3390020205.
- 739
- 740 Tallavaara, Miikka, and Petro Pesonen
- 741 2020 Human ecodynamics in the north-west coast of Finland 10,000–2000 years ago. *Quaternary International* 549:26–35. DOI:10.1016/j.quaint.2018.06.032.
- 742
- 743 Wang, Ian
- 744 2019 *topoDistance: Calculating topographic paths and distances. R package version 1.0.1.* [Https://CRAN.R-project.org/package=topoDistance](https://CRAN.R-project.org/package=topoDistance).
- 745
- 746 Wenn, Camilla Cecilie
- 747 2012 *Bosettingsspor, produksjonsområde og dyrkningsspor fra neolitikum til folkevandringstid. Bratsberg, 63/69, 244. Skien kommune, telemark.* University of Oslo, Museum of Cultural History, Oslo.
- 748
- 749 Wikell, Roger, Fredrik Molin, and Mattias Pettersson
- 750 2009 The archipelago of eastern middle sweden - mesolithic settlement in comparison with 14C and shoreline dating. In *Chronology and evolution within the mesolithic of north-west europe*, edited by Philippe Crombé, Mark van Strydonck, Joris Sergant, Mathieu Boudin, and Machteld Bats, pp. 417–434. Cambridge Scholar Publishing, Brussels.
- 751
- 752 Yubero-Gómez, María, Xavier Rubio-Campillo, and Javier López-Cachero
- 753 2016 The study of spatiotemporal patterns integrating temporal uncertainty in late prehistoric settlements in northeastern spain. *Archaeological and Anthropological Sciences* 8(3):477–490. DOI:10.1007/s12520-015-0231-x.
- 754