

shoredat: An R package for shoreline dating coastal Stone Age sites in south-eastern Norway

Summary

As a result of glacio-isostatic rebound, large regions of Fennoscandia have undergone a process of relative sea-level fall following the retreat of the Fennoscandian Ice Sheet. Furthermore, coastal Stone Age sites in the region appear to have been predominantly located on or close to the shoreline when they were in use. This can be combined with a reconstruction of past relative sea-level change to assign an approximate date to when the sites were in use, based on their altitude relative to the present-day sea-level. This method, called shoreline dating, has been used in the region since the early 1900s (e.g. Brøgger 1905) and is still widely applied today (e.g. Solheim and Persson 2018; Manninen et al. 2021).

Statement of need

shoredat is an R package for shoreline dating Stone Age sites on the coast of south-eastern Norway, based on local geological reconstructions of past relative sea-level change. Drawing on an empirically derived estimate of the likely elevation of the sites above sea-level when they were in use, the method for shoreline dating implemented in the package was recently published in Roalkvam (2023). No open-source software with which to perform shoreline dating has previously been available, and the only closed-source software available is **sealev** from the University of Tromsø, Tromsø Geophysical Observatory (<https://www.tgo.uit.no/sealev/>, see Møller 2003), which provides non-probabilistic point estimates of shoreline dates based on data last updated in 2002.

shoredat is aimed at providing researchers and students dealing with the coastal Stone Age of south-eastern Norway with a formalised tool for performing and handling shoreline dates. This complements already available software for handling radiocarbon dates and other sources of temporal data, such as the R packages **rcarbon** (Crema and Bevan 2021), **bchron** (Haslett and Parnell 2008), **oxcAAR** (Hinz et al. 2021), **kairos** (Frerebeau 2022) and **ArchaeoPhases** (Philippe and Vibet 2020), as well as proprietary software such as **OxCal** (Bronk Ramsey 2009).

Shoreline dating is frequently applied in the research and cultural resource management sectors in Norway, both to plan archaeological investigations and for establishing temporal frameworks with which to approach the archaeological material. Case-studies employing **shoredat** are currently being undertaken. Furthermore, future archaeological material can be drawn on to further test the method as it is implemented here, and potentially lead to adjustments in how it could be applied in a given setting.

Spatial and temporal coverage

As the method of shoreline dating is dependent on reliable reconstructions of relative sea-level change, the package is at present limited to being applicable in the coastal region between Horten in the north east to Arendal in the south west (Figure 1). Geologically derived displacement curves from this region have recently been compiled for Skoppum in Horten (Romundset 2021), Gunnarsrød in Porsgrunn (Sørensen et al. 2023), Hanto in Tvedestrand (Romundset, Lakeman, and Høgaas 2018) and Bjørnebu in Arendal (Romundset 2018).

The spatial coverage of `shoredate` will be extended to surrounding regions as forthcoming data on shoreline displacement becomes available.

As the latest start date of the displacement curves is 9469 BCE, and as the oldest known anthropogenic activity in Norway dates to around 9300 BCE (Glørstad 2016), 9469 BCE marks the lower temporal limit of the package. Furthermore, in Roalkvam (2023) it was found that sites tend to be more withdrawn from the shoreline after c. 2500 BCE, which thus marks the upper temporal limit for shoreline dating with the package.

Example of base functionality

To shoreline date a site, this has to be provided as a spatial object of class `sf` from the `sf` package (Pebesma 2018), and be set to the coordinate reference system WGS 84 / UTM zone 32N (EPSG:32632). The elevation of the site above present sea-level must be provided when running `shoreline_date()` by either manually specifying it, or by providing an elevation raster of class `SpatRaster` from the `terra` package (Hijmans et al. 2022), from where the site elevation is derived. When calling `shoreline_date()`, the trajectory of shoreline displacement at the location of the site is interpolated under the hood with the function `interpolate_curve()`, using inverse distance weighting. This is based on the distance between the site and the isobases of the displacement curves.

Figure 2 shows the location of an example site, plotted by passing it to `target_plot()`. Figure 3 displays the result of running the command `interpolate_curve()` on the example site, and plotting the resulting interpolated displacement curve with `displacement_plot()`. Finally, Figure 4 shows the result of dating the example site with `shoreline_date()` when manually specifying that the site is situated at 58.8m above present sea-level. The resulting date is plotted with the function `shoredate_plot()`.

Acknowledgements

I owe great thanks to David Wright, Anders Romundset, Ingrid Fuglestad, Per Persson, Steinar Solheim and Hallvard Bruvoll for valuable feedback during work with this project.

References

- Brøgger, Waldemar Christofer. 1905. *Strandliniens Beliggenhed under Stenalderen i Det Sydøstlige Norge*. Kristiania: Geological Survey of Norway.
- Bronk Ramsey, Christopher. 2009. “Bayesian Analysis of Radiocarbon Dates.” *Radiocarbon* 51 (1): 337–60. <https://doi.org/10.1017/S0033822200033865>.
- Crema, Enrico R., and Andrew Bevan. 2021. “Inference from Large Sets of Radiocarbon Dates: Software and Methods.” *Radiocarbon* 63: 23–39. <https://dx.doi.org/10.1017/RDC.2020.95>.
- Frerebeau, Nicolas. 2022. *Kairos: Analysis of Chronological Patterns from Archaeological Count Data*. Pessac, France: Université Bordeaux Montaigne. <https://doi.org/10.5281/zenodo.5653896>.
- Glørstad, Håkon. 2016. “Deglaciation, Sea-Level Change and the Holocene Colonization of Norway.” *Geological Society, London, Special Publications* 411: 9–25. <https://doi.org/10.1144/SP411.7>.
- Haslett, John, and Andrew C Parnell. 2008. “A Simple Monotone Process with Application to Radiocarbon-Dated Depth Chronologies.” *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 57 (4): 399–418. <https://rss.onlinelibrary.wiley.com/doi/full/10.1111/j.1467-9876.2008.00623.x>.
- Hijmans, Robert J, Roger Bivand, Edzer Pebesma, and Michael D Sumner. 2022. “Terra: Spatial Data Analysis.” <https://CRAN.R-project.org/package=terra>.
- Hinz, Martin, Clemens Schmid, Daniel Knitter, and Carolin Tietze. 2021. “oxcAAR: Interface to ‘OxCal’ Radiocarbon Calibration. R Package Version 1.1.0.” <https://CRAN.R-project.org/package=oxcAAR>.




Figure 1: The spatial and temporal coverage of the package. The first figure displays the location of the spatial extent in south-eastern Norway. The second figure displays the location of the isobases, representing contours along which the shoreline displacement have followed the same trajectory. The isobases correspond to the displacement curves in the third figure, where the temporal limits are marked with dashed lines.




Figure 2: The location of the example site relative to the isobases of the displacement curves. The basemap is a simplified and lightweight map of the region.




Figure 3: The curve interpolated to the example site by means of inverse distance weighting. This is based on the distance between the site and the isobases of the geologically derived displacement curves.




Figure 4: The resulting shoreline date for the example site. The blue gamma distribution on the y-axis indicates the likely elevation of the site above sea-level when it was occupied. The red envelope is the interpolated shoreline displacement curve for the site location. The resulting shoreline date in grey is the result of transferring the probability from the gamma distribution to the calendar scale by coupling it with the displacement curve. The date is underlined with the 95% highest density region (HDR) in black.

- Manninen, Mikael A, Hege Damlien, Jan Ingolf Kleppe, Kjel Knutsson, Anton Murashkin, Anja R Niemi, Carine S Rosenvinge, and Per Persson. 2021. “First Encounters in the North: Cultural Diversity and Gene Flow in Early Mesolithic Scandinavia.” *Antiquity* 95: 310–28. <https://doi.org/10.15184/aqy.2020.252>.
- Møller, Jakob J. 2003. “Late Quaternary Sea Level and Coastal Settlement in the European North.” *Journal of Coastal Research* 19: 731–37. <https://www.jstor.org/stable/4299209>.
- Pebesma, Edzer. 2018. “Simple Features for R: Standardized Support for Spatial Vector Data.” *The R Journal* 10 (1): 439–46. <https://doi.org/10.32614/RJ-2018-009>.
- Philippe, Anne, and Marie-Anne Vibet. 2020. “Analysis of Archaeological Phases Using the R Package ArchaeoPhases.” *Journal of Statistical Software, Code Snippets* 93 (1): 1–25. <https://doi.org/10.18637/jss.v093.c01>.
- Roalkvam, Isak. 2023. “A Simulation-Based Assessment of the Relation Between Stone Age Sites and Relative Sea-Level Change Along the Norwegian Skagerrak Coast.” *Quaternary Science Reviews* 299: 107880. <https://doi.org/10.1016/j.quascirev.2022.107880>.
- Romundset, Anders. 2018. “Postglacial shoreline displacement in the Tvedestrand-Arendal area.” In *The Stone Age Coastal Settlement in Aust-Agder, Southeast Norway*, edited by Gaute Reitan and Lars Sundström, 463–78. Oslo: Cappelen Damm Akademisk.
- . 2021. *Resultater fra NGUs undersøkelse av etteristidas strandforskyvning nord i Vestfold*. Trondheim: Geological Survey of Norway.
- Romundset, Anders, Thomas R. Lakeman, and Fredrik Høgaas. 2018. “Quantifying Variable Rates of Postglacial Relative Sea Level Fall from a Cluster of 24 Isolation Basins in Southern Norway.” *Quaternary Science Reviews* 197 (October): 175–92. <https://doi.org/10.1016/j.quascirev.2018.07.041>.
- Solheim, Steinar, and Per Persson. 2018. “Early and Mid-Holocene Coastal Settlement and Demography in Southeastern Norway: Comparing Distribution of Radiocarbon Dates and Shoreline-Dated Sites, 8500–2000 Cal. BCE.” *Journal of Archaeological Science: Reports* 19: 334–43. <https://doi.org/10.1016/j.jasrep.2018.03.007>.
- Sørensen, Rolf, Kari E Henningsmoen, Helge I Høeg, and Veronika Gälman. 2023. “Holocen vegetasjonshistorie og landhevning i søndre Vestfold og sørøstre Telemark.” In *The Stone Age in Telemark. Archaeological Results and Scientific Analysis from Vestfoldbaneprosjektet and E18 Rugtvedt–Dørdal*, edited by Persson Per and Solheim Steinar. University of Oslo, in press.