## Cod (Gadus morhua) in subareas 1 and 2 (Northeast Arctic)

## ICES advice on fishing opportunities

ICES advises that when the Joint Norwegian–Russian Fisheries Commission (JRNFC) management plan is applied, catches in 2022 should be no more than 708 480 tonnes.

## Stock development over time

Fishing pressure on the stock is at F<sub>MSY</sub> between F<sub>pa</sub> and F<sub>lim</sub> and spawning-stock size is above MSY B<sub>trigger</sub>, B<sub>pa</sub>, and B<sub>lim</sub>.




Figure 1<sup>†</sup> Cod in subareas 1 and 2 (Northeast Arctic). Catch, recruitment, F, and SSB. For this stock, F<sub>MSY</sub> and F<sub>MGT</sub> ranges from 0.40 to 0.60, and there are three SSB<sub>MGT</sub> values (460 000, 920 000 and 1 380 000 tonnes) which are not shown.

### **Catch scenarios**

**Table 1**<sup>‡</sup> Cod in subareas 1 and 2 (Northeast Arctic). Values in the forecast and for the interim year.

Variable	Value	Notes		
F <sub>ages 5-10</sub> (2021)	0.65	Based on a catch of 885 600 t for 2021.		
SSB (2022)	696 371	From assessment; tonnes		
R <sub>age 3</sub> (2021)	557 000	From recruitment model; thousands		
R <sub>age 3</sub> (2022)	620 000	From recruitment model; thousands		
Rage 3 (2023)	547 000	From recruitment model; thousands		
Total catch (2021)	885 600	TAC; tonnes		

<sup>&</sup>lt;sup>†</sup> Version 2: Figure changed. The assessment was revised due to discovery of errors in the calculation of some of the survey indices for 2021 used in the assessment (Joint Norwegian-Russian survey Barents Sea, Feb–Mar (G5348) bottom trawl (BTr) and acoustic (Aco) indices).

<sup>&</sup>lt;sup>‡</sup> Version 2: Values changed.

Table 2<sup>§</sup> Cod in subareas 1 and 2 (Northeast Arctic). Annual catch scenarios. All weights are in tonnes.

Table 2 Cou in subdices 1 and 2 (Northeast Arctic). Annual catch scenarios. An weights are in tonnes.									
Basis	Total catch (2022)	Ftotal (2022)	SSB (2023)	% SSB change *	% TAC change **	% Advice change  **			
ICES advice basis									
Management plan^	708 480	0.62	578 966	-17	-20	-20			
Other scenarios									
MSY approach: F <sub>MSY</sub>	504 244	0.40	720 844	4	-43	-43			
F = 0	0	0	1 103 288	58	-100	-100			
$F = F_{2020}$	735 105	0.65	561 135	-19	-17	-17			
F <sub>pa</sub>	504 244	0.40	720 844	4	-43	-43			
F <sub>lim</sub>	810 202	0.74	511 742	-27	-9	-9			

<sup>\*</sup> SSB 2023 relative to SSB 2022.

The advice for 2022 is 20% lower than the advice for 2021 due to a declining stock trend and the application of the 20% TAC change constraint.

#### Basis of the advice

**Table 3** Cod in subareas 1 and 2 (Northeast Arctic). The basis of the advice.

Advice basis	Joint Russian–Norwegian Fisheries Commission management plan
	At the 46th meeting of the Joint Russian–Norwegian Fisheries Commission (JRNFC) in October 2016, the previously used management plan was amended, and the current plan is as follows:
	The TAC is calculated as the average catch predicted for the coming three years, using the target level of exploitation ( $F_{\rm tr}$ ).
	The target level of exploitation is calculated according to the spawning-stock biomass (SSB) in the first year of the forecast as follows:
Management plan	- if SSB < B <sub>pa</sub> , then F <sub>tr</sub> = SSB/B <sub>pa</sub> × F <sub>MSY</sub> ; - if B <sub>pa</sub> ≤ SSB ≤ 2×B <sub>pa</sub> , then F <sub>tr</sub> = F <sub>MSY</sub> ; - if 2 × B <sub>pa</sub> < SSB < 3 × B <sub>pa</sub> , then F <sub>tr</sub> = F <sub>MSY</sub> × (1 + 0.5 × (SSB – 2 × B <sub>pa</sub> )/B <sub>pa</sub> ); - if SSB ≥ 3 × B <sub>pa</sub> , then F <sub>tr</sub> = 1.5 × F <sub>MSY</sub> ; where F <sub>MSY</sub> = 0.40 and B <sub>pa</sub> = 460 000 tonnes. If the SSB in the present year, the previous year, and each of the three years of prediction is above B <sub>pa</sub> , the TAC should not be changed by more than ±20% compared with the previous year's TAC. In this case, however,
	F <sub>tr</sub> should not be below 0.30.
	In 2014, JNRFC decided that from 2015 onwards, Norway and Russia can transfer to or borrow from the following year up to 10% of their country's quota.
	ICES evaluated this harvest control rule in 2016 (ICES, 2016a) and 2021 (ICES, 2021a) and concluded that it is precautionary.

## Quality of the assessment

Input data and assessment model settings were changed at the benchmark (ICES, 2021a), and this resulted in a downward revision of SSB.

ICES Advice 2021

<sup>\*\*</sup> Advice value for 2022 relative to TAC and advice value for 2021 (885 600 tonnes).

 $<sup>^{\</sup>circ}$  Since SSB in 2022 is between B<sub>pa</sub> = 460 000 tonnes and 2 × B<sub>pa</sub> = 920 000 tonnes, F = 0.40 is used in the 3-year prediction, giving catches of 504 244, 528 049 and 557 768 tonnes in 2022, 2023, and 2024, respectively. The average of this is 530 020 tonnes. According to the harvest control rule (HCR), the maximum decrease in TAC is limited by 20%, giving a catch of 708 480 tonnes, which corresponds to an F of 0.62 in 2022.

<sup>§</sup> Version 2: Values for 'Other scenarios' changed.

All surveys, in particular the ecosystem survey, indicate a decrease in abundance from 2020 to 2021 that is larger than expected and therefore the assessment is more uncertain than usual.




Figure 2\*\* Cod in subareas 1 and 2 (Northeast Arctic). Historical assessment results. There was a benchmark revision in 2021.

#### Issues relevant for the advice

Fisheries targeting Northeast Arctic (NEA) cod take as bycatch a considerable part of the total golden redfish (*Sebastes norvegicus*) catch, and this bycatch is still above any sustainable catch level. Measures to minimize bycatch levels are essential.

Bycatch of coastal cod should be kept as low as possible in order to promote rebuilding of the coastal cod (*Gadus morhua*) stock.

The advice this year is considerably lower than last year. The stock abundance is revised downwards due to both a downward revision following the benchmark (ICES, 2021a) and to unexpectedly low survey indices.

The advice for 2022 is robust to the intermediate year catch assumption due to the 20% constraint on annual TAC change.

#### **Reference points**

**Table 4** Cod in subareas 1 and 2 (Northeast Arctic). Reference points, values, and their technical basis.

Framework	Reference point	Value	Technical basis	Source
MSY approach	MSY B <sub>trigger</sub>	460 000 t	B <sub>pa</sub> and trigger point in HCR	ICES (2003, 2021a)
тизт арргоаст	F <sub>MSY</sub>	0.40-0.60	Long-term simulations	ICES (2003, 2021a)
	B <sub>lim</sub>	220 000 t	Change point regression	ICES (2003, 2021a)
Precautionary	B <sub>pa</sub>	460 000 t	The lowest SSB estimate having > 90% probability of remaining above B <sub>lim</sub>	ICES (2003, 2021a)
approach	Flim	0.74	F corresponding to an equilibrium stock = B <sub>lim</sub>	ICES (2003, 2021a)
	F <sub>pa</sub>	0.40	The highest F estimate having > 90% probability of remaining below F <sub>lim</sub>	ICES (2003, 2021a)
Management	SSB <sub>mgt</sub>	460 000 t	Two-step (double hockey stick) HCR, see Table 3	ICES (2017)
plan	F <sub>mgt</sub>	0.40-0.60	Two-step (double hockey stick) HCR, see Table 3	ICES (2017)

<sup>\*\*</sup> Version 2: Figure changed.

ICES Advice 2021

## Basis of the assessment

 Table 5
 Cod in subareas 1 and 2 (Northeast Arctic). Basis of the assessment and advice.

ICES stock data category	1 ( <u>ICES, 2021b</u> ).
Assessment type	Age-based analytical assessment (SAM; ICES, 2021c) that uses catches in the model and in the forecast
Input data	Commercial catches (international landings, ages and length frequencies from catch sampling); four survey indices (Joint Norwegian-Russian survey Barents Sea, Feb–Mar (G5348) bottom trawl (BTr) and acoustic (Aco) indices; Russian bottom trawl survey, Oct–Dec (G5348); Joint Norwegian-Russian Ecosystem survey (A5216; Btr)); annual maturity data from the four surveys; natural mortalities from annual stomach sampling.
Discards and bycatch	Discarding is considered negligible in recent years (below 5%). Bycatch is included.
Indicators	None.
Other information	Last benchmarked in February 2021 (ICES, 2021a).
Working group	Arctic Fisheries Working Group (AFWG).

## History of the advice, catch, and management

Table 6 Cod in subareas 1 and 2 (Northeast Arctic). ICES advice, agreed TACs, the official and unreported landings, and ICES catches. All weights are in tonnes.

	catches. All weights are in ton	1163.				
Year	ICES advice	Catch corresponding to advice	Agreed TAC	Official catches	Unreported landings (included in ICES catches)	ICES catches
1987	Gradual reduction in F	595000	560000	552000		523071
1988	F = 0.51; TAC (Advice November 1987, revised advice May 1988)	530000 (320000– 360000)	590000 (451000)	459000		434939
1989	Large reduction in F	335000	300000	348000		332481
1990	F at Flow; TAC	172000	160000	210000	25000	212000
1991	F at Flow; TAC	215000	215000	294000	50000	319158
1992	Within safe biological limits	250000	356000	421000	130000	513234
1993	Healthy stock	256000	500000	575000	50000	581611
1994	No long-term gains in increased F	649000	700000	795000	25000	771086
1995	No long-term gains in increased F	681000	700000	763000		739999
1996	No long-term gains in increased F	746000	700000	759000		732228
1997	Well below F <sub>med</sub>	< 993000	850000	792000		762403
1998	F less than F <sub>med</sub>	514000	654000	615000		592624
1999	Reduce F to below F <sub>pa</sub>	360000	480000	506000		484910
2000	Increase B above B <sub>pa</sub> in 2001	110000	390000			414870
2001	High probability of SSB > B <sub>pa</sub> in 2003	263000	395000			426471
2002	Reduce F to well below 0.25	181000	395000		90000	535045
2003	Reduce F to below F <sub>pa</sub>	305000	395000		115000	551990
2004	Reduce F to below F <sub>pa</sub>	398000	486000		117000	606445
2005	Take into account coastal cod and redfish bycatches; apply catch rule	485000	485000		166000	641276
2006	Take into account coastal cod and redfish bycatches; apply amended catch rule	471000	471000		67100	537642
2007	Take into account coastal cod and redfish bycatches; Fpa	309000	424000		41087	486883
2008	Take into account coastal cod and redfish bycatches; apply catch rule	409000	430000		15000	464171
2009	Take into account coastal cod and redfish bycatches; apply catch rule	473000	525000		0	523431
2010	Take into account coastal cod and redfish bycatches; apply catch rule.	577500	607000		0	609983
2011	Take into account coastal cod and redfish bycatches; apply catch rule	703000	703000		0	719829

Year	ICES advice	Catch corresponding to advice	Agreed TAC	Official catches	Unreported landings (included in ICES catches)	ICES catches
2012	Take into account coastal cod and redfish bycatches; apply catch rule	751000	751000		0	727663
2013	Take into account coastal cod and <i>S. marinus</i> ^^ bycatches; apply catch rule	940000	1000000		0	966209
2014	Take into account coastal cod and <i>S. marinus</i> ^^ bycatches; apply catch rule	993000	993000		0	986449
2015	Take into account coastal cod and <i>S. norvegicus</i> bycatches. Apply catch rule.	894000	894000		0	864384
2016	Take into account coastal cod and <i>S. norvegicus</i> bycatches; apply catch rule	805000	894000		0	849422
2017	Take into account coastal cod and <i>S. norvegicus</i> bycatches; apply management plan	≤ 805000	890000 ^		0	868276
2018	Take into account coastal cod and <i>S. norvegicus</i> bycatches; apply management plan	712000	775000		0	778627
2019	Take into account coastal cod and <i>S. norvegicus</i> bycatches; apply management plan	674678	725000		0	692609
2020	Apply management plan	≤ 689672	738000		0	692903
2021	Apply management plan	≤ 885600	885600			
2022	Apply management plan	≤ 708480				

<sup>^</sup> The 2017 TAC was set according to the management plan agreed by JNRFC in October 2016.

## History of the catch and landings

Table 7 Cod in subareas 1 and 2 (Northeast Arctic). Catches inside and outside the NEAFC Regulatory Area (RA) as estimated by ICES.

Year	Inside the NEAFC RA (tonnes)	Outside the NEAFC RA (tonnes)	Total catches (tonnes)	Proportion inside the NEAFC RA (%)	
2018	1726	776901	778627	0.22%	
2019	1094	691515	692609	0.16%	
2020	1616	691296	692903	0.23%	

Table 8 Cod in subareas 1 and 2 (Northeast Arctic). Catch distribution by fleet in 2020 as estimated by ICES.

Catch (2020)	Landir	Discards	
	74% demersal trawls	Unknown, but discarding	
692 903 tonnes	692 903 t	is considered to be	
	092 903 1	offiles	negligible

<sup>^^</sup> Until 2014 this species was named Sebastes marinus, thereafter Sebastes norvegicus.

Table 9 Cod in subareas 1 and 2 (Northeast Arctic). History of commercial landings by country. All weights are in tonnes.

Table 9	9 Cod in subareas 1 and 2 (Northeast Arctic). History of commercial landings by country. All weights are in tonnes.								tonnes.				
Year	Faroe Islands	France	German Dem.Rep.	Fed.Rep.Germany	Greenland	Iceland	Norway	Poland	United Kingdom	Russia**	Spain	Others	Total
1961	3934	13755	3921	8129			268377	-	158113	325780		1212	783221
1962	3109	20482	1532	6503			225615	-	175020	476760		245	909266
1963	-	18318	129	4223			205056	108	129779	417964		-	775577
1964	-	8634	297	3202			149878	-	94549	180550		585	437695
1965	-	526	91	3670			197085	-	89962	152780		816	444930
1966	-	2967	228	4284			203792	-	103012	169300		121	483704
1967	-	664	45	3632			218910	-	87008	262340		6	572605
1968	-	-	225	1073			255611	-	140387	676758		-	1074084
1969	29374	-	5907	5543			305241	7856	231066	612215		133	1197226
1970	26265	44245	12413	9451			377606	5153	181481	276632		-	933246
1971	5877	34772	4998	9726			407044	1512	80102	144802		215	689048
1972	1393	8915	1300	3405			394181	892	58382	96653		166	565287
1973	1916	17028	4684	16751			285184	843	78808	387196		276	792686
1974	5717	46028	4860	78507			287276	9898	90894	540801		38453	1102434
1975	11309	28734	9981	30037			277099	7435	101843	343580		19368	829377
1976	11511	20941	8946	24369			344502	6986	89061	343057		18090	867463
1977	9167	15414	3463	12763			388982	1084	86781	369876		17771	905301
1978	9092	9394	3029	5434			363088	566	35449	267138		5525	698715
1979	6320	3046	547	2513			294821	15	17991	105846		9439	440538
1980	9981	1705	233	1921			232242	3	10366	115194		8789	380434
1981	12825	3106	298	2228			277818		5262	83000	14500	-	399037
1982	11998	761	302	1717			287525		6601	40311	14515	-	363730
1983	11106	126	473	1243			234000		5840	22975	14229	-	289992
1984	10674	11	686	1010			230743		3663	22256	8608	-	277651
1985	13418	23	1019	4395			211065		3335	62489	7846	4330	307920
1986	18667	591	1543	10092			232096		7581	150541	5497	3505	430113
1987	15036	1	986	7035			268004		10957	202314	16223	2515	523071
1988	15329	2551	605	2803			223412		8107	169365	10905	1862	434939
1989	15625	3231	326	3291			158684		7056	134593	7802	1273	332481
1990	9584	592	169	1437			88737		3412	74609	7950	510	187000
1991	8981	975		2613			126226		3981	119427***	3677	3278	269158
1992	11663	2		3911	3337		168460		6120	182315	6217	1209	383234
1993	17435	3572		5887	5389	9374	221051		11336	244860	8800	3907	531611
1994	22826	1962		8283	6882	36737	318395		15579	291925	14929	28568	746086
1995	22262	4912		7428	7462	34214	319987		16329	296158	15505	15742	739999
1996	17758	5352		8326	6529	23005	319158		16061	305317	15871	14851	732228
1997	20076	5353		6680	6426	4200	357825		18066	313344	17130	13303	762403
1998	14290	1197		3841	6388	1423	284647		14294	244115	14212	8217	592624
1999	13700	2137		3019	4093	1985	223390		11315	210379	8994	5898	484910
2000	13350	2621		3513	5787	7562	192860		9165	166202	8695	5115	414870
2001	12500	2681		4524	5727	5917	188431		8698	183572	9196	5225	426471
2002	15693	2934		4517	6419	5975	202559		8977	184072	8414	5484	445045
2003	19427	2921		4732	7026	5963	191977		8711	182160	7924	6149	436990
2004	19226	3621		6187	8196	7201	212117		14004	201525	11285	6082	489445
2005	16273	3491		5848	8135	5874	207825		10744	200077	9349	7660	475276
2006	16327	4376		3837	8164	5972	201987		10594	203782	9219	6271	470527

Year	Faroe Islands	France	German Dem.Rep.	Fed.Rep.Germany	Greenland	Iceland	Norway	Poland	United Kingdom	Russia**	Spain	Others	Total
2007	14788	3190		4619	5951	7316	199809		9298	186229	9496	5101	445796
2008	15812	3149		4955	5617	7535	196598		8287	190225	9658	7336	449171
2009	16905	3908		8585	4977	7380	224298		8632	229291	12013	7442	523431
2010	15977	4499		8442	6584	11299	264701		9091	267547	12657	9185	609983
2011	13429	1173		4621	7155	12734	331535		8210	310326	13291	17354^	719829
2012	17523	2841		8500	8520	9536	315739		11166	329943	12814	11081	727663
2013	13833	7858		8010	7885	14734	438734		12536	432314	15042	15263	966209
2014	33298	8149		6225	10864	18205	431846		14762	433479	16378	13243	986449
2015	26568	7480		6427	7055	16120	377983		11778	381778	19905	9880	864384
2016	24084	7946		6336	8607	16031	348949		13583	394107	14640	15139	849422
2017	28637	9554		5977	13638	11925	357419		16731	396180	14414	13802	868276
2018	26152	6605		9768	12743	10708	333539		11533	340364	13143	14071	778627
2019	22270	6371		8470	7553	12294	282120		11214	316813	13939	11565	692609
2020*	21679	5796		9725	7391	9734	289472		12113	312683	11403	12908	692903

<sup>\*</sup> Provisional figures.

ICES Advice 2021 7

<sup>\*\*</sup> USSR prior to 1991. \*\*\* Includes Baltic countries.

<sup>^</sup> Includes unspecified EU catches.

# Summary of the assessment

Table 10<sup>††</sup> Cod in subareas 1 and 2 (Northeast Arctic). Assessment summary. High and low refer to 95% confidence bounds.

Table 10 <sup>††</sup>	Cod in su	bareas 1 and 2 (	Northeast Arcti	rctic). Assessment summary. High and low refer to 95% confidence bounds.							
		Recruitment		Spawn	ing-stock bi	omass	Total	Fishin	g morta	lity	
Year	Recruitment (Age 3)	Low	High	SSB	Low	High	Total catch	F (ages	Low	High	
		thousands			tonnes		tonnes	5–10)			
1946	1135995	775392	1664300	951351	798462	1133515	706000	0.25	0.21	0.30	
1947	582175	402972	841070	903141	766027	1064797	882017	0.31	0.27	0.36	
1948	438817	301320	639054	784918	661093	931937	774295	0.35	0.30	0.40	
1949	625639	436782	896156	595063	510613	693481	800122	0.37	0.32	0.42	
1950	1026124	718983	1464473	535994	470419	610709	731982	0.38	0.34	0.44	
1951	2444083	1715479	3482143	494938	439423	557466	827180	0.41	0.36	0.47	
1952	2342824	1660042	3306439	489050	430962	554966	876795	0.46	0.40	0.52	
1953	2420556	1718678	3409069	411898	360751	470297	695546	0.41	0.36	0.47	
1954	831335	591130	1169146	407947	360167	462066	826021	0.44	0.39	0.50	
1955	383577	272584	539765	328217	294202	366166	1147841	0.52	0.46	0.59	
1956	746701	531949	1048148	281774	253769	312868	1343068	0.57	0.50	0.64	
1957	1428657	1020201	2000647	212407	190851	236397	792557	0.53	0.47	0.60	
1958	937176	671425	1308110	205291	182597	230807	769313	0.53	0.47	0.59	
1959	1314632	945618	1827649	434184	384376	490445	744607	0.55	0.49	0.62	
1960	1483174	1066675	2062301	384276	337777	437177	622042	0.54	0.48	0.61	
1961	1554231	1116484	2163608	386369	342627	435696	783221	0.63	0.57	0.71	
1962	1252301	898767	1744901	315437	283260	351270	909266	0.74	0.66	0.83	
1963	900871	642811	1262530	216360	194726	240397	776337	0.81	0.72	0.91	
1964	468188	331586	661064	200613	179723	223931	437695	0.68	0.60	0.76	
1965	870801	618860	1225309	108004	95946	121577	444930	0.58	0.51	0.65	
1966	1842798	1317206	2578112	120913	108626	134590	483711	0.55	0.49	0.62	
1967	1311651	937444	1835231	128608	115457	143258	572605	0.56	0.50	0.63	
1968	183682	131056	257440	222811	202522	245132	1074084	0.60	0.54	0.67	
1969	110464	78751	154948	149052	134259	165475	1197226	0.71	0.63	0.79	
1970	205677	146379	288996	242287	218015	269260	933246	0.70	0.62	0.78	
1971	402694	287919	563224	330586	293512	372343	689048	0.65	0.58	0.73	
1972	1046188	753726	1452130	353299	311265	401009	565254	0.66	0.59	0.74	
1973	1723234	1235822	2402884	334009	290079	384592	792685	0.63	0.56	0.71	
1974	568075	412932	781506	158891	135418	186433	1102433	0.61	0.55	0.69	
1975	608747	442105	838199	133454	119319	149264	829377	0.66	0.59	0.73	
1976	606891	437764	841359	167175	151435	184552	867463	0.71	0.63	0.78	
1977	372774	271228	512338	336183	299424	377455	905301	0.82	0.73	0.91	
1978	622702	451059	859659	228073	199602	260604	698715	0.85	0.77	0.95	
1979	202676	146812	279797	180482	157416	206927	440538	0.77	0.69	0.86	
1980	130336	96706	175662	108430	96660	121633	380434	0.76	0.68	0.85	
1981	143894	108777	190348	161309	145938	178300	399038	0.79	0.71	0.88	
1982	183757	141725	238257	321072	287930	358029	363730	0.78	0.70	0.86	
1983	141573	109202	183541	311254	279887	346136	289992	0.79	0.72	0.88	
1984	442660	345055	567875	243541	222162	266976	277651	0.87	0.79	0.96	
1985	534453	426408	669874	195202	178120	213922	307920	0.81	0.73	0.90	
1986	1375465	1084974	1743732	164279	150132	179759	430113	0.87	0.79	0.96	
1987	360098	283602	457228	115232	104641	126894	523071	0.92	0.84	1.02	
1988	335686	264970	425274	191408	172832	211980	434939	0.88	0.79	0.98	
1989	157733	125676	197968	236934	211880	264951	332481	0.67	0.59	0.75	
1990	130259	102010	166331	300702	263370	343327	212000	0.43	0.37	0.49	
1991	296072	233113	376035	632183	560439	713112	319158	0.41	0.36	0.47	
1992	716248	571817	897162	801556	718826	893808	513234	0.49	0.43	0.55	
1993	988663	789520	1238038	701373	633832	776112	581611	0.59	0.53	0.65	
1994	752644	600198	943811	571967	521464	627361	771086	0.75	0.67	0.83	

<sup>††</sup> Version 2: Values changed.

	Recruitment			Spawning-stock biomass			Total	Fishing mortality		
Year	Recruitment (Age 3)	Low	High	SSB	Low	High	catch	F (ages	Low	High
	thousands			tonnes			tonnes	5–10)		
1995	539502	430537	676044	534275	486310	586972	739999	0.77	0.69	0.85
1996	407340	324138	511898	550559	496305	610744	732228	0.79	0.71	0.88
1997	785440	624690	987554	545333	487063	610573	762403	0.94	0.85	1.03
1998	1063525	842819	1342025	385716	345141	431060	592624	0.94	0.85	1.04
1999	632242	500734	798289	280733	252211	312481	484910	0.94	0.85	1.03
2000	750091	596852	942675	255605	233854	279379	414868	0.85	0.77	0.94
2001	593491	472683	745175	383171	346238	424044	426471	0.74	0.67	0.82
2002	374323	298574	469290	520976	470545	576812	535045	0.68	0.61	0.75
2003	757065	605595	946421	571225	517255	630827	551990	0.63	0.57	0.70
2004	242114	196508	298304	665716	603904	733856	606445	0.70	0.63	0.78
2005	693555	561762	856268	579178	526767	636803	641276	0.70	0.63	0.79
2006	536915	434421	663591	584022	531619	641590	537642	0.60	0.54	0.68
2007	1244698	1001618	1546772	651314	588970	720257	486883	0.44	0.39	0.50
2008	1003692	802201	1255793	722403	653337	798771	464171	0.36	0.32	0.41
2009	582481	464438	730526	1011809	916652	1116845	523430	0.31	0.27	0.35
2010	202041	156697	260506	1244121	1125664	1375043	609983	0.29	0.25	0.33
2011	358414	280491	457985	1806496	1627939	2004636	719830	0.31	0.27	0.35
2012	503503	400530	632949	2027169	1819728	2258258	727663	0.29	0.25	0.33
2013	465036	365513	591657	2262649	2022187	2531706	966209	0.31	0.27	0.36
2014	852367	674396	1077303	2160382	1918011	2433380	986449	0.33	0.29	0.38
2015	451911	360009	567273	1759211	1549221	1997663	864384	0.33	0.29	0.38
2016	286123	224783	364202	1416694	1244845	1612266	849422	0.35	0.31	0.40
2017	782183	613967	996487	1439236	1280270	1617940	868276	0.38	0.33	0.44
2018	507397	384793	669066	1298311	1149222	1466741	778627	0.40	0.34	0.46
2019	657814	493945	876048	1238878	1074599	1428273	692609	0.41	0.34	0.49
2020	561552	402771	782928	1013636	843651	1217870	692903	0.44	0.35	0.54
2021	557000*			902035	695846	1169322				

<sup>\*</sup> Recruitment model estimate.

#### **Sources and references**

ICES. 2003. Report of the Study Group on Biological Reference Points for Northeast Arctic Cod, 13–17 January 2003, Svanhovd, Norway. ICES CM 2003/ACFM:11. 39 pp.

ICES. 2017. Report of the Arctic Fisheries Working Group (AFWG), 19–25 April 2017, ICES HQ, Copenhagen, Denmark. ICES CM 2017/ACOM:06. 493 pp.

ICES. 2021a. Benchmark Workshop for Barents Sea and Faroese Stocks (WKBARFAR 2021).

ICES Scientific Reports. 3:21. 205 pp. https://doi.org/10.17895/ices.pub.7920.

ICES. 2021b. Advice on fishing opportunities. *In* Report of the ICES Advisory Committee, 2021. ICES Advice 2021, section 1.1.1. https://doi.org/10.17895/ices.advice.7720.

ICES. 2021c. Arctic Fisheries Working Group (AFWG). ICES Scientific Reports. 3:58. https://doi.org/10.17895/ices.pub.8196.

Download the stock assessment data and figures.

Recommended citation: ICES. 2021. Cod (*Gadus morhua*) in subareas 1 and 2 (Northeast Arctic). *In* Report of the ICES Advisory Committee, 2021. ICES Advice 2021, cod.27.1-2, <a href="https://doi.org/10.17895/ices.advice.7741">https://doi.org/10.17895/ices.advice.7741</a>.