

Exercise 1b: Differential Kinematics of the ABB IRB 120

Prof. Marco Hutter*

Teaching Assistants:

Jean-Pierre Sleiman, Maria Vittoria Minniti, Shao Zhang,
Jan Preisig and Mayank Mittal

October 6, 2021

Abstract

The aim of this exercise is to calculate the differential kinematics of an ABB robot arm. You will practice on the derivation of velocities for a multi-body system, as well as derive the mapping between generalized velocities and end-effector velocities. A separate MATLAB script will be provided for the 3D visualization of the robot arm.



Figure 1: The ABB IRW 120 robot arm.

1 Introduction

The following exercise is based on an ABB IRB 120 depicted in figure 2. It is a 6-link robotic manipulator with a fixed base. During the exercise you will implement

*original contributors include Michael Blösch, Dario Bellicoso, and Samuel Bachmann




Figure 2: ABB IRB 120 with coordinate systems and joints.

several different MATLAB functions, which you should test carefully since the next exercises will depend on them. To help you with this, we have provided the script prototypes (download from Moodle).

Throughout this document, we will employ I for denoting the inertial world coordinate system (which has the same pose as the coordinate system $P0$ in figure 2) and E for the coordinate system attached to the end-effector (which has the same pose as the coordinate system $P6$ in figure 2).

2 Differential Kinematics

Exercise 2.1

In this exercise, we seek to compute an analytical expression for the twist ${}^I\mathbf{w}_E = [{}^I\mathbf{v}_E^T \quad {}^I\boldsymbol{\omega}_E^T]^T$ of the end-effector. To this end, find the analytical expression of the end-effector linear velocity vector ${}^I\mathbf{v}_E$ and angular velocity vector ${}^I\boldsymbol{\omega}_E$ as a function of the linear and angular velocities of the coordinate frames attached to each link.

Hint: start by writing the rigid body motion theorem and extend it to the case of a 6DoF arm.

Exercise 2.2

This exercise focuses on deriving the mapping between the generalized velocities $\dot{\mathbf{q}}$ and the end-effector twist ${}^I\mathbf{w}_E$, namely the *basic* or *geometric* Jacobian ${}^I\mathbf{J}_{e0} = [{}^I\mathbf{J}_P^T \quad {}^I\mathbf{J}_R^T]^T$. To this end, you should derive the translational and rotational Jacobians of the end-effector, respectively ${}^I\mathbf{J}_P$ and ${}^I\mathbf{J}_R$. To do this, you can start from the derivation you found in exercise 1. The Jacobians should depend on the minimal coordinates \mathbf{q} only. Remember that Jacobians map joint space generalized

velocities to operational space generalized velocities:

$${}^I\mathbf{v}_{IE} = {}^I\mathbf{J}_P(\mathbf{q})\dot{\mathbf{q}} \quad (1)$$

$${}^I\boldsymbol{\omega}_{IE} = {}^I\mathbf{J}_R(\mathbf{q})\dot{\mathbf{q}} \quad (2)$$

Please implement the following two functions:

```
1 function J_P = jointToPosJac(q)
2 % Input: vector of generalized coordinates (joint angles)
3 % Output: Jacobian of the end-effector translation which maps joint
4 % velocities to end-effector linear velocities in I frame.
5
6 % Compute the translational jacobian.
7 J_P = zeros(3, 6);
8 end
9
10 function J_R = jointToRotJac(q)
11 % Input: vector of generalized coordinates (joint angles)
12 % Output: Jacobian of the end-effector orientation which maps joint
13 % velocities to end-effector angular velocities in I frame.
14
15 % Compute the rotational jacobian.
16 J_R = zeros(3, 6);
17 end
```