

0.0 TOPIC

$x + y$ some example equation related to the topic (1)

Mynd? Always try to have a visual representation of the equation

Topic: describes. minute details about the equation

Theorem: *Definition of "Topic"* theorem + tab

Uses: What is it be used for in linear algebra

Example: "We can use **this** to calculate **that**", followed by an example

Mynd? Always try to have a visual along with the example

Exercise: Exercise for the reader

Revision

0.1 LINEAR EQUATIONS

$$x + 2 = y \quad (2)$$


This is an example of a **linear equation**, one that has two variables, x and y , and it describes how the value of one of the variables depends on the value of the other variable.

What makes it **linear** is that every variable is only raised to the first power, so *this*

$$x^2 + 2 = y$$

is **not** a linear equation.

Lets plot (draw) equation (2), for a few diferent values of x , say, -6 , -2 , 1 and 5 .



As you can see, the graph of this equation is a **straight line**, which is true for all linear equations

0.1 LINEAR EQUATION WITH MULTIPLE VARIABLES

$$x + 9y + z = 3 \quad (3)$$

Lets now extend our definition of a linear equation to include more variables. (2) had only x and y , but (3) has 3 variables, x , y and z .

The 9 in front of the y ? That is called a *coefficient*, and the 3 on the right-hand-side is called a *constant*

Because we will (eventually) run out of letters in the alphabet, we write our equations like this:

$$c_1x_1 + c_2x_2 + c_3x_3 + \dots + c_nx_n = b$$

Here the c 's are the *coefficients*, the x 's are the *variables* and the b is (spoiler) the *constant*.

It may look complex the first time, but you'll get used to reading equations like this.

Lets look at an example: Jimmy goes to the store to buy cokes, snickers and apples. Jimmy knows that a can of coke is 2.2\$, snickers is 1.6\$ and an apple is 3.0\$

If c_1 is price of coke, c_2 is price of snickers and c_3 is price of apples, our equation would look like this

$$2.2x_1 + 1.6x_2 + 3.0x_3 = b$$

Jimmy needs a couple of cokes (x_1) and four apples (x_3). Jimmy has 20\$.

How many snickers bars can he buy with the leftover money?

$$2.2 \cdot 2 + 1.6x_2 + 3.0 \cdot 4 = 20$$

Do the math and help Jimmy get his snickers by solving for x_2 .

0.1 SYSTEM OF EQUATIONS

$$2x_1 + x_2 = 17$$

$$x_1 + 3x_2 = 26$$

The equations above are an example of a **system of equations**.

We want to solve these systems by finding the correct values for the *variables*, in this case, x_1 and x_2 , so that both equations work out.

We could start by *guessing* that $x_1 = 3$ and $x_2 = 5$, which would give us

$$2 \cdot 3 + 2 = 8 \neq 17$$

$$3 + 3 \cdot 5 = 18 \neq 26$$

This is far from correct, we need the **the substitution method**.

The substitution method consists of **two** steps, that you use over and over again until the system has been solved. These steps are:

1. Isolating a variable

2. Substitution

3. Simplification

Lets take another look at our system

$$2x_1 + x_2 = 17 \tag{4}$$

$$x_1 + 3x_2 = 26 \tag{5}$$

and solve it using the substitution method.

1. isolate the x_1 from (4), $x_1 = 26 - 3x_2$
2. substitute x_1 into (3), $2(26 - 3x_2) + x_2 = 17$
3. now the system looks like

$$2(26 - 3x_2) + x_2 = 17 \quad (6)$$

$$x_1 + 3x_2 = 26 \quad (7)$$

4. which simplifies to

$$x_2 = 7 \quad (8)$$

$$x_1 + 3x_2 = 26 \quad (9)$$

5. now we just insert x_2 into (8) to get

$$x_2 = 7 \quad (10)$$

$$x_1 + 21 = 26 \quad (11)$$

6. so (8) simplifies to $x_1 = 5$

Now its your turn to solve the following


$$6x_1 + 2x_2 = 70$$

$$3x_1 + 3x_2 = 45$$

Vectors

0.2 VECTOR PROPERTIES

$$\vec{v} = [4 \ 3] = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$



Vectors: Its easy to think of vectors as a **length** and a **direction**, usually denoted $v = [x_1 \ x_2 \ x_3 \ \dots \ x_n]$, where the **x's** are usually called **elements**.

Above we have an example of a **(2-dimensional)** vector, written both as a **row vector** and a **column vector**. The vector represents a "travel" by 4 steps along the x_1 axis and 3 steps along the x_2 axis. So if you find yourself positioned at the point $(2, -4)$ and someone "applies" this vector to you, you'll be moved to $(2 + 4, -4 + 3) = (6, -1)$.

Properties: The **length** (also known as *norm* or *size*) of a vector is written as $|\vec{v}|$ and found with the Pythagorean-theorem:

$$|\vec{v}| = \sqrt{4^2 + 3^2} = \sqrt{25} = 5$$

For higher dimensional vectors, the calculations look similar

$$|\mathbf{u}| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$$

A two-dimensional vector also has a **direction** written as $\theta_{\vec{v}}$ and calculated using absolute classic geometry

$$\theta_{\vec{v}} = \tan^{-1} \frac{3}{4}$$

Exercise: Find the length of the 5-dimensional vector $\vec{p} = [6 \ 2 \ 3 \ 9 \ 1]$

0.2 ELEMENTARY VECTOR OPERATIONS


$3 \cdot [1 \ 2]$	scalar multiplication
$[1 \ 2]^T$	transpose
$[2 \ 1] + [1 \ 2]$	addition
$[2 \ 1] - [1 \ 2]$	subtraction

Operations: These 4 operations are the most elementary and common operations you will be using in linear algebra. They are

- **Scalar multiplication** simply multiply every element of the vector with the scalar.

$$3 \cdot [1 \ 2] = [3 \ 6]$$

All this operation does is
lengthen the vector.




- **Transposing** a vector is simply converting it from a row-vector to a column-vector, or a column-vector back to a row-vector.

$$[1 \ 2]^T = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

- Adding two vectors is straightforward

$$\begin{bmatrix} 2 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 3 \end{bmatrix}$$


Visually, when adding two vectors you simply add one to the end of the other. Note that it does not matter in which order you add them.



- Subtraction, like addition, is intuitive

$$\begin{bmatrix} 2 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & -1 \end{bmatrix}$$

Here we negate the vector we are subtracting (the red one in this case) and just like addition we add it to the end of the other.



Exercise: Here are 3 vectors, $u = \begin{bmatrix} 3 & 5 \end{bmatrix}$, $v = \begin{bmatrix} 8 & 10 \end{bmatrix}$, $p = \begin{bmatrix} 10 & 1 & 1 \end{bmatrix}$, can you add them together?

0.2 VECTOR DOT PRODUCT

$$[1 \ 3] \bullet [4 \ 1] = 1 \cdot 4 + 3 \cdot 1 = 7$$

Dot product: When taking a dot product of two vectors we multiply their corresponding elements together and then add up the products. The result is a single number.

Definition: Say we have two vectors $u = [u_1 \ u_2 \ \dots \ u_n]$ and $v = [v_1 \ v_2 \ \dots \ v_n]$, their dot product is defined as

$$\vec{u} \bullet \vec{v} = u_1 \cdot v_1 + u_2 \cdot v_2 + \dots u_n \cdot v_n$$

Uses: The dot product tells us a lot about the direction of the vectors relative to each other:

- A positive dot product: The angle between the vectors is less than 90° .
- The dot product is 0: The vectors are perpendicular to each other.
- A negative dot product: The angle between the vectors is more than 90° .

Example: "We can use **this** to calculate **that**", followed by an example

Mynd? Always try to have a visual along with the example

Exercise: Exercise for the reader