

Satellite Imagery


assisting in


Crop yield Prediction

Observing Satellite Images with human eye

Following are satellite images of **2** places on the planet,

The Amazon Rainforest, and
The Sahara Desert





We (humans) can easily tell by **looking** at both the images, which place is which.

The **greener** the place is, more likely it is for a crop to grow and sustain.

“greenness measurement”

There are various indices used to calculate various metrics from the satellite image data.

One such index is used to determine the **greenness**.

This index is called **Normalized Difference Vegetation Index**.

Calculation of NDVI

$$\frac{\left(\begin{array}{|c|} \hline \text{Near IR} \\ \hline \end{array} \text{ minus } \begin{array}{|c|} \hline \text{Red} \\ \hline \text{Green} \\ \hline \text{Blue} \\ \hline \end{array} \right)}{\left(\begin{array}{|c|} \hline \text{Near IR} \\ \hline \end{array} \text{ plus } \begin{array}{|c|} \hline \text{Red} \\ \hline \text{Green} \\ \hline \text{Blue} \\ \hline \end{array} \right)}$$


Name	Units	Scale	Wavelength	Description
B1		0.0001	0.435-0.451 μm	Band 1 (ultra blue) surface reflectance
B2		0.0001	0.452-0.512 μm	Band 2 (blue) surface reflectance
B3		0.0001	0.533-0.590 μm	Band 3 (green) surface reflectance
B4		0.0001	0.636-0.673 μm	Band 4 (red) surface reflectance
B5		0.0001	0.851-0.879 μm	Band 5 (near infrared) surface reflectance
B6		0.0001	1.566-1.651 μm	Band 6 (shortwave infrared 1) surface reflectance
B7		0.0001	2.107-2.294 μm	Band 7 (shortwave infrared 2) surface reflectance

Value of NDVI can range from -1 to +1

+1 being very green

-1 being not green (red)

COLOR WHEEL



- ★ Therefore, we collect satellite image data of the input coordinates.
- ★ We determine/measure the “greenness” of the area.
- ★ We provide this **greenness measurement (NDVI)** to the ML model.
- ★ ML model does its job and gives the final crop that can be grown