

# ESO201A : THERMODYNAMICS

## 2021-22 1st semester

### IIT Kanpur

Instructor : P.A.Apte

## Lecture 4

**Copyright:** The instructor of this course (ESO201A) owns the copyright of all the course materials. The lecture material was distributed only to the students attending the course ESO201A of IIT Kanpur, and should not be distributed in print or through electronic media without the consent of the instructor. Students can make their own copies of the course materials for their use.

# Changes in internal energy:

As engineers, we are concerned with processes involving changes of temperature, pressure, phase, and composition.



**Steam boiler**

Changes of phase from water to steam

Changes in composition due to burning of fuel (usually coal)



**Air compressor**

Changes in Temperature and Pressure

# Changes in internal energy:

The energy transfer in the form of heat and work in all such processes results in changes of internal energy  $U$ . Hence it is important to have a better understanding of changes in  $U$ .

At a microscopic level, changes in  $U$  are due to changes of average kinetic and potential energies of constituent atoms and molecules




$$\Delta U = \Delta \langle \text{k.e.} \rangle + \Delta \langle \text{p.e.} \rangle$$

$\Delta \langle \text{k.e.} \rangle$  = Change in average kinetic energy of atoms and molecules

$\Delta \langle \text{p.e.} \rangle$  = Change in average potential energy of atoms and molecules

$$\Delta \langle \text{k.e.} \rangle$$

(translational k.e. + rotational k.e. + vibrational k.e.)




Processes involving changes of temperature are affected by changes in kinetic energies due to translation, rotation, and (intra-molecular) vibration of molecules

The amount of heat per kg or per mole of a substance required to increase the temperature by  $1^{\circ}\text{C}$  is known as specific heat .

So, these microscopic kinetic energies directly affect specific heat.

Specific heat at constant volume ( $C_v$ ) refer to amount of heat absorbed per unit mass or per mole to increase temperature by  $1^{\circ}\text{C}$  at constant volume. Similarly specific heat at constant pressure ( $C_p$ ) is defined.

Hydrogen gas is composed of diatomic molecules of  $H_2$ . The following plot shows how specific heat at constant volume ( $C_v$ ) is affected by kinetic energies due to translation, rotation, and vibration of  $H_2$  molecules



$$\Delta\langle\text{p.e.}\rangle$$

(Intermolecular p.e. + Intramolecular p.e.)

Intermolecular p. e. Due to binding forces between the molecules. These are important in changes in which average distance between the molecules change, e.g., change in T (fluids, solids), P (for gases), and phase (e.g. liquid→vapor, solid→liquid) and changes of composition without a chemical reaction (dissolution). These binding forces include


- (i) Van der Waal's forces : These are the non-ionic forces between the molecules
- (ii) Ionic forces : due to charge-charge interactions. For example, common salt (NaCl) has a large melting point (801 °C) due to strong ionic forces between Na<sup>+</sup> and Cl<sup>-</sup> ions.

Intermolecular p. e. directly affect physical properties. For example :

$\text{CH}_4$  (Methane) is a gas at ambient conditions because intermolecular forces are weakly attractive due to non-ionic nature of interactions

In  $\text{CH}_3\text{-OH}$  (Methanol), due to strong electronegativity of oxygen in  $\text{-OH}$  group, it attracts electron cloud towards itself. As a result, H atom of  $\text{-OH}$  group develops a slightly positive charge and O atom of  $\text{-OH}$  group develops a slightly negative charge. This leads to strong hydrogen bonded (ionic) interactions between different molecules. As a result of these strong interactions, methanol exists in a liquid state at ambient conditions.

Intermolecular p. e. These binding forces are important during heating/cooling of a material, compression/expansion of a gas, phase changes, dissolution, and mixing processes



Dissolution of sugar in water



Intramolecular p. e. These binding forces are important during changes of composition (e.g., reactions)

## 1. Combustion



## 2. Reforming of methane (used in Ammonia production)

**Table 1:** Basic processes for syngas production from methane.

Process	Reaction	H <sub>2</sub> /CO ratio	$\Delta H_{1173\text{K}}$ [kJ mol <sup>-1</sup> ]
Steam reforming	$\text{CH}_4 + \text{H}_2\text{O} \rightarrow \text{CO} + 3\text{H}_2$	3:1	225.7
CO <sub>2</sub> reforming (dry reforming)	$\text{CH}_4 + \text{CO}_2 \rightarrow 2\text{CO} + 2\text{H}_2$	1:1	258.8
Oxy reforming (partial oxidation)	$\text{CH}_4 + 1/2\text{O}_2 \rightarrow \text{CO} + 2\text{H}_2$	2:1	-23.1

# Analysis of processes involving **pure** substance:

If a substance (which has a fixed chemical composition throughout) does not undergo a change in composition, it can be considered as a pure substance


A pure substance need not contain a single element or a single compound.




Air contains  $\text{N}_2$ ,  $\text{O}_2$ , and small amounts of other gases

In processes NOT involving change of composition, air is treated as a pure substance

# Analysis of processes involving **pure** substance:




(water + steam) is a pure substance




(water + air) is NOT a pure substance

## A pure substance can exist in different phases


Ref. : Cengel and Boles , 8th Edition (2015)



SOLID



LIQUID



GAS

# State postulate :

## (for a single phase of a pure substance)

It is known experimentally that whenever temperature and pressure of a single phase of a pure substance (e.g., liquid water or air) are specified, all other intensive properties of water have unique values.

For example, water at  $T = 50^{\circ}\text{C}$  and  $P = 1 \text{ atm}$  has a density of  $988 \text{ kg/m}^3$  and a viscosity of  $0.5474 \times 10^{-3} \text{ Ns/m}^2$ .

Thus,  $\theta = f(T, P)$  where  $\theta$  = density, viscosity, refractive index, specific heat or any other intensive property [Recall that mathematically if  $z = f(x, y)$ , then  $z$  has a unique value for given values of  $x$  and  $y$ ]

Similarly, temperature ( $T$ ) and specific volume ( $v$ ) of water can determine all other intensive properties uniquely, i.e.,  $\theta = f(T, v)$ .

The above experimental observations are formally stated as a **state postulate** :

**In processes not involving electrical, magnetic, stress-strain or surface tension effects, the state of a given phase of a pure substance is uniquely determined by specifying two independent, intensive properties.**

Throughout this course, we will consider that  
THERE ARE NO ELECTRIC, MAGNETIC, STRESS-  
STRAIN, AND SURFACE TENSION EFFECTS,  
unless stated explicitly.


# Expansion of a pure substance :

In an expansion process of a pure substance NOT involving changes in kinetic or potential energies, first law of thermodynamics states that :

$$U_2 - U_1 = Q_{\text{in}} - W_{\text{out}}$$

where,

$$W_{\text{out}} = \int_1^2 P_{\text{ext}} dV$$



For an expansion process,  $P > P_{\text{ext}}$   
Therefore, one can write  
 $(P - \Delta P) = P_{\text{ext}}$

# Expansion of a pure substance :

$$Q_{in} = (U_2 - U_1) + \int_1^2 P dV - \int_1^2 (\Delta P) dV$$

For a sufficiently slow expansion process,  $\Delta P \rightarrow 0$ . Hence third term on the right hand side of above equation can be neglected in comparison with the second term. So, for a slow expansion,

$$Q_{in} = (U_2 - U_1) + \int_1^2 P dV$$

If the expansion is sufficiently slow and pressure is constant throughout the process, the above equation reduces to

$$Q_{in} = (U_2 - U_1) + P(V_2 - V_1)$$


$$Q_{in} = H_2 - H_1$$

Here,  $H = (U + PV)$  is the **Enthalpy**. Enthalpy is a state property

# Constant pressure expansion of a pure substance :


The above derivation shows that the heat absorbed by system in a sufficiently slow expansion at constant pressure is equal to increase of enthalpy of the substance.

## Constant pressure expansion of water to steam at $P = 1 \text{ atm}$ ( $101.325 \text{ N/m}^2$ )





# Constant pressure expansion of water at 1 atm :



Note that based on Joule's experiments, these changes of state of water from 1 to 5 can also be brought about by performing mechanical work by enclosing water in adiabatic container.

# Constant pressure expansion of water at 1 atm :

According to the derivation in previous slides, for a constant pressure expansion process from 1→2 (see diagram on last slide)

$$\begin{aligned} Q_{\text{in}} (1 \rightarrow 2) &= H_2 - H_1 \\ &= m (h_2 - h_1) \end{aligned}$$

$h$  = specific enthalpy of liquid water (kJ/kg)

$m$  = mass of water (kg)

The quantity ( $H_2 - H_1$ ) for a change of state from compressed liquid(1) to saturated liquid (2) is called as “sensible” heat since temperature increases as a result of providing heat, i.e., water becomes hotter which can be sensed by hand.

Similarly, the quantity ( $H_5 - H_4$ ) for a change of state from saturated vapor (4) to superheated vapor (5) is also called as “sensible heat”

# Constant pressure expansion of water at 1 atm :

For the process from 2→4 (see previous slides)

$$\begin{aligned} Q_{\text{in}} (2 \rightarrow 4) &= H_4 - H_2 \\ &= m (h_4 - h_2) \\ &= m (h_{g@1\text{atm}} - h_{f@1\text{atm}}) \end{aligned}$$

$h_g$  = specific enthalpy of saturated vapor (kJ/kg)

$h_f$  = specific enthalpy of saturated liquid (kJ/kg)

The quantity ( $H_4 - H_2$ ) for a change of state from saturated liquid(2) to saturated vapor (2) is called as “latent” heat since temperature does NOT change in this process (“latent” means hidden).

The quantity  $h_{fg} = (h_g - h_f)$  is the latent heat of vaporization in kJ/kg

The values of  $h_f$ ,  $h_g$ ,  $h_{fg}$  are listed in tables known as “steam tables”.

# Constant pressure expansion of water :

Note that expansion process of water by heating may be carried out at different (constant) pressures. At 1 atm,  $T_{\text{sat}} = 100^\circ\text{C}$ . This is the saturation temperature, i.e., the temperature at which liquid  $\rightarrow$  vapor transition occurs at constant pressure

It is observed that  $T_{\text{sat}}$  is uniquely determined by the pressure.

Thus,  $T_{\text{sat}} = f(P)$ .

Press., $P$ kPa	Sat. temp., $T_{\text{sat}}$ $^\circ\text{C}$
101.325	99.97
125	105.97
150	111.35
175	116.04
200	120.21
225	123.97
250	127.41
275	130.58
300	133.52