

ESO 208A: Computational Methods in Engineering

Richa Ojha

Department of Civil Engineering
IIT Kanpur



Acknowledgements: Profs. Abhas Singh and Shivam Tripathi (CE)



- **Copyright:**

The instructor of this course owns the copyright of all the course materials. This lecture material was distributed only to the students attending the course *ESO208A: Computational methods in Engineering of IIT Kanpur* and should not be distributed in print or through electronic media without the consent of the instructor. Students can make their own copies of the course materials for their use.



Recap

- Pitfalls of Gauss Elimination Method
- Gauss Jordan Method
- LU decomposition-Gauss Elimination, Dolittle, Crout

Today's lecture

- Thomas Algorithm
- Cholesky Decomposition
- Forward Error Analysis
- Indirect Methods-Gauss-Seidal, Jacobi iterative method



LU decomposition

Thomas Algorithm (Tri-diagonal Matrix)

Thomas Algorithm - Tri diagonal

$$\begin{bmatrix} & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \end{bmatrix}$$

$$\begin{matrix} \rightarrow \\ \rightarrow \\ \rightarrow \end{matrix} \begin{bmatrix} d_1 & u_1 & & & \\ l_2 & d_2 & u_2 & & \\ & l_3 & d_3 & u_3 & \\ & & & \ddots & \\ & & & & l_n & d_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

Define

$$\alpha_1 = d_1$$

$$\beta_1 = b_1$$

$$\lambda_2 = \frac{l_2}{\alpha_1} \quad \begin{matrix} R_1 - \alpha_1 x_1 & + & u_1 x_2 & = & \beta_1 \\ R_2 - l_2 x_1 & + & d_2 x_2 + u_2 x_3 & = & b_2 \end{matrix}$$

$$R_2 \rightarrow R_2 - \lambda_2 R_1$$

$$x_2 (d_2 - \lambda_2 u_1) + u_2 x_3 = b_2 - \lambda_2 \beta_1$$

Define

$$\alpha_2 = d_2 - \lambda_2 u_1$$

$$\beta_2 = b_2 - \lambda_2 \beta_1$$

$$\Rightarrow \begin{matrix} \alpha_2 x_2 & + & u_2 x_3 & = & \beta_2 \\ l_3 x_2 & + & d_2 x_3 + u_3 x_4 & = & b_3 \end{matrix}$$

$$\lambda_3 = \frac{l_3}{\alpha_2}$$

$$\lambda_i^o = \frac{l_i^o}{\alpha_{i-1}^o}$$

$$\alpha_i^o = d_i - \lambda_i^o u_{i-1}$$

$$\beta_i^o = b_i - \lambda_i^o \beta_{i-1}$$

$$x_i = \frac{\beta_i - u_i x_{i+1}}{\alpha_i} \quad \leftarrow \alpha_i x_i + u_i x_{i+1} = \beta_i$$

$$i=n$$

$$\alpha_n x_n = \beta_n \Rightarrow x_n = \beta_n / \alpha_n$$

LU decomposition

Thomas Algorithm (Tri-diagonal Matrix)

GE : $O(2/3n^3)$
 Thomas : $O(n)$

Example

$$\begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \\ 8 \end{bmatrix}$$

$$\alpha_1 = d_1 \quad \beta_1 = b_1 \quad \lambda_i = \frac{l_i}{\alpha_{i-1}}$$

$$\alpha_i = d_i - \lambda_i u_{i-1}$$

$$\beta_i = b_i - \lambda_i \beta_{i-1}$$

$$x_n = \beta_n / \alpha_n$$

$$x_i = \frac{\beta_i - u_i x_{i+1}}{\alpha_i}$$

i	l	d	u	b	λ	α	β	x
1		2	1	4		2	4	$x_1 = 1$
2	1	2	1	8	$\frac{l_2}{\alpha_1} = 0.5$	$2 - 0.5 \times 1 = 1.5$	$8 - 0.5 \times 4 = 6$	$x_2 = \frac{6 - 1 \times 3}{3/2} = 2$
3	1	2		8	$\frac{l_3}{\alpha_2} = \frac{1}{1.5} = 2/3$	$4/3$	4	$x_3 = 4/4/3 = 3$

LU decomposition

Cholesky Decomposition (for +ve definite matrix)

Diagonalization (*LDU* theorem):

Let A be a $n \times n$ invertible matrix then there exists a decomposition of the form $A = LDU$ where, L is a $n \times n$ lower triangular matrix with diagonal elements as 1, U is a $n \times n$ upper triangular matrix with diagonal elements as 1, and D is a $n \times n$ diagonal matrix.

Example of a 3×3 matrix:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$
$$= \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} d_{11} = u_{11} & 0 & 0 \\ 0 & d_{22} = u_{22} & 0 \\ 0 & 0 & d_{33} = u_{33} \end{bmatrix} \begin{bmatrix} 1 & u_{12}/u_{11} & u_{13}/u_{11} \\ 0 & 1 & u_{23}/u_{22} \\ 0 & 0 & 1 \end{bmatrix}$$

LU decomposition

Cholesky Decomposition (for +ve definite matrix)

For a symmetric Matrix

$$A^T = A$$

$$A = LDU$$

$$A = A^T = U^T D L^T$$

This implies,

$$L = U^T$$

$$U = L^T$$

For symmetric matrix: $U = L^T$ and $A = LDL^T$

Note that the entries of the diagonal matrix D are the *pivots*!

LU decomposition

Cholesky Decomposition (for +ve definite matrix)

- For **positive definite** matrices, *pivots* are positive!
- Therefore, a diagonal matrix \mathbf{D} containing the *pivots* can be factorized as: $\mathbf{D} = \mathbf{D}^{1/2} \mathbf{D}^{1/2}$
- Example of a 3×3 matrix

$$\begin{bmatrix} d_{11} & 0 & 0 \\ 0 & d_{22} & 0 \\ 0 & 0 & d_{33} \end{bmatrix} = \begin{bmatrix} \sqrt{d_{11}} & 0 & 0 \\ 0 & \sqrt{d_{22}} & 0 \\ 0 & 0 & \sqrt{d_{33}} \end{bmatrix} \begin{bmatrix} \sqrt{d_{11}} & 0 & 0 \\ 0 & \sqrt{d_{22}} & 0 \\ 0 & 0 & \sqrt{d_{33}} \end{bmatrix}$$

- For **positive definite** matrices: $\mathbf{A} = \mathbf{L} \mathbf{D} \mathbf{L}^T = \mathbf{L} \mathbf{D}^{1/2} \mathbf{D}^{1/2} \mathbf{L}^T$
- However, $\mathbf{D}^{1/2} \mathbf{L}^T = (\mathbf{L} \mathbf{D}^{1/2})^T$. Denote: $\mathbf{L} \mathbf{D}^{1/2} = \mathbf{L}_1$
- Therefore, $\mathbf{A} = \mathbf{L}_1 \mathbf{L}_1^T$. This is also a *LU-Decomposition* where one needs to evaluate only one triangular matrix \mathbf{L}_1 .

LU decomposition

Cholesky Decomposition (for +ve definite matrix)

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{bmatrix} = \begin{bmatrix} l_{11} & & & & \\ l_{21} & l_{22} & & & \\ l_{31} & l_{32} & l_{33} & & \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ l_{n1} & l_{n2} & \dots & & l_{nn} \end{bmatrix} \begin{bmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & \ddots & \\ & & & & 1 \end{bmatrix}$$

$$a_{ij} = q_{ji}$$

$$l_{11}^2 = a_{11}$$

$$l_{11} = \sqrt{a_{11}}$$

$$l_{j1} = \frac{a_{j1}}{l_{11}}$$

$$l_{jj} = \sqrt{a_{jj} - \sum_{k=1}^{j-1} l_{jk}^2} \quad j=2, \dots, n$$

$$l_{kj} = \left(a_{kj} - \sum_{s=1}^{j-1} l_{ks} l_{js} \right) / l_{jj} \quad k=j+1, \dots, n$$

LU decomposition

Cholesky Decomposition (for +ve definite matrix)

Example

$$\begin{bmatrix} 4 & 2 & 14 \\ 2 & 17 & -5 \\ 14 & -5 & 83 \end{bmatrix} \rightarrow \begin{bmatrix} \downarrow & & \\ l_{11} & 0 & 0 \\ \rightarrow l_{21} & l_{22} & 0 \\ \rightarrow l_{31} & l_{32} & l_{33} \end{bmatrix} \rightarrow \begin{bmatrix} \downarrow & \downarrow & \\ l_{11} & l_{21} & l_{31} \\ 0 & l_{22} & l_{32} \\ 0 & 0 & l_{33} \end{bmatrix}$$

$$\Rightarrow l_{11}^2 = a_{11} \Rightarrow \boxed{l_{11} = \sqrt{a_{11}}} = \underline{\underline{2}}$$

$$l_{21} l_{11} = a_{21}$$

$$\Rightarrow l_{21} = \frac{a_{21}}{l_{11}} \Rightarrow l_{21} = \frac{2}{2} = \underline{\underline{1}}$$

$$\boxed{l_{j1} = \frac{a_{j1}}{l_{11}}}$$

$$l_{31} = \frac{14}{2} = \underline{\underline{7}}$$

$$\rightarrow L = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 4 & 0 \\ 7 & -3 & 5 \end{bmatrix}$$

$$l_{21}^2 + l_{22}^2 = a_{22}$$

$$\Rightarrow l_{22} = \sqrt{a_{22} - l_{21}^2} = \sqrt{17 - 1} = \underline{\underline{4}}$$

$$\boxed{l_{ij} = \sqrt{a_{ij} - \sum_{k=1}^{j-1} l_{ik}^2}}$$

$$l_{31} l_{21} + l_{32} l_{22} = a_{32}$$

$$\Rightarrow l_{32} = \frac{a_{32} - l_{31} l_{21}}{l_{22}} = \frac{-5 - 7 \times 1}{4} = \underline{\underline{-3}}$$

$$l_{kj} = \frac{a_{kj} - \sum_{s=1}^{j-1} l_{ks} l_{js}}{l_{jj}} \quad k=j+1, \dots, n$$

$$l_{33} = \sqrt{83 - 7^2 - 3^2} = \underline{\underline{5}}$$

Summary

- Thomas Algorithm
- Cholesky Decomposition

