

ESO 208A: Computational Methods in Engineering

Richa Ojha

Department of Civil Engineering
IIT Kanpur



Acknowledgements: Profs. Abhas Singh and Shivam Tripathi (CE)



- **Copyright:**

The instructor of this course owns the copyright of all the course materials. This lecture material was distributed only to the students attending the course *ESO208A: Computational methods in Engineering of IIT Kanpur* and should not be distributed in print or through electronic media without the consent of the instructor. Students can make their own copies of the course materials for their use.



Non-linear equation

In computer, we have five approaches

- **Graphical method**
- **Bracketing methods:** Bisection, Regula-Falsi
- **Open methods:** Fixed point, Newton-Raphson, Secant
- **Special methods for polynomials:** Muller, Bairstow's
- **Hybrid methods:** Brent's



Open Methods



Distinguishing feature:

- Only one starting value
- Convergence is not always guaranteed
- If algorithm converges, the rate of convergence may be faster



Open Methods

1. Fixed Point



$$f(x) = 0 \\ \Rightarrow x = g(x), \quad x_{i+1} = g(x_i)$$

Fixed Point Method

- **Problem:** $f(x) = 0$, find a root $x = \alpha$ such that $f(\alpha) = 0$
 - **Re-arrange the function:** $f(x) = 0$ to $x = g(x)$
 - **Iteration:** $x_{k+1} = g(x_k)$
 - **Stopping criteria:** $\left| \frac{x_{k+1} - x_k}{x_{k+1}} \right| \leq \varepsilon$

Fixed Point Method

Example

$$f(x) = e^{-x} - x = 0$$

$$S = 0.5671$$

1 - $x = e^{-x}$

2 - $x = -\log(x)$

$$x_0 = 0$$

$$x_1 = e^0 = 1$$

$$x_2 = e^{-1} = 0.3678$$

$$x_3 = e^{-0.3678} = 0.692$$

$$x_4 = e^{-x_3} = 0.5$$

$$x_0 = 0.50$$

$$x_1 = 0.69$$

$$x_2 = \vdots$$



Fixed Point Method

Convergence of fixed point

$$x_{i+1} = g(x_i) \quad (1)$$

$$s = g(s) \quad (2)$$

2 - 1

$$s - x_{i+1} = \frac{g(s) - g(x_i)}{\text{MVT}}$$

$$e_{i+1} = g'(\xi) (s - x_i)$$

$\xi \in (s, x_i)$

$$\Rightarrow e_{i+1} = g'(\xi) e_i$$

$$\Rightarrow \frac{|e_{i+1}|}{|e_i|} = |g'(\xi)|$$

$$\text{as } i \rightarrow \infty \quad \left| \frac{e_{i+1}}{e_i} \right| = |g'(s)| \checkmark$$

If $|g'(s)| < 1$ algorithm converges

Linear convergence

If $|g'(s)| > 1$ - algorithm diverges

$$\boxed{\frac{|e_{i+1}|}{|e_i|^p} = C}$$

$$g'(s) = +ive$$

errors will reduce monotonically

$$g'(s) = -ive$$

errors will oscillate



Fixed Point Method

Convergence of fixed point

Example

(1) $x = e^{-x}$
 $g'(x) = -e^{-x}$

$$|g'(s)| < 1$$

(2) $x = -\log(x)$
 $g'(x) = -\frac{1}{x}$

$$s = 0.567$$

$$|g'(s)| > 1$$


$$s = \underline{0.567}$$




Open Methods

2. Newton Raphson Method

- **Problem:** $f(x) = 0$, find a root $x = \alpha$ such that $f(\alpha) = 0$
- **Principle:** Approximate the function as a straight line having same slope as the original function at the point of iteration.



Newton-Raphson Method



$$f'(x_i) = \frac{-f(x_i)}{x_{i+1} - x_i}$$

$$\Rightarrow \boxed{x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}}$$



Newton-Raphson Method

Example

$$x = \sqrt{a}$$

$$f(x) = x^2 - a = 0$$

$$f'(x) = 2x$$

$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}$$

$$= x_i - \frac{x_i^2 - a}{2x_i}$$

$$= \frac{x_i^2 + a}{2x_i}$$

$$\boxed{x_{i+1} = \frac{1}{2} \left[x_i + \frac{a}{x_i} \right]}$$



Newton-Raphson Method

Convergence

Convergence of NR method
Taylor series

$$f(x_{i+1}) = f(x_i) + (x_{i+1} - x_i) f'(x_i)$$

$$\Rightarrow \boxed{x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)}}$$

Let's assume that at i^{th} step
we are just one step away from
the true solution

$$f(s) = f(x_i) + (s - x_i) f'(x_i) + \frac{1}{2} (s - x_i)^2 f''(\xi)$$

$\xi \in (x_i, s)$



Newton-Raphson Method

Convergence

We know s is the solution

$$f(s) = 0$$

$$\Rightarrow 0 = f(x_i) + (s-x_i)f'(x_i) + \frac{1}{2}(s-x_i)^2 f''(\xi)$$

Divide $f'(x_i)$

$$\frac{-f(x_i)}{f'(x_i)} = (s-x_i) + \frac{1}{2}(s-x_i)^2 \frac{f''(\xi)}{f'(x_i)}$$

$$x_{i+1} - x_i - s + x_i = \frac{1}{2}(s-x_i)^2 \frac{f''(\xi)}{f'(x_i)}$$

$$e_{i+1} = -\frac{1}{2} e_i^2 \frac{f''(\xi)}{f'(x_i)}$$

$$\Rightarrow \frac{|e_{i+1}|}{|e_i|^2} = \left| \frac{1}{2} \frac{f''(\xi)}{f'(x_i)} \right|$$

$i \rightarrow \infty$

$$\boxed{\frac{|e_{i+1}|}{|e_i|^2} = \left| \frac{1}{2} \frac{f''(s)}{f'(s)} \right|}$$

- Quadratic convergence




Newton-Raphson Method

Advantages:

Faster convergence (quadratic)

Disadvantages:

Need to calculate derivate



Newton-Raphson
method may get stuck!




Newton-Raphson Method

Places where Newton-Raphson may not work


a) Inflection point

Double Derivative $= 0$; Solution is diverging




Newton-Raphson Method

b) If you have a local minima, it will trap your solution



Newton-Raphson Method

c) Multiple Solutions



- Convergence depends on the function
- Guess is close to the solution


“No substitute for understanding the problem”



Open Methods

3. Secant Method

- **Principle:** Use a difference approximation for the slope or derivative in the Newton-Raphson method. This is equivalent to approximating the tangent with a secant.
- **Problem:** $f(x) = 0$, find a root $x = \alpha$ such that $f(\alpha) = 0$



Secant Method


- **Problem:** $f(x) = 0$, find a root $x = \alpha$ such that $f(\alpha) = 0$
 - **Initialize:** choose two points x_0 and x_1 and evaluate $f(x_0)$ and $f(x_1)$
 - **Approximation:** $f'(x_k) \approx \frac{f(x_k) - f(x_{k-1})}{x_k - x_{k-1}}$, replace in Newton-Raphson
 - **Iteration Formula:** $x_{k+1} = x_k - f(x_k) \frac{x_k - x_{k-1}}{f(x_k) - f(x_{k-1})}$
 - **Stopping criteria:** $\left| \frac{x_{k+1} - x_k}{x_{k+1}} \right| \leq \varepsilon$



Secant Method

Advantages:

- Fast convergence (slightly less than quadratic)
- Overcomes the disadvantage of having to calculate derivate



Secant method may **also** get stuck!

Look for Modified Secant Method!



Summary

What are fixed-point, Newton-Raphson, and Secant method?

Under what conditions these methods will not work.

What is the convergence rate of these methods?

