

M 361K Homework 3

Ishan Shah

October 20, 2022

5.1

3. Let $a < b < c$. Suppose that f is continuous on $[a, b]$, that g is continuous on $[b, c]$, and that $f(b) = g(b)$. Define h on $[a, c]$ by $h(x) := f(x)$ for $x \in [a, b]$ and $h(x) := g(x)$ for $x \in [b, c]$. Prove that h is continuous on $[a, c]$.

7. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous at c and let $f(c) > 0$. Show that there exists a neighborhood $V_\delta(c)$ of c such that if $x \in V_\delta(c)$, then $f(x) > 0$.

12. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous on \mathbb{R} and that $f(r) = 0$ for every rational number r . Prove that $f(x) = 0$ for all $x \in \mathbb{R}$.

5.2

2. Show that if $f : A \rightarrow \mathbb{R}$ is continuous on $A \subseteq \mathbb{R}$ and if $n \in \mathbb{N}$, then the function f^n defined by $f^n(x) = (f(x))^n$, for $x \in A$, is continuous on A .

3. Give an example of functions f and g that are both discontinuous at a point c in \mathbb{R} such that

(a) The sum $f + g$ is continuous at c .

(b) The product $f \cdot g$ is continuous at c .

7. Give an example of a function $f : [0, 1] \rightarrow \mathbb{R}$ that is discontinuous at every point of $[0, 1]$ but such that $|f|$ is continuous on $[0, 1]$.

8. Let f, g be continuous from \mathbb{R} to \mathbb{R} , and suppose that $f(r) = g(r)$ for all rational numbers r . Is it true that $f(x) = g(x)$ for all $x \in \mathbb{R}$?

6.1

3. Let $I \subseteq \mathbb{R}$ be an interval, let $c \in I$, and let $f : I \rightarrow \mathbb{R}$ and $g : I \rightarrow \mathbb{R}$ be functions that are differentiable at c . Prove the following:

- (a) If $\alpha \in \mathbb{R}$, then the function αf is differentiable at c .
- (b) The function $f + g$ is differentiable at c and $(f + g)'(c) = f'(c) + g'(c)$.

4. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) := x^2$ for x rational, $f(x) := 0$ for x irrational. Show that f is differentiable at $x = 0$, and find $f'(0)$

7. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable at c and that $f(c) = 0$. Show that $g(x) := |f(x)|$ is differentiable at c if and only if $f'(c) = 0$.

9. Prove that if $f : \mathbb{R} \rightarrow \mathbb{R}$ is an even function (that is, $f(-x) = f(x)$ for all $x \in \mathbb{R}$) and has a derivative at every point, then the derivative f' is an odd function (that is, $f'(-x) = -f'(x)$ for all $x \in \mathbb{R}$). Also prove that if $g : \mathbb{R} \rightarrow \mathbb{R}$ is a differentiable odd function, then g' is an even function.

6.2

5. Let $a > b > 0$ and let $n \in \mathbb{N}$ satisfy $n \geq 2$. Prove that $a^{1/n} - b^{1/n} < (a - b)^{1/n}$. [Hint: Show that $f(x) := x^{1/n} - (x - 1)^{1/n}$ is decreasing for $x \geq 1$, and evaluate f at 1 and a/b .]

6. Use the Mean Value Theorem to prove that $|\sin x - \sin y| \leq |x - y|$ for all x, y in \mathbb{R} .

8. Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable in (a, b) . Show that if $\lim_{x \rightarrow a} f'(x) = A$, then $f'(a)$ exists and equals A . [Hint: Use the definition of $f'(a)$ and the Mean Value Theorem.]

10. Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $g(x) := x + 2x^2 \sin(1/x)$ for $x \neq 0$ and $g(0) := 0$. Show that $g'(0) = 1$, but in every neighborhood of 0 the derivative $g'(x)$ takes on both positive and negative values. Thus g is not monotonic in any neighborhood of 0.

13. Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable on I . Show that if f' is positive on I , then f is strictly increasing on I .