


Key features responsible for the proper diagnosis of PTSD

Ishan Khatri
July 20, 2019



References

[1] N. Howard, “Approach towards a natural language analysis for diagnosing mood disorders and comorbid conditions,” in *2013 12th Mexican International Conference on Artificial Intelligence*, pp. 234–243, IEEE, 2013.

[2] A. Bourla, S. Mouchabac, W. El Hage, and F. Ferreri, “e-ptsd: An overview on how new technologies can improve prediction and assessment of posttraumatic stress disorder (ptsd),” *European Journal of Psychotraumatology*, vol. 9, no. sup1, p. 1424448, 2018.

- [3] D. M. Sloan, A. T. Sawyer, S. E. Lowmaster, J. Wernick, and B. P. Marx, “Efficacy of narrative writing as an intervention for ptsd: does the evidence support its use?,” *Journal of contemporary psychotherapy*, vol. 45, no. 4, pp. 215–225, 2015.
- [4] K. T. Green, P. A. Dennis, L. C. Neal, A. L. Hobkirk, T. A. Hicks, L. L. Watkins, J. Hayano, A. Sherwood, P. S. Calhoun, and J. C. Beckham, “Exploring the relationship between posttraumatic stress disorder symptoms and momentary heart rate variability,” *Journal of psychosomatic research*, vol. 82, pp. 31–34, 2016.
- [5] P. A. Dennis, L. Watkins, P. S. Calhoun, A. Oddone, A. Sherwood, M. F. Dennis, M. B. Rissling, and J. C. Beckham, “Posttraumatic stress, heart-rate variability, and the mediating role of behavioral health risks,” *Psychosomatic medicine*, vol. 76, no. 8, p. 629, 2014.
- [6] F. Riganello, A. Candelieri, M. Quintieri, D. Conforti, and G. Dolce, “Heart rate variability: an index of brain processing in vegetative state? an artificial intelligence, data mining study,” *Clinical Neurophysiology*, vol. 121, no. 12, pp. 2024–2034, 2010.
- [7] J. L. Kibler, K. Joshi, and M. Ma, “Hypertension in relation to posttraumatic stress disorder and depression in the us national comorbidity survey,” *Behavioral Medicine*, vol. 34, no. 4, pp. 125–132, 2009.
- [8] M. Butt, E. Espinal, R. L. Aupperle, V. Nikulina, and J. L. Stewart, “The electrical aftermath: Brain signals of posttraumatic stress disorder filtered through a clinical lens,” *Frontiers in Psychiatry*, vol. 10, 2019.
- [9] E. L. van den Broek, F. van der Sluis, and T. Dijkstra, “Telling the story and re-living the past: How speech analysis can reveal emotions in post-traumatic stress disorder (ptsd) patients,” in *Sensing emotions*, pp. 153–180, Springer, 2010.
- [10] K. Ćosić, S. Popović, M. Horvat, D. Kukolja, B. Dropuljić, B. Kovač, and M. Jakovljević, “Computer-aided psychotherapy based on multimodal elicitation, estimation and regulation of emotion,” *Psychiatria Danubina*, vol. 25, no. 3, pp. 0–346, 2013.
- [11] J. L. Armony, V. Corbo, M.-H. Clément, and A. Brunet, “Amygdala response in patients with acute ptsd to masked and unmasked emotional facial expressions,” *American Journal of Psychiatry*, vol. 162, no. 10, pp. 1961–1963, 2005.
- [12] A. S. Garrett, V. Carrion, H. Kletter, A. Karchemskiy, C. F. Weems, and A. Reiss, “Brain activation to facial expressions in youth with ptsd symptoms,” *Depression and anxiety*, vol. 29, no. 5, pp. 449–459, 2012.
- [13] T. Fujiwara, R. Mizuki, T. Miki, and C. Chemtob, “Association between facial expression and ptsd symptoms among young children exposed to the great east japan earthquake: a pilot study,” *Frontiers in psychology*, vol. 6, p. 1534, 2015.
- [14] L. M. Shin, S. L. Rauch, and R. K. Pitman, “Amygdala, medial prefrontal cortex, and hippocampal function in ptsd,” *Annals of the New York Academy of Sciences*, vol. 1071, no. 1, pp. 67–79, 2006.
- [15] A. Powers, N. Fani, L. Murphy, M. Briscione, B. Bradley, E. B. Tone, S. D. Norrholm, and T. Jovanovic, “Attention bias toward threatening faces in women with ptsd: eye tracking correlates by symptom cluster,” *European journal of psychotraumatology*, vol. 10, no. 1, p. 1568133, 2019.
- [16] D. Spring, “Thirty-year study links neuroscience, specific trauma, ptsd, image conversion, and language translation,” *Art Therapy*, vol. 21, no. 4, pp. 200–209, 2004.
- [17] R. C. Kessler, A. Sonnega, E. Bromet, M. Hughes, and C. B. Nelson, “Posttraumatic stress disorder in the national comorbidity survey,” *Archives of general psychiatry*, vol. 52, no. 12, pp. 1048–1060, 1995.
- [18] J. I. Bisson, A. Ehlers, S. Pilling, P. Dix, A. Murphy, J. Johnston, D. Richards, S. Turner, W. Yule, C. Jones, *et al.*, “Post-traumatic stress disorder (ptsd): The management of ptsd in adults and children in primary and secondary care,” 2005.
- [19] A. O. Chan and C. Y. Huak, “Influence of work environment on emotional health in a health care setting,” *Occupational Medicine*, vol. 54, no. 3, pp. 207–212, 2004.
- [20] P. A. Joyce and R. Berger, “Which language does ptsd speak? the westernization of mr. sanchez,” *Journal of Trauma Practice*, vol. 5, no. 4, pp. 53–67, 2007.
- [21] E. De Neubourg and C. de Neubourg, “The impact of malnutrition and post traumatic stress disorder on the performance of working memory in children,” 2012.
- [22] Q. X. Ng, A. Y. S. Soh, W. Loke, N. Venkatanarayanan, D. Y. Lim, and W.-S. Yeo, “Systematic review with meta-analysis: The association between post-traumatic stress disorder and irritable bowel syndrome,” *Journal of gastroenterology and hepatology*, vol. 34, no. 1, pp. 68–73, 2019.