

MEDIVIEW IMAGING & DIAGNOSTICS

Accredited Vascular Ultrasound & Radiology Centre
#24, Residency Road, Bengaluru — 560025 • Tel: 31-48854721 • reports@mediviewdiagnostics.in

COLOR DOPPLER / DUPLEX ULTRASOUND REPORT

Left Lower Limb Arterial Study

Patient Details

Patient Name Ramesh Gupta
Age / Sex 46 Years / Male
Patient ID B-2025-1102

Referral / Exam

Referring Doctor Dr. P. Reddy
Examination Date 13-11-2025

Technique

Duplex evaluation of Left lower limb arteries was performed using B-mode, Color and Spectral Doppler imaging. Peak systolic velocities (PSV) and waveform patterns were analyzed in standard arterial segments — Common Femoral, Superficial Femoral (SFA), Popliteal and Tibial arteries. Measurements obtained at rest; care taken to optimize insonation angle.

Findings — Left Lower Limb

Artery	PSV (cm/s)	Waveform	Remarks
Common Femoral Artery	95.6	Triphasic	Normal PSV and triphasic waveform; no focal acceleration.
Profundus Femoris	82.4	Triphasic	Normal flow pattern; no hemodynamically significant lesion seen.
Proximal Superficial Femoral Artery	90.2	Triphasic	PSV within expected range; waveform triphasic.
Mid SFA	87.0	Triphasic	No focal stenosis or dampening of waveform.
Distal SFA	80.5	Triphasic	Physiologic velocity; triphasic waveform preserved.
Popliteal Artery	89.7	Triphasic	Normal spectral pattern; no turbulence or aliasing.
Peroneal / Posterior Tibial Artery	67.1	Triphasic	Distal tibial arteries show preserved triphasic flow and physiologic PSV.
Anterior Tibial Artery	70.3	Triphasic	Normal amplitude and contour for distal vessel.

Ankle—Brachial Index (ABI)

ABI: 0.88 (Left)

Impression

- Ankle—Brachial Index mildly reduced (ABI 0.88) on the left.
- Triphasic waveforms are preserved throughout the examined arterial tree of the left lower limb with physiologic peak systolic velocities at femoral, popliteal and tibial levels.
- No focal hemodynamically significant arterial stenosis or occlusion identified on Doppler interrogation. Overall flow patterns are consistent with preserved distal perfusion; correlate clinically for symptoms. (Note: ABI alone should be interpreted in clinical context.)

Dr. K. Iyer
Consultant Vascular Radiologist