

Clinical Decision-Support: Comprehensive Genetic & Dermatological Dossier

PHYSICIAN & MEDICAL GENETICS EDITION | Record ID: MM-ISHIT--0B53 | Model: Hybrid-XGB-AdaBoost-VQC-Derm-v2.0

Patient Demographics	Age: 19.0 yrs Biological Sex: Female Detected Phototype: Type II	Tri-Model Consensus Risk	29% (MODERATE RISK)
Dual Boosting + VQC	XGBoost (0.45) + AdaBoost (0.25) + 4-Qubit VQC (0.30)	Epistemic Uncertainty	Confidence: High ($\sigma = \pm 0.1174$)

1. Detailed Molecular & Genetic Deficiencies Analysis

Genetic Pathway / Defect	Status / Score	Classification	Clinical Oncologic Implications
TP53 Tumor Suppressor (17p13.1 DNA Binding)	Score: 0.00 / 1.0	Wild-Type	Loss of apoptotic checkpoint control, impaired p21/CDKN1A induction, elevated genomic destabilization.
BRCA1 / BRCA2 (HR Repair Pathway)	Absent (0.0)	Intact	Homologous recombination double-strand break repair failure; high sensitivity to PARP inhibition & platinum agents.
Tumor Mutational Burden (Somatic Density)	1.2 mut/Mb	Low / Moderate	Elevated somatic neoantigen expression. Biomarker for immune checkpoint inhibitor response (Anti-PD-1/PD-L1).
hs-CRP / Inflammation (Microenvironment)	0.5 mg/L	Normal	Sustained pro-inflammatory cytokines (IL-6 / TNF-alpha) facilitating cutaneous angiogenesis & immune evasion.
Hereditary Pedigree (First-Degree Relatives)	None	Negative	Germline susceptibility inheritance profile. Multi-generational screening indicated.

2. Dual Boosting (XGBoost + AdaBoost) & Parameterized Quantum VQC Ensemble

Inference Engine	Estimated Probability	Weight	Algorithm Architecture & Role
XGBoost (Gradient Boosted Trees)	33%	45%	Depth-6 gradient boosting over combined 17-dimensional cutaneous & genomic vector.
AdaBoost (Adaptive Stumps)	64%	25%	Sequential decision stumps emphasizing borderline cases with ambiguous melanin-genomic interplay.
Quantum VQC (Qiskit Aer / PennyLane)	22%	30%	4-qubit parameterized variational circuit with angle embedding exploring non-linear Hilbert space correlations.
Hybrid Calibrated Consensus	29%	100%	Calibrated consensus with discordance variance boundary (± 0.208). Stratified to HIGH RISK.

3. Model Risk Spectrum Across All Skin Phototypes (Fitzpatrick Types I–VI)

Simulates the patient's exact genetic mutations across all skin pigmentation categories to evaluate cutaneous susceptibility:

Fitzpatrick Phototype	Optical ITA / Melanin	XGBoost	AdaBoost	Hybrid Consensus	Stratification Tier
Type I (Very Light) [PATIENT PHENOTYPE]	ITA: 78.0° Mel: 3.5	40%	35%	38%	Moderate Risk
Type II (Light) [PATIENT PHENOTYPE]	ITA: 52.0° Mel: 14.0	38%	34%	36%	Moderate Risk

Type III (Intermediate)	ITA: 35.0° Mel: 22.0	37%	33%	35%	Moderate Risk
Type IV (Tan / Olive)	ITA: 18.0° Mel: 30.0	35%	31%	34%	Moderate Risk
Type V (Brown)	ITA: -12.0° Mel: 46.0	32%	29%	31%	Moderate Risk
Type VI (Dark / Deeply Pigmented)	ITA: -62.0° Mel: 72.0	27%	25%	27%	Moderate Risk

4. Actionable Clinical Guidance & Triage Protocol

Routine preventive maintenance

MANDATORY MEDICAL DISCLAIMER: AI-generated decision-support output. Not a final medical diagnosis. Final clinical decision remains with a qualified healthcare professional.