

# The Mardešić Conjecture for countably compact spaces

Tetsuya Ishiu

Department of Mathematics  
Miami University

Saturday, December 3rd, 2022

# Space-filling curves

A famous theorem of G. Peano in 1890 says that there exists a continuous surjection from  $[0, 1]$  onto  $[0, 1] \times [0, 1]$ .

It was a groundbreaking result, which challenged the notion of dimensions.

Such curves are called *space-filling curves*. The one used in Peano's proof is called the *Peano curve* while there are many other examples, including the *Hilbert curve*.

# Space-filling curves

A famous theorem of G. Peano in 1890 says that there exists a continuous surjection from  $[0, 1]$  onto  $[0, 1] \times [0, 1]$ .

It was a groundbreaking result, which challenged the notion of dimensions.

Such curves are called *space-filling curves*. The one used in Peano's proof is called the *Peano curve* while there are many other examples, including the *Hilbert curve*.

# Space-filling curves

A famous theorem of G. Peano in 1890 says that there exists a continuous surjection from  $[0, 1]$  onto  $[0, 1] \times [0, 1]$ .

It was a groundbreaking result, which challenged the notion of dimensions.

Such curves are called *space-filling curves*. The one used in Peano's proof is called the *Peano curve* while there are many other examples, including the *Hilbert curve*.

## D. Kurepa's theorem

D. Kurepa asked the following natural question.

### Question

For which linearly order topological spaces (LOTS)  $L$ , does there exist a continuous surjection from  $L$  onto  $L \times L$ ?

In 1952, D. Kurepa showed the following theorem.

### Theorem (D. Kurepa)

*For every nondegenerate connected compact Suslin line  $S$ , there is no continuous surjection from  $S$  onto  $S \times S$ .*

## D. Kurepa's theorem

D. Kurepa asked the following natural question.

### Question

For which linearly order topological spaces (LOTS)  $L$ , does there exist a continuous surjection from  $L$  onto  $L \times L$ ?

In 1952, D. Kurepa showed the following theorem.

### Theorem (D. Kurepa)

*For every nondegenerate connected compact Suslin line  $S$ , there is no continuous surjection from  $S$  onto  $S \times S$ .*

# Treybig's Theorem

In 1964, L. B. Treybig proved the following result.

## Theorem (L. B. Treybig)

*If  $X$  and  $Y$  are infinite Hausdorff spaces and  $X \times Y$  is a continuous image of a compact LOTS, then both  $X$  and  $Y$  are metrizable.*

Since  $X$  and  $Y$  are compact and metrizable, both of them are separable.

## Corollary

*If  $L$  is a compact LOTS that is not metrizable, then there exists no continuous surjection from  $L$  to  $L \times L$ .*

# Treybig's Theorem

In 1964, L. B. Treybig proved the following result.

## Theorem (L. B. Treybig)

*If  $X$  and  $Y$  are infinite Hausdorff spaces and  $X \times Y$  is a continuous image of a compact LOTS, then both  $X$  and  $Y$  are metrizable.*

Since  $X$  and  $Y$  are compact and metrizable, both of them are separable.

## Corollary

*If  $L$  is a compact LOTS that is not metrizable, then there exists no continuous surjection from  $L$  to  $L \times L$ .*



# Treybig's Theorem

In 1964, L. B. Treybig proved the following result.

## Theorem (L. B. Treybig)

*If  $X$  and  $Y$  are infinite Hausdorff spaces and  $X \times Y$  is a continuous image of a compact LOTS, then both  $X$  and  $Y$  are metrizable.*

Since  $X$  and  $Y$  are compact and metrizable, both of them are separable.

## Corollary

*If  $L$  is a compact LOTS that is not metrizable, then there exists no continuous surjection from  $L$  to  $L \times L$ .*

# Treybig's Theorem

In 1964, L. B. Treybig proved the following result.

## Theorem (L. B. Treybig)

*If  $X$  and  $Y$  are infinite Hausdorff spaces and  $X \times Y$  is a continuous image of a compact LOTS, then both  $X$  and  $Y$  are metrizable.*

Since  $X$  and  $Y$  are compact and metrizable, both of them are separable.

## Corollary

*If  $L$  is a compact LOTS that is not metrizable, then there exists no continuous surjection from  $L$  to  $L \times L$ .*

# Čertanov's Theorem

It seems that the researchers focus on continuous images of *compact LOTS*, but in 1976, G. I. Čertanov proved the following theorem, which says that Treybig's theorem holds even when we replace 'compact LOTS' by 'countably compact GO-space'.

## Theorem (G. I. Čertanov)

*If  $X$  and  $Y$  are infinite Hausdorff spaces and  $X \times Y$  is a continuous image of a countably compact GO-space, then both  $X$  and  $Y$  are compact and metrizable.*

It seems that this result was hardly recognized, and the paper was only cited once for another result.

# Čertanov's Theorem

It seems that the researchers focus on continuous images of *compact LOTS*, but in 1976, G. I. Čertanov proved the following theorem, which says that Treybig's theorem holds even when we replace 'compact LOTS' by 'countably compact GO-space'.

## Theorem (G. I. Čertanov)

*If  $X$  and  $Y$  are infinite Hausdorff spaces and  $X \times Y$  is a continuous image of a countably compact GO-space, then both  $X$  and  $Y$  are compact and metrizable.*

It seems that this result was hardly recognized, and the paper was only cited once for another result.

# Čertanov's Theorem

It seems that the researchers focus on continuous images of *compact LOTS*, but in 1976, G. I. Čertanov proved the following theorem, which says that Treybig's theorem holds even when we replace 'compact LOTS' by 'countably compact GO-space'.

## Theorem (G. I. Čertanov)

*If  $X$  and  $Y$  are infinite Hausdorff spaces and  $X \times Y$  is a continuous image of a countably compact GO-space, then both  $X$  and  $Y$  are compact and metrizable.*

It seems that this result was hardly recognized, and the paper was only cited once for another result.

# The Mardešić Conjecture

S. Mardešić proposed the following conjecture in 1970, which was proved by G. Martínez-Cervantes and G. Plebanek in 2019.

Theorem (G. Martínez-Cervantes and G. Plebanek)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a compact LOTS for each  $i < d$ ,  $X$  a compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many metrizable factors  $Z_j$ .*

(This statement is slightly stronger than the one presented in their paper, but it clearly holds from their argument).

# The Mardešić Conjecture

S. Mardešić proposed the following conjecture in 1970, which was proved by G. Martínez-Cervantes and G. Plebanek in 2019.

## Theorem (G. Martínez-Cervantes and G. Plebanek)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a compact LOTS for each  $i < d$ ,  $X$  a compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many metrizable factors  $Z_j$ .*

(This statement is slightly stronger than the one presented in their paper, but it clearly holds from their argument).

# The Mardešić Conjecture

S. Mardešić proposed the following conjecture in 1970, which was proved by G. Martínez-Cervantes and G. Plebanek in 2019.

## Theorem (G. Martínez-Cervantes and G. Plebanek)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a compact LOTS for each  $i < d$ ,  $X$  a compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many metrizable factors  $Z_j$ .*

(This statement is slightly stronger than the one presented in their paper, but it clearly holds from their argument).



# What does it mean?

If  $d = s = 1$ , then it coincides with Treybig's Theorem.

By using Peano's Theorem, it is easy to see that for all positive integers  $d$  and  $s$ , there exists a continuous surjection from  $[0, 1]^d$  onto  $[0, 1]^{d+s}$ . The Mardešić Conjecture implies even this seemingly weaker phenomenon does not happen to a nonmetrizable compact LOTS.

# What does it mean?

If  $d = s = 1$ , then it coincides with Treybig's Theorem. By using Peano's Theorem, it is easy to see that for all positive integers  $d$  and  $s$ , there exists a continuous surjection from  $[0, 1]^d$  onto  $[0, 1]^{d+s}$ . The Mardešić Conjecture implies even this seemingly weaker phenomenon does not happen to a nonmetrizable compact LOTS.

# What does it mean?

If  $d = s = 1$ , then it coincides with Treybig's Theorem. By using Peano's Theorem, it is easy to see that for all positive integers  $d$  and  $s$ , there exists a continuous surjection from  $[0, 1]^d$  onto  $[0, 1]^{d+s}$ . The Mardešić Conjecture implies even this seemingly weaker phenomenon does not happen to a nonmetrizable compact LOTS.

# Countably compact version

Recall that Čertanov's Theorem is the countably compact version of Treybig's Theorem. As the Mardešić Conjecture is a generalization of Treybig's Theorem, we may wonder if we can prove the countably compact version of the Mardešić Conjecture. We solved this problem positively, namely proved the following theorem.

## Theorem (T. Ishiu)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a compact LOTS for each  $i < d$ ,  $X$  a **countably** compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection  $f$  from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many compact and metrizable factors  $Z_j$ .*

# Countably compact version

Recall that Čertanov's Theorem is the countably compact version of Treybig's Theorem. As the Mardešić Conjecture is a generalization of Treybig's Theorem, we may wonder if we can prove the countably compact version of the Mardešić Conjecture. We solved this problem positively, namely proved the following theorem.

## Theorem (T. Ishiu)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a compact LOTS for each  $i < d$ ,  $X$  a **countably** compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection  $f$  from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many compact and metrizable factors  $Z_j$ .*

## Countably compact version

Recall that Čertanov's Theorem is the countably compact version of Treybig's Theorem. As the Mardešić Conjecture is a generalization of Treybig's Theorem, we may wonder if we can prove the countably compact version of the Mardešić Conjecture. We solved this problem positively, namely proved the following theorem.

### Theorem (T. Ishiu)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a compact LOTS for each  $i < d$ ,  $X$  a **countably** compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection  $f$  from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many compact and metrizable factors  $Z_j$ .*

## Countably compact version

Recall that Čertanov's Theorem is the countably compact version of Treybig's Theorem. As the Mardešić Conjecture is a generalization of Treybig's Theorem, we may wonder if we can prove the countably compact version of the Mardešić Conjecture. We solved this problem positively, namely proved the following theorem.

### Theorem (T. Ishiu)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a compact LOTS for each  $i < d$ ,  $X$  a **countably** compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection  $f$  from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many compact and metrizable factors  $Z_j$ .*

# Countably compact GO-spaces

S. Purish proved that Stone-Čech compactification of any countably compact GO-space is a LOTS. By using this result, we can obtain the following corollary.

Corollary (T. Ishiu)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a **countably** compact GO-space for each  $i < d$ ,  $X$  a **countably** compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many compact and metrizable factors  $Z_j$ .*



# Countably compact GO-spaces

S. Purish proved that Stone-Čech compactification of any countably compact GO-space is a LOTS. By using this result, we can obtain the following corollary.

Corollary (T. Ishiu)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a **countably** compact GO-space for each  $i < d$ ,  $X$  a **countably** compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection from  $X$  onto  $\prod_{j < d+s} Z_j$ , then there exist at least  $s + 1$ -many compact and metrizable factors  $Z_j$ .*

# Countably compact GO-spaces

S. Purish proved that Stone-Čech compactification of any countably compact GO-space is a LOTS. By using this result, we can obtain the following corollary.

## Corollary (T. Ishiu)

*Let  $d$  and  $s$  be positive integers. Let  $K_i$  be a **countably** compact GO-space for each  $i < d$ ,  $X$  a **countably** compact subspace of  $\prod_{i < d} K_i$ ,  $Z_j$  an infinite Hausdorff space for each  $j < d + s$ . If there exists a continuous surjection from  $X$  onto  $\prod_{j < d+s} X_j$ , then there exist at least  $s + 1$ -many compact and metrizable factors  $Z_j$ .*

# Beginning

Let me outline the proof of the main theorem.

By the observation of G. Martínez-Cervantez and G. Plebanek,  
we can focus on the case  $s = 1$ .

# Beginning

Let me outline the proof of the main theorem.  
By the observation of G. Martínez-Cervantez and G. Plebanek,  
we can focus on the case  $s = 1$ .

# Main Lemma

The following lemma is the most significant piece of the proof of the main theorem.

## Lemma

*Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem. Suppose that  $Z_d$  is not separable. Then, there exists a compact LOTS  $\tilde{K}_i$  for each  $i < d - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < d-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < d} Z_j$ .*

Namely, if one of  $Z_j$ 's is nonseparable, then there exist  $\tilde{K}_i$ ,  $\tilde{X}$ , and  $\tilde{f}$  that satisfies the assumption of the main theorem for  $d - 1$ . This lemma is proved by the heavy use of countable elementary submodels.

# Main Lemma

The following lemma is the most significant piece of the proof of the main theorem.

## Lemma

*Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem. Suppose that  $Z_d$  is not separable. Then, there exists a compact LOTS  $\tilde{K}_i$  for each  $i < d - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < d-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < d} Z_j$ .*

Namely, if one of  $Z_j$ 's is nonseparable, then there exist  $\tilde{K}_i$ ,  $\tilde{X}$ , and  $\tilde{f}$  that satisfies the assumption of the main theorem for  $d - 1$ . This lemma is proved by the heavy use of countable elementary submodels.

# Main Lemma

The following lemma is the most significant piece of the proof of the main theorem.

## Lemma

*Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem. Suppose that  $Z_d$  is not separable. Then, there exists a compact LOTS  $\tilde{K}_i$  for each  $i < d - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < d-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < d} Z_j$ .*

Namely, if one of  $Z_j$ 's is nonseparable, then there exist  $\tilde{K}_i$ ,  $\tilde{X}$ , and  $\tilde{f}$  that satisfies the assumption of the main theorem for  $d - 1$ . This lemma is proved by the heavy use of countable elementary submodels.

# Main Lemma

The following lemma is the most significant piece of the proof of the main theorem.

## Lemma

*Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem. Suppose that  $Z_d$  is not separable. Then, there exists a compact LOTS  $\tilde{K}_i$  for each  $i < d - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < d-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < d} Z_j$ .*

Namely, if one of  $Z_j$ 's is nonseparable, then there exist  $\tilde{K}_i$ ,  $\tilde{X}$ , and  $\tilde{f}$  that satisfies the assumption of the main theorem for  $d - 1$ . This lemma is proved by the heavy use of countable elementary submodels.



# $I(K, M, p)$

We shall sketch the proof of the main lemma.  
In general, we shall define the following notations.

## Definition

Let  $K$  be a compact LOTS and  $M$  a countable set. Then, for all  $p \in K$ , define

$$\eta(K, M, p) = \sup \{ u \in \text{cl}(K \cap M) \mid u \leq p \}$$

$$\zeta(K, M, p) = \inf \{ u \in \text{cl}(K \cap M) \mid u \geq p \}$$

$$I(K, M, p) = [\eta(K, M, p), \zeta(K, M, p)]$$

These play an essential role in my results about LOTS.

# $I(K, M, p)$

We shall sketch the proof of the main lemma.  
In general, we shall define the following notations.

## Definition

Let  $K$  be a compact LOTS and  $M$  a countable set. Then, for all  $p \in K$ , define

$$\eta(K, M, p) = \sup \{ u \in \text{cl}(K \cap M) \mid u \leq p \}$$

$$\zeta(K, M, p) = \inf \{ u \in \text{cl}(K \cap M) \mid u \geq p \}$$

$$I(K, M, p) = [\eta(K, M, p), \zeta(K, M, p)]$$

These play an essential role in my results about LOTS.

# $I(K, M, p)$

We shall sketch the proof of the main lemma.  
In general, we shall define the following notations.

## Definition

Let  $K$  be a compact LOTS and  $M$  a countable set. Then, for all  $p \in K$ , define

$$\eta(K, M, p) = \sup \{ u \in \text{cl}(K \cap M) \mid u \leq p \}$$

$$\zeta(K, M, p) = \inf \{ u \in \text{cl}(K \cap M) \mid u \geq p \}$$

$$I(K, M, p) = [\eta(K, M, p), \zeta(K, M, p)]$$

These play an essential role in my results about LOTS.

# Set up

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main lemma.

- Let  $f_d$  be the  $d$ -th component function of  $f$ .
- Let  $g : X \rightarrow \prod_{j < d} Z_j$  be defined by  $g(x) = f(x) \upharpoonright d$ .  
(Technically,  $g$  must be a slight extension of this, but never mind.)
- Let  $M$  be a good countable elementary submodel of  $H(\theta)$  for a sufficiently large regular cardinal  $\theta$ .

# Set up

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main lemma.

- Let  $f_d$  be the  $d$ -th component function of  $f$ .
- Let  $g : X \rightarrow \prod_{j < d} Z_j$  be defined by  $g(x) = f(x) \restriction d$ .  
(Technically,  $g$  must be a slight extension of this, but never mind.)
- Let  $M$  be a good countable elementary submodel of  $H(\theta)$  for a sufficiently large regular cardinal  $\theta$ .

# Set up

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main lemma.

- Let  $f_d$  be the  $d$ -th component function of  $f$ .
- Let  $g : X \rightarrow \prod_{j < d} Z_j$  be defined by  $g(x) = f(x) \restriction d$ .  
(Technically,  $g$  must be a slight extension of this, but never mind.)
- Let  $M$  be a good countable elementary submodel of  $H(\theta)$  for a sufficiently large regular cardinal  $\theta$ .

# Set up

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main lemma.

- Let  $f_d$  be the  $d$ -th component function of  $f$ .
- Let  $g : X \rightarrow \prod_{j < d} Z_j$  be defined by  $g(x) = f(x) \upharpoonright d$ .  
(Technically,  $g$  must be a slight extension of this, but never mind.)
- Let  $M$  be a good countable elementary submodel of  $H(\theta)$  for a sufficiently large regular cardinal  $\theta$ .

# Set up

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main lemma.

- Let  $f_d$  be the  $d$ -th component function of  $f$ .
- Let  $g : X \rightarrow \prod_{j < d} Z_j$  be defined by  $g(x) = f(x) \restriction d$ .  
(Technically,  $g$  must be a slight extension of this, but never mind.)
- Let  $M$  be a good countable elementary submodel of  $H(\theta)$  for a sufficiently large regular cardinal  $\theta$ .



# The First Claim

First, we can prove the following claim.

## Claim

Let  $r \in Z_d \setminus \text{Cl}(Z_d \cap M)$ . Then, there exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$f_d^{\leftarrow} \{r\} \subseteq \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} I(K_{i_n}, M, p_n)$$

# The First Claim

First, we can prove the following claim.

## Claim

Let  $r \in Z_d \setminus \text{Cl}(Z_d \cap M)$ . Then, there exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$f_d^{\leftarrow} \{r\} \subseteq \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} I(K_{i_n}, M, p_n)$$

## The Second Claim

We can remove the reference to  $r$ , which does not belong to  $M$ , by proving the following claim.

### Claim

There exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$\prod_{j < d} Z_j = g^{\rightarrow} \left( X \cap \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} I(K_{i_n}, M, p_n) \right)$$

$I(K_{i_n}, M, p_n)$  can be 'approximated' by elements of  $M$ . So, we can almost describe this phenomenon within  $M$ .

## The Second Claim

We can remove the reference to  $r$ , which does not belong to  $M$ , by proving the following claim.

### Claim

There exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$\prod_{j < d} Z_j = g^{\rightarrow} \left( X \cap \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} I(K_{i_n}, M, p_n) \right)$$

$I(K_{i_n}, M, p_n)$  can be 'approximated' by elements of  $M$ . So, we can almost describe this phenomenon within  $M$ .

## The Second Claim

We can remove the reference to  $r$ , which does not belong to  $M$ , by proving the following claim.

### Claim

There exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$\prod_{j < d} Z_j = g^{\rightarrow} \left( X \cap \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} I(K_{i_n}, M, p_n) \right)$$

$I(K_{i_n}, M, p_n)$  can be 'approximated' by elements of  $M$ . So, we can almost describe this phenomenon within  $M$ .

# The Third Claim

An argument with elementarity proves the following third claim.

## Claim

There exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$\prod_{j < d} Z_j = g^{\rightarrow} \left( X \cap \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} \{ \eta(K_{i_n}, M, p_n), \zeta(K_{i_n}, M, p_n) \} \right)$$

By 'combining'  $\pi_{i_n}^{\leftarrow} \{ \eta(K_{i_n}, M, p_n), \zeta(K_{i_n}, M, p_n) \}$  for all  $n < \hat{n}$ , we can finish proving the main lemma.

## The Third Claim

An argument with elementarity proves the following third claim.

### Claim

There exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$\prod_{j < d} Z_j = g^{\rightarrow} \left( X \cap \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} \{ \eta(K_{i_n}, M, p_n), \zeta(K_{i_n}, M, p_n) \} \right)$$

By 'combining'  $\pi_{i_n}^{\leftarrow} \{ \eta(K_{i_n}, M, p_n), \zeta(K_{i_n}, M, p_n) \}$  for all  $n < \hat{n}$ , we can finish proving the main lemma.

## The Third Claim

An argument with elementarity proves the following third claim.

### Claim

There exist finite sequences  $\langle i_n | n < \hat{n} \rangle$  and  $\langle p_n | n < \hat{n} \rangle$  such that for all  $n < \hat{n}$ ,  $i_n < d$  and  $p_n \in K_{i_n}$ , and

$$\prod_{j < d} Z_j = g^{\rightarrow} \left( X \cap \bigcup_{n < \hat{n}} \pi_{i_n}^{\leftarrow} \{ \eta(K_{i_n}, M, p_n), \zeta(K_{i_n}, M, p_n) \} \right)$$

By 'combining'  $\pi_{i_n}^{\leftarrow} \{ \eta(K_{i_n}, M, p_n), \zeta(K_{i_n}, M, p_n) \}$  for all  $n < \hat{n}$ , we can finish proving the main lemma.



# Toward separability

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem.

Let  $e = |\{j < d + 1 \mid Z_j \text{ is separable}\}|$ . Without loss of generality, we may assume that for all  $j < d + 1$ ,  $Z_j$  is separable if and only if  $j < e$ .

By repeatedly applying the main lemma, we can prove the following:

- $e \geq 2$ , and
- there exists a compact LOTS  $\tilde{K}_i$  for each  $i < e - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < e-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < e} Z_j$ .

Note that  $\prod_{j < e} Z_j$  is separable.

# Toward separability

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem.

Let  $e = |\{j < d + 1 \mid Z_j \text{ is separable}\}|$ . Without loss of generality, we may assume that for all  $j < d + 1$ ,  $Z_j$  is separable if and only if  $j < e$ .

By repeatedly applying the main lemma, we can prove the following:

- $e \geq 2$ , and
- there exists a compact LOTS  $\tilde{K}_i$  for each  $i < e - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < e-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < e} Z_j$ .

Note that  $\prod_{j < e} Z_j$  is separable.

## Toward separability

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem.

Let  $e = |\{j < d + 1 \mid Z_j \text{ is separable}\}|$ . Without loss of generality, we may assume that for all  $j < d + 1$ ,  $Z_j$  is separable if and only if  $j < e$ .

By repeatedly applying the main lemma, we can prove the following:

- $e \geq 2$ , and
- there exists a compact LOTS  $\tilde{K}_i$  for each  $i < e - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < e-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < e} Z_j$ .

Note that  $\prod_{j < e} Z_j$  is separable.

## Toward separability

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem.

Let  $e = |\{j < d + 1 \mid Z_j \text{ is separable}\}|$ . Without loss of generality, we may assume that for all  $j < d + 1$ ,  $Z_j$  is separable if and only if  $j < e$ .

By repeatedly applying the main lemma, we can prove the following:

- $e \geq 2$ , and
- there exists a compact LOTS  $\tilde{K}_i$  for each  $i < e - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < e-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < e} Z_j$ .

Note that  $\prod_{j < e} Z_j$  is separable.

## Toward separability

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem.

Let  $e = |\{j < d + 1 \mid Z_j \text{ is separable}\}|$ . Without loss of generality, we may assume that for all  $j < d + 1$ ,  $Z_j$  is separable if and only if  $j < e$ .

By repeatedly applying the main lemma, we can prove the following:

- $e \geq 2$ , and
- there exists a compact LOTS  $\tilde{K}_i$  for each  $i < e - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < e-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < e} Z_j$ .

Note that  $\prod_{j < e} Z_j$  is separable.

## Toward separability

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem.

Let  $e = |\{j < d + 1 \mid Z_j \text{ is separable}\}|$ . Without loss of generality, we may assume that for all  $j < d + 1$ ,  $Z_j$  is separable if and only if  $j < e$ .

By repeatedly applying the main lemma, we can prove the following:

- $e \geq 2$ , and
- there exists a compact LOTS  $\tilde{K}_i$  for each  $i < e - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < e-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < e} Z_j$ .

Note that  $\prod_{j < e} Z_j$  is separable.

## Toward separability

Let  $d$ ,  $K_i$ ,  $X$ ,  $Z_j$ , and  $f$  be as in the assumption of the main theorem.

Let  $e = |\{j < d + 1 \mid Z_j \text{ is separable}\}|$ . Without loss of generality, we may assume that for all  $j < d + 1$ ,  $Z_j$  is separable if and only if  $j < e$ .

By repeatedly applying the main lemma, we can prove the following:

- $e \geq 2$ , and
- there exists a compact LOTS  $\tilde{K}_i$  for each  $i < e - 1$ , a countably compact subspace  $\tilde{X}$  of  $\prod_{i < e-1} \tilde{K}_i$ , and a continuous surjection  $\tilde{f} : \tilde{X} \rightarrow \prod_{j < e} Z_j$ .

Note that  $\prod_{j < e} Z_j$  is separable.

# Compactness

By using the fact that  $\prod_{j < e} Z_j$  is separable, there exist a separable compact LOTS  $K'_i$  for each  $i < e - 1$ ,  $X'$  a compact subspace of  $\prod_{i < e-1} K'_i$ , and a continuous surjection  $f' : X' \rightarrow \prod_{j < e} Z_j$ .

Now, by using the original Mardešić Conjecture, there exist at least two factors  $Z_j$  that are compact and metrizable.



# Compactness

By using the fact that  $\prod_{j < e} Z_j$  is separable, there exist a separable compact LOTS  $K'_i$  for each  $i < e - 1$ ,  $X'$  a compact subspace of  $\prod_{i < e-1} K'_i$ , and a continuous surjection  $f' : X' \rightarrow \prod_{j < e} Z_j$ .

Now, by using the original Mardešić Conjecture, there exist at least two factors  $Z_j$  that are compact and metrizable.

# Mardešić's Theorem

In fact, S. Mardešić proved that the Mardešić Conjecture holds when all  $Z_j$ 's are separable. So, we can finish proving the main theorem without using the full Mardešić Conjecture. Thus, this argument gives another proof of the (original) Mardešić Conjecture.

# Mardešić's Theorem

In fact, S. Mardešić proved that the Mardešić Conjecture holds when all  $Z_j$ 's are separable. So, we can finish proving the main theorem without using the full Mardešić Conjecture.

Thus, this argument gives another proof of the (original) Mardešić Conjecture.

# Mardešić's Theorem

In fact, S. Mardešić proved that the Mardešić Conjecture holds when all  $Z_j$ 's are separable. So, we can finish proving the main theorem without using the full Mardešić Conjecture. Thus, this argument gives another proof of the (original) Mardešić Conjecture.

# Open Questions

## Question

Can we extend the main theorem to a wider class than LOTS?

## Question

What can we say when there is a continuous surjection from  $\prod_{i < d} K_i$  onto  $\prod_{i < d} Z_i$ ?

We know that if all  $Z_i$ 's are nonseparable, then  $Z_i$  is an image of some LOTS.

# Open Questions

## Question

Can we extend the main theorem to a wider class than LOTS?

## Question

What can we say when there is a continuous surjection from  $\prod_{i < d} K_i$  onto  $\prod_{i < d} Z_i$ ?

We know that if all  $Z_i$ 's are nonseparable, then  $Z_i$  is an image of some LOTS.