ISI-BUDS Lab for Generalized Linear Models

Babak Shahbaba, PhD University of California at Irvine (UCI)

 Examine how the left hippocampus volume lhippo changes as we get older (use age).

- Examine how the left hippocampus volume lhippo changes as we get older (use age).
- Visualize your model.

- Examine how the left hippocampus volume lhippo changes as we get older (use age).
- Visualize your model.
- Examine the relationship between education on lhippo.

- Examine how the left hippocampus volume lhippo changes as we get older (use age).
- Visualize your model.
- Examine the relationship between education on lhippo.
- Visualize your model.

- Examine how the left hippocampus volume lhippo changes as we get older (use age).
- Visualize your model.
- Examine the relationship between education on lhippo.
- Visualize your model.
- Now consider the relationship between the left hippocampus volume and both age and education.

- Examine how the left hippocampus volume lhippo changes as we get older (use age).
- Visualize your model.
- Examine the relationship between education on lhippo.
- Visualize your model.
- Now consider the relationship between the left hippocampus volume and both age and education.
- Interpret your estimates of the regression coefficients.

 Divide the data into two equal subsets called "Training" and "Test".

- Divide the data into two equal subsets called "Training" and "Test".
- Use the Training set to build a linear regression model with naccmmsse as the response and age, gender, and education as predictors.

- Divide the data into two equal subsets called "Training" and "Test".
- Use the Training set to build a linear regression model with naccmmsse as the response and age, gender, and education as predictors.
- Use your model to predict the values of naccmmsse in the test set.

- Divide the data into two equal subsets called "Training" and "Test".
- Use the Training set to build a linear regression model with naccmmsse as the response and age, gender, and education as predictors.
- Use your model to predict the values of naccmmsse in the test set.
- Compare your predictions with the true values.

• Create a new variable called disease.status by transforming the variable diagnosis to a new feature with two categories: no symptoms (0) versus mild or strong symptoms (1).

- Create a new variable called disease.status by transforming the variable diagnosis to a new feature with two categories: no symptoms (0) versus mild or strong symptoms (1).
- Next, we want to build a logistic regression model for disease status with age, education, naccicv, female as its predictors.

- Create a new variable called disease.status by transforming the variable diagnosis to a new feature with two categories: no symptoms (0) versus mild or strong symptoms (1).
- Next, we want to build a logistic regression model for disease status with age, education, naccicv, female as its predictors.
- Interpret your estimates of the regression coefficients.

- Create a new variable called disease.status by transforming the variable diagnosis to a new feature with two categories: no symptoms (0) versus mild or strong symptoms (1).
- Next, we want to build a logistic regression model for disease status with age, education, naccicv, female as its predictors.
- Interpret your estimates of the regression coefficients.
- As before, divide the dataset into two subsets, fit the above logistic regression model to one subset and use the resulting model to predict the disease status for the other subset.
 Evaluate the performance of your model.