R6.11: relation yield across a prime conductor gap

Question. Is the index-calculus class factor c the same on the crater ($\mathrm{End} = \mathcal{O}_K$) and the floor ($\mathrm{End} = \mathbb{Z} + \ell\mathcal{O}_K$) of an ℓ-volcano, $\ell \in \{5, 7, 11, 13\}$, $p \in [2^{20}, 2^{22})$? (Toy test of Conjecture 10′.)

Verdict: confirmed at toy size; the falsifier did not fire. 24 instances (6 per ℓ), same $\#E = N$ prime on both levels (certified), $|\mathrm{Aut}| = 2$, $h = 1$; every crater⟶floor edge certified by an Elkies kernel and Vélu codomain. Exact census at $m = 2, 3$, $|F| \in \{60, 80, 100, 200\}$: every crater/floor ratio mean lies within 0.5% of 1 and every mean ± 2 sd interval contains 1 (measured).




Figure 1: Crater/floor$_0$ class-factor ratio vs ℓ; bars mean ± 2 sd over 6 instances, dots per instance.




Figure 2: Measured c over the derived birthday reference, every curve; crater and floor interleave.

Limits. At $m = 2$, $|F|^2_2 \ll N$ forces $c \approx 1$ on any curve; $m = 3$ is the informative arm. Plain x-frame bases only: an oracle that uses $\mathrm{End}(E)$ explicitly is not tested. Toy sizes; $h(D) = 1$ craters; one vertical step. No sub-rho or deployed-curve claim.

Full tables, commands and raw data: `RESULTS.md`, `raw/`.