Residual quartics and a complete Hess transfer

Exact GF(16)/GF(4096) curves, verified subgroup transport, and measured attack cost

Result. A fixed 30,000-draw search independently verified three plane-quartic equations in Jacobian isogeny classes absent from the earlier local screen. A separate, published genus-four Hess construction carried all four exact source subgroups through full split-divisor relation solves. Each measured transfer was at least 512 times slower than same-source rho in this backend. The new quartics still lack explicit source-to-quartic maps.

1. Exact quartic witnesses

Let K=GF(2)[w]/(w^12+w^7+w^6+w^5+w^3+w+1), k=GF(2)[a]/(a^4+a+1). Each source is E: Y^2+XY=X^3+w^5 X^2+b, with polynomial-basis integer encoding for b and point coordinates. Quartic vectors use x^i y^j z^(4-i-j), descending i and j.

#E=#Jac	b	prime r	G / Q=[7]G	(#C(k),#C(k^2),#C(K))	LMFDB class
4006	198	2003	(199,2783) / (738,3018)	(17,257,3824)	3.16.a_a_adn
4054	32	2027	(33,349) / (2098,1182)	(17,257,3968)	3.16.a_a_abr
4078	44	2039	(45,1367) / (47,3257)	(17,257,4040)	3.16.a_a_at
4126	63	2063	(62,1499) / (2633,1798)	(17,257,4184)	3.16.a_a_bd

```
4006: [1,14,11,0,8,3,6,8,7,1,15,9,15,12,3]
4054: [1,15,8,7,1,6,10,2,10,7,10,3,12,3,12]
4078: [1,4,10,9,0,6,9,5,3,8,15,8,2,12,2]
4126: [1,3,13,3,5,4,5,4,15,11,15,7,4,9,1]
```

Every quartic is geometrically smooth. Alternate-chart recounting and exact Jacobian orders were independently replayed; rational point-difference classes reach the selected prime subgroups. The first three orders were absent from the earlier 2,048-draw screen. Order 4126 already had a different local quartic witness. All four isogeny classes are cataloged in LMFDB, whose stored models for these labels are hyperelliptic only. Publication novelty of the equations remains unresolved.

2. Known direct Hess genus-four subgroup transfers

For each source in the table, choose Hess descent parameter c=1. The verified genus is four and Magma produced a nonzero order-r divisor-class image. For the order-4126 source E: Y^2+XY=X^3+w^5 X^2+63, the exact model and evaluated images are:

```
C: y^2+(a^8 x^4+a^3 x^2+x+a^4)y
   = a^11 x^9+a^13 x^8+a x^5+a^10 x^3
     +a x^2+a^12 x+a^12.
#Jac(C)(k)=57764=28*2063.

h(G)=(x^4+a^6 x^3+a^13 x^2+x+a^8,
      a^2 x^3+x^2+a^6 x+a^9, 4)
h(Q)=(x^4+a^4 x^2+a^5 x+a^10,
      a^12 x^3+a^13 x^2+a^2 x+a^12, 4)
```

The other three exact genus-four equations and Mumford image triples appear in the appendix. For every source, Magma checked h(G) nonzero, [r]h(G)=0, [r]h(Q)=0, and h(Q)=[7]h(G). The alternate plane model of the 4126 control has degree nine; Magma's documented line-sieve gate q >= d! fails because 16 < 9!. All four full runs used hyperelliptic split-divisor factor bases.

3. Full relation solve and cost

Each run factored the Mumford u-polynomial of sampled R=u h(G)+v h(Q) and reconstructed every retained R in Jac(C). Projecting by Jacobian cofactor m gives rows sum(n_i t_i)-m v x = m u (mod r). Independent Python elimination of the saved full-rank rows recovered x=7 for all four sources; Magma checked Q=[7]G on each source.

#E / r	Jacobian / m	FB; attempts; rows/rank	Full transfer	Rho per target	Ratio
4006 / 2003	88132 / 44	21; 211; 22/22	0.350 s	0.000684 s	512x
4054 / 2027	56756 / 28	13; 506; 14/14	0.450 s	0.000645 s	698x
4078 / 2039	57092 / 28	13; 471; 18/14	0.450 s	0.000645 s	698x
4126 / 2063	57764 / 28	13; 431; 15/14	0.430 s	0.000625 s	688x

Each rho time is amortized over 512 target-verified trials on that same source subgroup. Full transfer times include cover, map, factor-base setup, relations, linear solve, and source verification. They exclude field/source-object setup, as does the rho timer. The ratio range is 512-698. These are calculator-backend measurements, not EKS-qualified benchmarks; the configured AWS caller is outside the required Iso Lab account. These small-order timings are not asymptotic bounds.

4. Claim boundary and sources

The quartics have exact equations and matching Jacobian classes, but no evaluated elliptic-to-quartic maps or complete attack costs. The genus-four controls have complete maps and solves, but are published and slower than rho. No newly vulnerable family survives these gates. The prior degree-at-most-three exclusion applies only to the listed order-4126 hyperelliptic genus-three models; it does not rule out the genus-four construction or new quartic correspondences.

Primary sources: Hess, *Generalising the GHS Attack on the Elliptic Curve Discrete Logarithm Problem* (LMS J. Comput. Math.); Magma Handbook, *Weil Descent in Characteristic Two* and *Linear Equivalence of Divisors / Index Calculus*. Exact scripts, receipts, audits, and checked Lean arithmetic/projection theorem accompany this PDF in the PR.

Appendix. Exact Hess outputs for the other three sources

All equations are over k=GF(16) with a^4+a+1=0. Source points and quartic vectors are on page 1. Each triple below is a reduced Mumford representative of the evaluated subgroup map.

```
ORDER 4006, b=198, r=2003, #Jac=88132=44*2003
C: y^2+(a^8*x^4+a^10*x^2+a^11*x+a^4)y
   = a^5*x^9+a^4*x^8+a^9*x^5+x^4+a^11*x^3
     +a^2*x^2+a^14*x+a^4.
h(G)=(x^4+a^3*x^2+a^6*x+a^8, a^6*x^2+a^12*x+a^13, 4)
h(Q)=(x^4+x^3+a^10*x^2+a^9*x+a^12,
      a^10*x^3+a^3*x^2+a^4*x+a^11, 4)

ORDER 4054, b=32, r=2027, #Jac=56756=28*2027
C: y^2+(a^8*x^4+a^13*x+a^9)y
   = a^10*x^9+a^12*x^8+a^10*x^4+a^5*x^3
     +a^7*x^2+a^11*x+a^10.
h(G)=(x^3+a^2*x^2+a^4*x+a^8, x^2+a^11*x+a, 3)
h(Q)=(x^4+a^10*x^3+a^3*x^2+a^2*x+a^9,
      a^9*x^3+a^3*x^2+a^5*x, 4)

ORDER 4078, b=44, r=2039, #Jac=57092=28*2039
C: y^2+(a^8*x^4+a^14*x^2+x+a^9)y
   = a^14*x^9+x^8+a^11*x^5+a^11*x^4
     +a^13*x^3+a^8*x^2+a^14*x+a^5.
h(G)=(x^4+a^6*x^3+a^5*x^2+x+a^10,
      a^5*x^3+a*x^2+a^13*x+a^14, 4)
h(Q)=(x^4+a^12*x^3+a^5*x^2+a^2,
      a^6*x^3+x^2+a^6*x+a^4, 4)
```