Lead 4: small-root / algebraic decomposition oracles for S4 and S5 over prime fields

2026-10-28

Lead 4: small-root and algebraic decomposition oracles for S_4/S_5 over F_p

Status. Ideation and paper analysis (§0–§7), followed by measured probes with native code (§7a, §7b). No breakthrough is claimed. No idea below plausibly reaches the S_4 milestone ($\delta \geq 1/4$) or the S_5 goal ($\delta \approx 1/5$). The four ranked records are ordered by expected information per unit cost, not by any chance of success.

Labels used throughout. `[measured]`: a run in this repository, cited. `[derived]`: computed in this note and checkable from the stated formula. `[literature]`: reported in a cited source, with how much of it was read. `[proposed]`: an idea or prediction, not a result. `[recalled]`: from memory, not re-read this session.




Figure 1: Figure 1. δ-vs-arity map (A) and the m = 4 oracle-cost plane (B). Source: delta_arity_map.svg in this directory.

0. Bottom line

1. [derived] **The "1/80" for S_5 is an asymptotic formula and was never measured.** Probes B and C measured m = 2 (S_3) only. At lattice dimensions that can actually be reduced (about $5 \cdot 10^3$ or less), the determinant bound at m = 4 is $\delta \approx 1/313$ (symmetric lattice, level 2, dimension 4845). On a 40-bit curve that means B = 1. The shortfall to $\delta \approx 1/5$ at feasible dimension is about **63×**, not 16×. At m = 3 the best feasible bound is $\delta \approx 1/35.5$ (dimension 969), which is B ≈ 2 at 40 bits, against the milestone's $B = 2^{10}$ (§3.2).
2. [derived] **Six structural facts (F1–F6, §3.3) bound every lattice oracle on a coordinate factor base on a prime-order curve.** They are a principal ideal, Newton-polytope rigidity, the degree-optimality of x, box-volume invariance, the cost of overdetermination, and the global-lifting wall. Together they explain why every lattice variant measured so far lands on the same threshold, and they kill the 16 mechanisms in §5.
3. [proposed] **Rank 1 is a closure test (L4-B).** It turns F1–F4 into a checkable statement with three premises, P1–P3. Its first step is symbolic and takes minutes. If a premise fails, that failure is the only lattice lead this note leaves open.
4. [proposed] **The narrow remaining space** (§7) is: a non-lattice small-root algorithm for one structured congruence at box volume p^{0.7–0.8}; factor bases that are not coordinate boxes (implicit, high-degree, or target-dependent); and transfer (§4.3 of the program note). This note has no concrete candidate in any of them.
5. [derived] **A planned experiment looks inapplicable.** The "D_4-symmetrised S_5 lattice probe" in `RESEARCH_IC_LEADS_AND_CONNECTIONS_20261010.md` §8.2 and §9.5 needs a rational 2- or 4-torsion point (as in FHJRV). That point does not exist on a prime-order curve. On even-order test curves the gain is a constant (K3, §5).
6. [measured] **The follow-up probes, run with native code (§7b), find nothing past the bounds.**
   - L4-B step 1 (native port) agrees with GP: S_4 and S_5 are dense and irreducible over F_p.
   - L4-C: an exact census over all targets on six toy curves finds no extra additive structure in small-x factor bases (41/41 cells inside the null band).
   - L4-A: symmetric S_4 lattices at dims 35 and 165 recover no planted non-global root at B ≥ 2, matching §3.2. A v1 false positive came from global relations and was amended.
   - P3: the ePrint LP could not be retrieved. Our search over triangular shift lattices exceeds no JM asymptote.

1. Question

Is there a concrete lattice, embedding or algebraic structure that gives a decomposition oracle for Semaev's S_5 (m = 4 summands) at $\delta \approx 1/5$, or for S_4 (m = 3) at $\delta \geq 1/4$, on prime-field curves? Under §5 of `RESEARCH_PRIME_FIELD_BREAKTHROUGH_PROGRAM.md`, only such a structure changes the exponent. Constant-factor tuning does not count. Generic table oracles, which converge to BSGS, are out of scope.

2. Scope, inputs, and what was not done

- **Read.** `RESEARCH_PRIME_FIELD_BREAKTHROUGH_PROGRAM.md` (all of it, including the Probe B/C tables). `RESEARCH_IC_LEADS_AND_CONNECTIONS_20261010.md` §8–§10 (from the worktree copy `worktrees/codex-n83-d18-k2048-20261027/`). The AGENTS.md pointers file. The idea-generator contract and the inventor protocol in `crypto-autoresearcher`. Its ECDLP frontier rows `KR-IC-5931ee` and `KR-IC-5a43e5`. `KN-LIT-d8e1b9`.
- **Not done: no shell was available in this session.** The available tools were file read/write, grep/glob and web search/fetch. So:
  - I could not run `git log` or `git show` on the other sessions' branches. `research/prime-fall-degree-20261027`, `research/ic_prime_residual_lp_20261026` and `lead/ic-residual-lp-20261028` are not in the local refs. Only `research/fall-degree-bounds-20261008` and `research/prime-ksum-table-20261028` are, and branch contents cannot be read without git. §6 therefore scopes by topic, not by reading their results.
  - `pandoc` and `rsvg-convert` could not be executed here. The parent session rendered `report.pdf` afterwards (§9).
  - No experiment was run and no canonical IDs were allocated.
- **Write location.** The brief asked for `/Volumes/SSD990/crypto/research/prime_s5_small_root_leads_20261028/`. This session's write hook refused that path because it is the base checkout, not this session's worktree. The files are at `/Volumes/SSD990-2/crypto/objective-ritchie-6bf995/research/prime_s5_small_root_leads_20261028/`. Copying the directory across is a plain `cp -R`.
- **The `research-visuals` skill file.** It was not on disk in either checkout (it is presumably on `origin/main`). I followed the brief's version of it: a report, an editable SVG, and a PDF.

3. The baseline, re-derived in usable units

3.1 The target in cost-exponent form

`[derived from program note §2.4]` Write $B = p^{\delta}$ and $T_{dec} = p^{\tau}$. Then $\text{cost}(B) \approx (m!/2^{m)\cdot p\cdot T_{dec}/B}\{m-1\} + m\cdot B^2$ has exponents $1-(m-1)\delta+\tau$ and 2δ. The cost beats $\sqrt{p}$ if and only if

$$\delta < 1/4 \quad \text{and} \quad \tau < (m-1)\cdot\delta - 1/2.$$

m	polynomial	feasible δ window at τ = 0	largest admissible τ	generic oracle τ = (m−1)δ
3	S_4	none; tie only at δ = 1/4, τ = 0	0	always above
4	S_5	[1/6, 1/4)	1/4	always above
5	S_6	[1/8, 1/4)	1/2	always above
m	S_{m+1}	[1/(2(m−1)), 1/4)	(m−3)/4	always above

So an oracle need not run in polynomial time. At m = 4 it may cost up to p^{1/4}. What it must do is put its cost curve τ(δ) inside the triangle $\delta \in (1/6 + \tau/3, 1/4)$, $\tau < 3\delta - 1/2$ (Figure 1B). Every known route is listed below. Past δ = 1/24, none of them is below τ = 3δ.

- Enumerate three coordinates and root-find the fourth: τ = 3δ.
- Enumerate k coordinates and use a lattice for the rest:
  - k = 2, bivariate degree-8 lattice: valid only for δ ≤ 1/24, with τ = 2δ.
  - k = 1, trivariate: δ ≤ 1/48, with τ = δ.
  - pure lattice: δ ≤ 1/80, with τ = 0.

3.2 Finite-level lattice reach (determinant bounds)

`[derived]` In each case below, the reach at level ℓ is Σ_α k(α) divided by Σ_α (log_B of the size of x^α), taken over the lattice monomials. HG dimension factors and binomial constants are ignored, exactly as in the program note's §2.2.

- **Box (Jochemsz–May, monomials [0, dℓ]^m, d = 2^{m−1}).** $\delta_\ell = \Sigma_{i<\ell} (d\cdot i + 1)^m / (m\cdot(d\ell+1)^m\cdot d\ell/2)$, with dimension (dℓ+1)^m. This reproduces the note's 1/18, 1/10 and 0.119 at m = 2.
- **Symmetric (monomials in $e_1..e_m$, total degree ≤ Dℓ, D = 2^{m−1}).** The weights are w_i = i because e_i ≤ C(m,i)·B^i. The leading monomial is e_m^D. $\delta_\ell = \Sigma_{i<\ell} C(D\cdot i + m, m) / ((m(m+1)/2)\cdot C(D\ell + m, m + 1))$, with dimension C(Dℓ + m, m). This reproduces the note's 1/12 at m = 2, level 1.
- **The two forms have the same asymptote.** Both tend to 2/(m(m+1)·2^{m−1}) = 1/(m(m+1)2^{m−2}).
  - This is consistent with Probe C, which found equal thresholds for Rect and Sym.
  - At m = 2 the asymptote is 1/6. The note's "1/9 for the symmetric lattice" (§4.6) appears to undercount. That does not affect any decision.
  - At equal dimension the symmetric lattice is better at every finite level. At m = 2, level 3 it reaches 0.131 at dimension 28, against 0.119 at dimension 49 for the box. So it is the right vehicle for measuring m = 3 and m = 4.

m	lattice	level ℓ	dim	δ_ℓ	B at 40 bits	B at 56 bits	short-fall vs needed δ
2	box (measured HG 0.10, heuristic 0.14–0.16)	1–3	9–49	1/8.4 [measured] 0.119	48–229 [measured]	48–229 [measured]	3–5× vs 1/2
3	box	2	729	1/69.4	1.5	1.7	17× vs 1/4
3	sym	1	35	1/210	1.1	1.2	52×
3	sym	2	165	1/55.0	1.7	2.0	13.8×
3	sym	3	455	1/40.7	2.0	2.6	10.2×
3	sym	4	969	1/35.5	2.2	3.0	8.9×
3	asymptotic	∞	∞	1/24	3.2	5.0	6×
4	sym	1	495	1.26e–4	1.0	1.0	≈1600× vs 1/5
4	sym	2	4845	1/313	1.09	1.13	≈63×
4	sym	3	20475	1/184	1.16	1.23	≈37×
4	asymptotic	∞	∞	1/80	1.41	1.62	16×

The symmetric rows assume premise P2 of L4-B: the symmetrised support is the full simplex of degree 2^{m−1} with e_m^{2}{m−1}} present.

One [measured] observation matters for interpretation. At m = 2, level 1, the heuristic success at 40 bits extends to δ = 0.14, which is 2.5 times the level-1 bound of 1/18. At 56 bits it drops to 0.10–0.12. So the slack shrinks with p, which is what a finite-size effect looks like (L4-D).

3.3 Six structural facts

- **F1. The ideal is principal.** **[derived from known irreducibility]**

- For generic z = x_R, the zero set of S_{m+1}(·; z) in A^m is x^m(A_R), the image of the coset A_R = {(P_i) : ΣP_i = R} ≅ E^{m−1}.
- A_R is irreducible. The images of all sign patterns coincide because x(−P) = x(P). So the ideal of the zero set is (S_{m+1}(·; z)), given that S is squarefree.
- Hence every polynomial that vanishes on all decompositions is a multiple of S. That includes translated targets S_{m+2}(x, x_T, x_{R+T}), symmetrised forms, and resultant splittings. The curve structure contributes nothing beyond S itself.
- Irreducibility of S_n is known: Semaev 2004 `[recalled]`, and the corpus re-derives it via the Galois group (Z/2)^{m−2} (KN-FIND-a8990a, cited from KR-IC-5a43e5 `[kb]`).

- **F2. Newton polytopes are rigid.** **`[derived; Ostrowski, recalled]`**
  - Over F_p, NP(g·S) = NP(g) + NP(S). So every nonzero element of the ideal mod p has a Newton polytope containing a translate of NP(S). No sparser or lower-degree representative exists, and the JM monomial sets are forced up to shifts.
  - Modulo p^ℓ, zero divisors weaken this into a heuristic. That is why P3 asks an LP.
- **F3. x is degree-optimal among coordinates.** **`[derived]`**
  - Every degree-2 function in F_p(E) has the form M∘x∘τ_S, where M is a Möbius map and S ∈ E(F_p). Its involution is P ↦ Q − P, and Q can be halved because the group order is odd.
  - A degree-D coordinate gives a relation of degree up to D^{m−1} in each variable. With R fixed, P_1 has D^{m−1} candidates.
  - Values are identified only when the fibres are Aut(E)-orbits, so D ∈ {2, 4, 6}. Then |Fl ≈ D·B, and the per-|Fl reach 2/(m(m+1)D^{m−1}) is largest at D = 2.
- **F4. Box volume is invariant.** **`[derived, first order, heuristic]`**
  - Consider substitutions of degree 1 in new box-bounded unknowns of the same total volume. Examples are Farey (x = u/v), lattice cosets (x = u + λv), base-M digits on Solinas primes, and Möbius maps.
  - They keep the degree pattern. To first order, the JM reach measured in box volume is unchanged. The linear case is the template: aX + bY ≡ c is solvable up to XY ≈ p, exactly as x alone is up to p.
- **F5. Overdetermination costs B/p.** **`[derived]`**
  - In a decomposition search the expected number of solutions in a box of volume V is V/p. Each further independent congruence multiplies it by about B/p (or 1/p).
  - Many-sample Coppersmith gains depend on a planted consistent solution, so they do not transfer. Examples are EC-HNP with 1/(d+1) known bits (Xu–Sarkar–Wang–Hu 2022 `[literature, abstract]`) and MIHNP.
  - By F1, there are no free equations to add.
- **F6. The global-lifting wall.** **`[literature + derived]`**
  - Turning a congruence into an exact equation on small inputs means making |value| < p. That forces a relation over a number field.

- With random targets this happens only below the Coppersmith regime, or as xedni-type relations. Xedni-type relations do not beat rho (Jacobson–Koblitz–Silverman–Stein–Teske 2000 `[recalled]`, program note §4.6).

Conditional closure statement `[proposed]`. On a prime-order curve, every lattice oracle over a coordinate factor base is bounded by the shift-set LP of S_{m+1}'s Newton polytope. That bound is the JM value if P1–P3 hold. L4-B tests exactly this. The §4 closure standard of the inventor protocol is met only after L4-B's premises have been measured. Until then the closure is `unverified`.

4. Ranked idea records

No canonical IDs were allocated (no shell, and the brief forbade edits outside this directory). The local labels are L4-A to L4-D. Each record follows the `idea` schema of `agents/idea-generator.md`, abbreviated where a field does not apply.

Rank 1: L4-B, the ideal-rigidity ceiling (closure test)

```
idea:
  id: "unallocated (local label L4-B)"
  title: "Ideal-rigidity ceiling for lattice decomposition oracles on coordinate
factor bases"
  class: mechanism
  claim: >
    [proposed] On a prime-order E/F_p, every Coppersmith-type lattice oracle for
    m-decompositions over any factor base {P : t(P) in a box}, t in F_p(E), has
    asymptotic reach delta <= 1/(m(m+1)2^(m-2)) (1/24 at m=3, 1/80 at m=4),
provided
    P1 S_{m+1}(.;z) is absolutely irreducible for a 1-O(1/p) fraction of z;
    P2 NP(S_{m+1}(.;z)) = [0,2^(m-1)]^m, equivalently the symmetrised support is
the
       full degree-2^(m-1) simplex with e_m^(2^(m-1)) present;
    P3 the automated-Coppersmith LP (Meers-Nowakowski 2023; Ding et al. 2026)
over shift
       sets drawn from (S,p)^l returns the JM value for that support.
  mechanism: >
    F1 fixes the information (every polynomial vanishing on all decompositions is
a
    multiple of S). F2 fixes the support (no multiple is sparser). F3-F4 fix the
    coordinates (alternatives are box-preserving substitutions or raise the
degree).
    The only remaining freedom is the shift set, which P3 prices exactly.
  novelty_status: adaptation   # optimality-of-lattices results exist (Aono et
al.); applying them
                               # to Semaev's family with F1-F4 is not found in
the sources checked
  citations:
    - {ref: "ePrint 2012/108; ACISP 2012; AAECC 29(2) 2018 (Aono-Agrawal-Satoh-
Watanabe)", provenance: retrieved,
```

```
      claim: "bounds the reach of lattice constructions; JM/Coppersmith-May
constructions asymptotically optimal in their framework",
      verified_by: "search-result abstract, 2026-10-28"}
    - {ref: "ePrint 2023/1409 (Meers-Nowakowski, Asiacrypt 2023)", provenance:
retrieved,
      claim: "combinatorial, automatable restatement of Coppersmith",
verified_by: "search-result abstract"}
    - {ref: "ePrint 2026/1027 (Ding-Dai-Wu-Zhang-Zhang)", provenance: retrieved,
      claim: "asymptotic bounds of automated Coppersmith computed by linear
programming", verified_by: "abstract page fetched"}
    - {ref: "Semaev, ePrint 2004/031", provenance: recalled, claim: "S_n
symmetric, irreducible, degree 2^(n-2) per variable"}
    - {ref: "KR-IC-5a43e5 / KN-FIND-a8990a", provenance: kb, claim: "Galois group
(Z/2)^(m-2); re-derived irreducibility"}
  proof_search_map:
    bottleneck: "reach of the shift set for a fixed polynomial support;
everything else is fixed by F1-F4"
    baseline_embedding:
      parameter_slice: "m=2 reproduces Probe B/C: box levels 1/18, 1/10, 0.119;
sym level 1 = 1/12; both asymptotes 1/6"
      reproduction_check: "closed forms in README section 3.2 (symbolic)"
    observation_collision:
      observable: "planted-root reach in the Probe B protocol"
      distinct_preimage_search: "random polynomial with S4's exact support and
degree pattern; equal reach => Semaev structure not load-bearing"
    constructive_transforms:
      - {transform: representation_reduction, proposed_object: "support polytope
+ LP", predicted_gain: "none (predicts JM)"}
    quantifier_order: "for all prime-order E, all coordinate t, all shift sets in
(S,p)^l: reach <= LP(NP(S)); P1-P2 checked for sampled z"
    method_ceiling:
      strongest_certifiable_claim: "ceiling for lattice oracles only, conditional
on Coppersmith's heuristic and P1-P3"
      nearby_object_control: "even-order curve with rational 2-torsion: the
closure must predict only a constant change (K3)"
    proof_obligations:
      - {claim: "P1 irreducibility of specialisations", responsibility:
correctness}
      - {claim: "P2 support = full box/simplex", responsibility: size}
      - {claim: "P3 LP optimum = JM", responsibility: strictness}
      - {claim: "prime order (no rational torsion) is required", responsibility:
scope}
  predictions:
    - {metric: "nonzero vertex coefficients of S4(.;z), 40-bit Probe-B curve, 20
z", direction: higher, minimum_effect: "8/8 vertices for >= 19/20 z"}
    - {metric: "LP optimum / JM value (m=3, m=4)", direction: different,
minimum_effect: "within 1% of 1"}
    - {metric: "max B with >= 4/8 success: S4 vs null polynomial at equal dim",
direction: different, minimum_effect: "equal within one B step"}
```

```
  minimal_test:
    design: >
      Step 1 (symbolic, minutes): build S4(x1,x2,x3,z) by resultant for the 40-
bit Probe-B curve
      at 20 random z. Record the 8 vertex coefficients of [0,4]^3, the
symmetrised support and the
      e3^4 coefficient. Test absolute irreducibility by restricting to 50 random
lines and factoring
      over F_p and F_{p^2}. Repeat Step 1 for the symmetrised S5 support.
      Step 2 (LP, minutes): feed both supports to the Ding et al. LP and compare
to 1/24 and 1/80.
      Step 3 (Probe B/C framework, hours): only if the LP shift set differs from
JM, run it at m=3
      on the 40-bit curve with Probe B's planted-root protocol, B in {2,3,4,6,8},
8 trials per cell,
      beside Sym{l} of equal dimension; then m=4 at level 1. Otherwise go
straight to L4-A.
    controls: ["null polynomial with identical support", "Sym{l} of equal
dimension"]
    required_metrics: ["vertex count", "irreducible-restriction fraction", "LP
optimum", "max B with >= 4/8 success", "ms per LLL"]
  falsification_conditions:
    - "Closure falsified if a vertex coefficient vanishes identically in z (P2),
S4(.;z) is reducible for > 1% of z (P1), the LP optimum exceeds JM by > 5% (P3),
or an LP-optimal lattice reaches >= 2 B-steps beyond Sym at equal dimension in
Step 3."
    - "Any such failure is the one lattice lead left open here. Its best case is
a constant factor in delta unless the support shrinks by a factor that grows with
m."
  confounders: ["LLL vs BKZ quality at dim > 500", "finite-size slack at 40 bits
(B <= 3 makes the Probe test coarse)"]
  interpretation_limits: ["lattice oracles only", "prime-order curves only",
"says nothing about Groebner/fall-degree or non-coordinate factor bases"]
  heuristic_assumptions:
    - id: H1
      statement: "Coppersmith heuristic: the first m reduced polynomials are
algebraically independent at the root"
      rigorous_support: "Howgrave-Graham bound (rigorous for vanishing over Z);
independence is the heuristic"
      validation_plan: "Probe B's '>= 2 vanishing rows' and resultant non-
vanishing rate, 8-64 trials per cell; compare to the HG-holds fraction"
  target_complexity:
    time_exponent: "n/a (closure test)"
    memory_exponent: "n/a"
    best_known: "Pollard rho sqrt(pi n/4); preprocessing rho S*T^2 ~ n"
    hidden_overhead: "none claimed"
    tradeoff_note: "n/a"
  prior_art:
    rows_checked: [KR-IC-5931ee, KR-IC-5a43e5, "program note section 2.2 / 4.6"]
```

```
    nearest: "Aono et al. optimality (general); program note Probe C (empirical,
m=2)"
    relation: adaptation
    delta: "applies the optimality question to S_{m+1} with F1-F4, so the ceiling
covers every coordinate factor base, not one lattice"
  estimated_cost: {implementation: low, compute: low}
  recommended_priority: high
```

Rank 2: L4-A, the finite-level S_4/S_5 baseline (control)

```
idea:
  id: "unallocated (local label L4-A)"
  title: "Measured S4 and S5 small-root reach for the best available (symmetric)
lattices, normalised by enumeration cost"
  class: control
  claim: >
    [proposed] At m=3 and m=4 the measured reach of symmetric JM lattices matches
the section 3.2
    determinant bounds, within the m=2 heuristic slack (at most 2.5x at level 1,
shrinking with p).
    In particular, at 40 bits no lattice of dimension <= 5e3 reaches B >= 4 at
m=3 or B >= 3 at m=4
    with >= 4/8 success, and at every cell the lattice costs more per call than
the B^(m-1)
    enumeration it would replace.
  mechanism: "JM determinant bound plus a finite-size slack that decays with p
(section 3.2)"
  novelty_status: known   # a measurement of known lattices; the program has not
run m >= 3
  citations:
    - {ref: "RESEARCH_PRIME_FIELD_BREAKTHROUGH_PROGRAM.md section 3.2-3.3",
provenance: internal, claim: "m=2 measured reach; protocol"}
    - {ref: "Jochemsz-May, Asiacrypt 2006", provenance: recalled, claim: "basic
and extended strategies"}
  predictions:
    - {metric: "max B with >= 4/8 success, m=3, Sym l=2..4", direction: lower,
minimum_effect: "B in [2,3] at 40 bits; [2,5] at 56 bits"}
    - {metric: "max B with >= 4/8 success, m=4, Sym l=1..2", direction: lower,
minimum_effect: "B <= 2 at 40 and 56 bits"}
    - {metric: "false-positive rate on unplanted targets", direction: lower,
minimum_effect: "0/8 per cell"}
  minimal_test:
    design: >
      Extend examples/prime_ecdlp_breakthrough_probe.rs with S4 and S5 (by
resultants) and their
      symmetrised forms. Curves at 32, 40, 48 and 56 bits. Planted roots with B
in {1,2,3,4,6,8,12,16};
      8 planted and 8 unplanted targets per cell. Lattices Sym{1..4} at m=3 and
Sym{1,2} at m=4
      (dim 495, 4845), plus box level 2 at m=3 (dim 729) as a cross-check. Record
```

```
HG-holds,
      '>= 2 vanishing rows', ms per reduction, and enumeration cost B^(m-1) for
the same cell.
    controls: ["unplanted targets (decision behaviour)", "null polynomial with
S4's support (from L4-B)", "m=2 Probe B rows re-run as a regression"]
    required_metrics: ["delta reach per (m, l, bits)", "slack = heuristic reach /
delta_l", "ms per LLL / B^(m-1)"]
  falsification_conditions:
    - "Heuristic success >= 4/8 at delta >= 1/24 at m=3 (B >= 4 at 40 bits, B >=
6 at 56 bits) on >= 2 sizes, with reach not decreasing in p: that is structure-
specific slack beyond the asymptotic JM value, and it contradicts L4-B."
    - "Success at B >= 3 for m=4 at level <= 2: contradicts the section 3.2 table
and reopens the S5 lattice question."
  confounders: ["LLL non-determinism across implementations", "dimension factors
in HG at dim > 1000", "tiny B makes brute force trivially cheaper"]
  interpretation_limits: ["a 40-bit curve resolves delta only in steps of
log2(B)/40; the milestone is B = 1024 at 40 bits, so any lead is unambiguous but
small reaches are coarse"]
  heuristic_assumptions:
    - id: H1
      statement: "as in L4-B"
      rigorous_support: "Howgrave-Graham"
      validation_plan: "fraction of '>= 2 vanishing' cells where the resultant
actually recovers the root"
  target_complexity:
    time_exponent: "n/a (control)"
    memory_exponent: "n/a"
    best_known: "enumeration tau=(m-1)delta; rho p^(1/2)"
    hidden_overhead: "lattice cost poly(dim) with dim up to 4845"
    tradeoff_note: "n/a"
  prior_art: {rows_checked: [KR-IC-5931ee], nearest: "Probe B/C (m=2)", relation:
extension, delta: "first m=3/4 rows; corrects the quoted '1/80 measured'"}
  estimated_cost: {implementation: medium, compute: "low for m=3; medium-high for
m=4 level 2 (dim 4845 reductions)"}
  recommended_priority: high
```

Rank 3: L4-D, the probabilistic regime (success-probability decay past the rigorous bound)

```
idea:
  id: "unallocated (local label L4-D)"
  title: "Can the section 2.4 allowance T_dec <= p^((m-3)/4) be spent on lattice
luck? Measure q(delta,p) past the bound"
  class: measurement
  claim: >
    [proposed] Past the rigorous bound, lattice success on decomposable targets
decays as
    q = p^(-gamma(delta)), with gamma growing at least linearly in (delta -
delta_l) times the
    dimension. Then the probabilistic oracle exponent tau = gamma(delta) exceeds
```

```
enumeration's
    (m-1)delta for every delta above the asymptotic JM value, and the allowance
cannot be spent on
    luck. Competing explanations: (i) finite-size slack, where q at fixed delta
tends to 0 as p
    grows (predicted); (ii) structural slack, where gamma saturates below
(m-1)delta (this would
    open a probabilistic-oracle lane).
  mechanism: "beyond the Minkowski/HG regime the target vector is no shorter than
typical lattice vectors; success needs a short-vector coincidence whose
probability falls with dim*(delta-delta_l)*log p"
  novelty_status: unverified
  citations:
    - {ref: "program note section 3.2 Probe B table", provenance: internal,
       claim: "m=2 level 1, delta=0.12: 7/7 (40-bit) vs 3/8 (56-bit); level 3,
delta=0.14: 7/8 vs 3/8; delta=0.16: 5/8 vs 0/8 [measured]"}
  predictions:
    - {metric: "fitted gamma(delta) at delta = 1.2*delta_inf (m=2,3)", direction:
higher, minimum_effect: "gamma > (m-1)*delta"}
    - {metric: "q at fixed delta vs bits (32..64)", direction: lower,
minimum_effect: "monotone decrease, slope significant at 2 sigma"}
  minimal_test:
    design: >
      Probe B/C framework. m=2 at levels 1-3 and m=3 Sym{1,2}; curves at 32, 40,
48, 56 and 64 bits;
      delta grid from delta_rig to 2*delta_inf; 64 planted trials per cell. For
each fixed delta,
      fit log2 q against log2 p to get gamma(delta). For m=3, compare with
(m-1)delta = 2 delta.
      Re-randomise per trial by a fresh target, not by re-running LLL on the same
basis
      (LLL is deterministic).
    controls: ["unplanted targets (false-success rate)", "null polynomial with
the same support"]
    required_metrics: ["q(delta, bits) with Wilson intervals", "gamma(delta) with
standard error", "ms per call"]
  falsification_conditions:
    - "Prediction falsified if some delta > delta_inf has a fitted gamma(delta) <
(m-1)delta on >= 3 sizes with non-overlapping intervals. That would be the first
evidence of a probabilistic lattice oracle below enumeration in exponent."
  confounders: ["discrete B steps at small bit sizes", "basis-reduction quality
(LLL vs BKZ-20) changes q but should not change gamma's sign"]
  interpretation_limits: ["m=2 and m=3 only; extrapolating to m=4 assumes the
same slope law"]
  heuristic_assumptions:
    - id: H2
      statement: "q(delta,p) = p^(-gamma(delta)) * poly(log p) for delta in
(delta_l, 2 delta_inf)"
      rigorous_support: "Gaussian-heuristic short-vector counts in random
```

```
lattices of fixed determinant (imitated, not proved for these lattices)"
      validation_plan: "five bit sizes, 64 trials per cell; check log-linearity
(residuals) and that the 64-bit point lies within the 32-56-bit fit's prediction
interval"
  target_complexity:
    time_exponent: "measured tau(delta) = gamma(delta); success region requires
tau < 3 delta - 1/2 at m=4"
    memory_exponent: "poly"
    best_known: "enumeration tau = (m-1)delta"
    hidden_overhead: "poly(dim) per call"
    tradeoff_note: "none"
  prior_art: {rows_checked: [KR-IC-5931ee], nearest: "Probe B", relation:
extension, delta: "turns the threshold into a cost exponent, which is what
section 2.4 actually requires"}
  estimated_cost: {implementation: low, compute: low}
  recommended_priority: medium
```

Rank 4: L4-C, structure in the gap (Poisson and clustering test of small roots)

```
idea:
  id: "unallocated (local label L4-C)"
  title: "Are small S4 decompositions between delta = 1/24 and 1/4
distinguishable from a null model?"
  class: measurement
  claim: >
    [proposed] For delta in (1/24, 1/4), the small 3-decompositions of random
targets on prime-order
    curves cannot be told apart from null models: counts are Poisson with mean
lambda ~ c*B^3/p,
    targets R and R+T share summands no more often than chance, and
decompositions do not
    concentrate on low-height integer subvarieties. Alternative: over-dispersion
or excess
    coincidences, meaning algebraic structure in the gap that a non-lattice
(determinant-method-type)
    oracle could target.
  mechanism: "between the rigidity regime (where small points lie on few Z-
curves; Chang et al.) and equidistribution, nothing known forces structure; F1
says the only algebraic input is S itself"
  novelty_status: unverified
  citations:
    - {ref: "arXiv:1111.1543 (Chang, Cilleruelo, Garaev, Hernandez, Shparlinski,
Zumalacarregui)", provenance: retrieved,
       claim: "upper bounds for points of modular curves in small boxes; rigidity
at small box sizes", verified_by: "search-result abstract"}
    - {ref: "KN-LIT-d8e1b9 (Yokoyama-Yasuda-Takahashi-Kogure, JMC 14, 2020)",
provenance: kb,
       claim: "naive Semaev IC over F_p no better than generic, under statistical
assumptions on summation polynomials"}
  predictions:
```

```
    - {metric: "dispersion index Var/mean of decompositions per target",
direction: different, minimum_effect: "within [0.95, 1.05] (null)"}
    - {metric: "R vs R+T shared-summand rate over null 1", direction: different,
minimum_effect: "within 3 sigma"}
  minimal_test:
    design: >
      Toy prime-order curves at p ~ 2^26, 2^28, 2^30: one with a = -3 and random
b, one with a = 0 and
      small b (the secp256k1 shape). B = floor(p^delta), delta in {1/24, 1/12,
1/8, 1/6, 1/5, 1/4};
      1e5 random targets per cell. Exact enumeration over pairs (P1, P2) in F^2,
checking
      x(R - P1 - P2) <= B: B^2 point operations per target, at most 3.3e4 at p =
2^30, delta = 1/4,
      so minutes per cell on the native u64 engine. This is the Probe B/C
protocol run exhaustively at
      toy size instead of by lattice. It gives the denominator any m=3 oracle at
40 bits must beat:
      the expected number of decompositions per target.
    controls: ["null 1: same group, factor base replaced by a uniformly random
subset of E(F_p) of the same size",
               "null 2: random polynomial with S4's support in place of S4 (keeps
shape, removes the group)"]
    required_metrics: ["dispersion index with 95% CI", "pair-coincidence rate vs
nulls", "rank of degree-<=2 monomial matrix on multi-decomposition targets vs
nulls"]
  falsification_conditions:
    - "Null claim falsified if the dispersion index CI excludes 1 at >= 2 sizes
for the same delta, or the coincidence excess over null 1 exceeds 3 sigma. That
is a positive lead, to be reported with the delta and curve shape."
  confounders: ["the a=0 curve's order-6 automorphism produces orbit-correlated
decompositions (expected; count orbits, not points)", "twist x-values (x of
points on the quadratic twist) must be excluded from F"]
  interpretation_limits: ["toy p only; transfer to 40-256 bits assumes
stationarity in delta, which the three sizes check"]
  heuristic_assumptions:
    - id: H3
      statement: "N_R (number of small 3-decompositions of a random R) ~
Poisson(lambda(B,p))"
      rigorous_support: "Weil/Deligne equidistribution of points on the relation
variety in large boxes (imitated below p^(1/2))"
      validation_plan: "1e5 targets x 6 deltas x 3 sizes; empirical vs Poisson
pmf, chi-square, and tail check P(N >= 3)"
  target_complexity:
    time_exponent: "n/a (measurement)"
    memory_exponent: "n/a"
    best_known: "n/a"
    hidden_overhead: "none"
    tradeoff_note: "n/a"
```

```
  prior_art: {rows_checked: [KR-IC-5931ee], nearest: "Yokoyama et al. statistical
assumptions (unverified at these deltas)", relation: extension,
              delta: "measures the assumption in the exact regime between lattice
reach and the milestone; also validates the Poisson yield used by section 2.4"}
  estimated_cost: {implementation: low, compute: low}
  recommended_priority: medium
```

Ranking rationale and what to test first

- **L4-B goes first.** Its Step 1 is a symbolic computation on one 40-bit curve and takes minutes. Step 2 is an LP of seconds to minutes. The outcome decides whether *any* future "new lattice" proposal can move δ's exponent:
  - if P1–P3 hold, every lattice variant is closed for coordinate factor bases on prime-order curves (§3.3);
  - if one fails, the failure names the only lattice lead left.

  It is also the cheapest valid discriminator in this note, because it needs no lattice reduction. L4-A and L4-D need the support facts P2 that L4-B produces before their symmetric-lattice dimensions mean anything.
- **L4-A fills in the measured rows the brief assumed already exist.** Information is moderate, since the predictions are fairly confident. Cost is medium for m = 4.
- **L4-D asks the right §2.4 question: cost exponent, not threshold.** It is cheap. A positive result would matter, but its prior is low.
- **L4-C is the only test aimed at non-lattice structure.** It is very cheap. A null result strengthens the closure. A positive result would be the only new lead.

5. Killed on paper

Every entry is `[derived]` unless marked otherwise. "Reach" means the JM asymptotic δ in factor-base-size units.

#	idea examined	mechanism hoped for	why it fails	effect on δ
K1	Möbius-transformed or translated x-intervals; any other degree-2 coordinate	a better-conditioned box	F3: every degree-2 F_p-function is $M \circ x \circ \tau_S$. Homogenising keeps degree d per variable and the same box	none (≡ JM)
K2	Farey ($x = u/v$), GLV/lattice-coset ($x = u + \lambda v$), SNFS base-M digit bases on Solinas primes ($x = \Sigma a_j M^j$)	"approximate subfield"; Weil restriction over Z	F4: degree-1 substitution, same box volume, same first-order reach. Exact splitting into several equations over Z needs $\lvert S(\ldots)\rvert < M$, i.e. small x_R and small b: that is F6 (xedni)	none
K3	Torsion symmetrisation (FHJRV), including the "D_4-symmetrised S_5 probe"	lower-degree invariants	needs $E(F_p)[2]$ or $[4] \neq 0$, which prime-order curves lack. On even-order curves, quotienting by $\langle T \rangle$ maps the problem onto $E/\langle T \rangle$ with $\lvert F\rvert \times \lvert T\rvert$ at fixed box: a constant	constant
K4	Interval membership polynomial, [illegible] falling-factorial; Bostan–Schost products	evaluate "∃ root in the box" faster than enumeration	the m-dimensional box product costs $B^{m^2/(m+1)}$ (balance $s^{m+1} = B^d$), which is above enumeration's	$\tau = 3.2\delta > 3\delta$ at $m = 4$
K5	[illegible]	[illegible]	[illegible]	≈ 1.28 at (m = 2)

Also checked and already closed in the program note, not re-proposed: the symmetric (e_1, e_2) rewrite of S_3 (§3.2 above shows its asymptote is 1/6, not 1/9, which changes nothing); multiplicative-subgroup bases; global lifting / xedni; embedding into an extension; lattice-assisted MITM; low-degree-map factor bases.

6. Relation to the other active sessions

- **Prime-field fall degree / Gröbner** (`research/prime-fall-degree-20261027`, `research/fall-degree-bounds-20261008`, `codex/fall-degree-*`).
  - Nothing here proposes a Gröbner run.
  - K12 hands the Boolean-variable encoding to them.
  - If their solving degrees for S_4/S_5 with interval constraints come out sub-enumerative, that lies outside F1–F4, which are lattice-specific, and would supersede this note's closure for that solver class.
  - I could not read their branches (§2).
- **Residual large-prime IC** (`research/ic_prime_residual_lp_20261026`, `lead/ic-residual-lp-20261028`). Large-prime variants change constants in the relation phase. They do not change the §3.1 exponent condition, because the oracle's τ(δ) is unaffected. Not duplicated.
- **k-sum table** (`research/prime-ksum-table-20261028`). Generic, out of scope by the brief. Figure 1B's black line is its exponent.

7. Honest accounting (inventor protocol §5)

```
honest_accounting:
  objects:
    - "tracked object: the decomposition tuple (x_1..x_m) as a small point on
V(S_{m+1}(.;z)); operation set: none beyond the ideal (S)"
    - "representations examined: K1-K16 (coordinate changes, divisor/line/orbit/
bit bases, lifts, covers)"
  lossy_projection_test: >
    No new tracked object is proposed. Of the representations examined, K1, K2
and K11 are lossless
    changes of coordinates (they fail the test). K6-K9 are lossy but raise the
degree. Index calculus
    stays Class I partial-action (IDEA-20260806-c5d183). None of K1-K16 escapes
KN-OPEN-020: all are
    explicit low-degree descriptions.
  depth_of_verified_structure: "paper-level (derived) only; nothing measured this
session"
  dominated_by: "n/a (no result claimed). Rows checked: Pollard rho sqrt(pi n/4);
preprocessing rho (S*T^2 ~ n, Probe A); generic IC (m-1)-enumeration; JM
asymptotic lattice; McGuire-Mueller O(p log^2 p) (ePrint 2017/1262); naive Semaev
IC (KR-IC-5931ee)"
  sota_delta: >
    No attack. Corrected baseline: at feasible dimension (<= 5e3) the S5 reach is
about 1/313
```

```
    against 1/5 needed (about 63x), and the S4 reach is 1/35.5 against 1/4
(8.9x). Asymptotic
    gaps are 16x and 6x.
  closures:
    - "lattice oracles on coordinate factor bases, prime order: F1+F2+F3+F4 ->
shift-set LP; status unverified until L4-B measures P1-P3"
    - "many-sample / overdetermined lattice imports: F5 (no planted solution, no
free equations)"
    - "exact-vanishing tricks (small coefficients, digit bases, quadratic
orders): F6 (global lifting, xedni)"
    - "K1-K16 individually, each with its mechanism in section 5"
  open_directions:
    - "a non-lattice small-root algorithm for one structured congruence at box
volume p^0.7-0.8, which would need a named external ingredient; none found
(searches below)"
    - "factor bases whose membership is implicit, high-degree or target-dependent
(the KN-OPEN-020 escape), with a polynomial-time membership test and no
coordinate-box description"
    - "transfer (program note section 4.3) and Groebner/fall degree (other
sessions)"
    - "if L4-C shows over-dispersion: a determinant-method oracle aimed at the
observed structure"
```

7a. L4-B step 1, run afterwards (measured, 2026-10-28)

The parent session ran L4-B's first step with PARI/GP (`l4b_step1.gp`, output `l4b_step1.out`). Setup: the frontier campaign's 40-bit `a = -3` curve `icv1-fp40-tm924137-0ecaaeb9` ($p = 2^{40} - 213$, $b = 56$); 20 seeded random `z`; S_4 = `Res_X(S3(x1,x2,X), S3(x3,z,X))`. It takes under a second.

- **[measured] P2 holds on every sample.** For all 20 values of `z`, all 8 vertex coefficients of the box $[0,4]^3$ are nonzero, so the Newton polytope of $S_4(\cdot; z)$ is the full box. No sparser support exists for a lattice to exploit at `m = 3`.
- **[measured] Irreducible over `F_p` on every sample.** For all 20 values of `z`, a restriction to a random line has full degree 12 and is irreducible over `F_p`.
- **[not tested] Absolute irreducibility.** Line restrictions cannot decide it. A degree-12 restriction that is irreducible over `F_p` splits over `F_{p^k}` into `gcd(k, 12)` factors whatever S_4 does. A first version of the script that "tested" `F_{p^2}` was wrong for this reason and was corrected. Absolute irreducibility of summation polynomials is [recalled] a known result (Diem); it was not re-read here.
- **P3 not run.** The automated-Coppersmith linear program of Ding et al. (ePrint 2026/1027) has no implementation in this repository.
- **[measured] `m = 4` (S_5), the arity where beating rho is possible.** `l4b_step1_s5.gp` (`l4b_step1_s5.out`): S_5 = `Res_X(S3(x1,x2,X), S4(X,x3,x4))` on the same curve, 20 seeded `z`.

- All 16 vertex coefficients of $[0,8]^4$ are nonzero for all 20 values, so P2 holds.
- A restriction to a random line has full degree 32 and is irreducible over F_p for all 20 values.

Reading. The failure of P2 was the only lattice lead L4-B could open at $m = 3$ or $m = 4$, and P2 holds at both. P1 holds over F_p. Only P3, the linear program, is left open. Under L4-B's claim, every lattice oracle on a coordinate factor base is then capped at $1/(m(m+1)2^{m-2})$: $1/24$ at $m = 3$ and $1/80$ at $m = 4$, against the $1/4$ milestone and the $1/5$ goal. **Lead 4 is closed for lattice methods on coordinate factor bases**, unless P3 returns a better shift set than Jochemsz–May. What remains is the non-lattice space of §7, where this note has no concrete candidate.

7b. Lead-4 code and the follow-up probes (measured, 2026-10-28)

Five pieces of native Rust, all on main's crate and with no off-main dependency:

- `src/cryptanalysis/semaev_prime_poly.rs`: dense $S_3 \ldots S_6$ over F_p by the GP recursion, plus the structure checks;
- `examples/semaev_prime_structure.rs`: L4-B step 1, natively;
- `examples/semaev_gap_census.rs`: L4-C;
- `examples/semaev_lattice_reach.rs`: L4-A;
- `examples/semaev_shift_search.rs`: the P3 fallback.

Each probe has a hashed `PROTOCOL.md`, registered before its measured run (`REGISTRATION.sha256`), and the hashes of the binary and source it ran (`BINARY.sha256`). Raw outputs are under `*/data/`. Derived tables and figures come from `*/derive.mjs`, which reads only those outputs.

After the runs, CI's clippy (`-D warnings`) required five lint-only edits to `semaev_prime_poly.rs` and two to `examples/semaev_gap_census.rs`: `is_multiple_of`, `div_ceil`, `repeat_n`, and $2 \cdot d < \text{len}$ for $2 \cdot d \le \text{len} - 1$. Each edit computes the same values, and the 6 unit tests still pass. The source hashes in the probes' `BINARY.sha256` refer to the sources as run, at commit f91405e.

7b.1 The GP checks, ported (L4-B step 1, native)

`semaev_prime_poly` builds `S_{m+1}(x_1…x_m; z)` as `Res_X(S_3(x_1,x_2,X), S_m(X,x_3,…; z))`, with formal-degree Sylvester resultants and tensor interpolation.

- **[measured] The values match PARI/GP exactly.** The unit test `values_match_pari` checks S_3, S_4 and S_5 at three points each on the 40-bit curve, against `polresultant` output.
- **[measured] S_4 and S_5 vanish on real sums.** Tested at $x(P_1+P_2+P_3)$ and $x(P_1+\ldots+P_4)$.
- **Irreducibility over F_p.** It is decided by intersecting factor-degree patterns of line restrictions (distinct-degree factorisation). A factor of total degree a forces a to be a subset sum on every full-degree line. This replaces the GP script's "one irreducible line" test. That

test needs about `n` lines for a degree-`n` restriction, and in a first native run it missed for S_5 (degree 32) within 60 lines.

- **[measured, `l4b/native_step1.jsonl`]** On the same curve and the same count (20 seeded `z` per arity): S_4 and S_5 have the full box (8/8 and 16/16 vertices nonzero on 20/20 samples). They are irreducible over `F_p` on 20/20, settled within at most 8 lines for S_4 and 6 for S_5.
  - New: the polynomials are **completely dense**. All 125 coefficients of $S_4(\cdot; z)$ on $[0,4]^3$ and all 6561 of $S_5(\cdot; z)$ on $[0,8]^4$ are nonzero on every sample.
  - It takes 0.1 s (S_4) and 14 s (S_5) for all 20 values of `z`.
- **[derived] Simplex vertices.** For the symmetrised form `g(e)`, the vertex coefficient of `e_i^D` equals the box coefficient of `Π_{j≤i} x_j^d`. That is the only `x`-monomial of that multidegree among `e^β` with `|β| ≤ D`. So box vertices nonzero implies simplex vertices nonzero, which is the premise §7b.4 needs.

7b.2 L4-C: is there structure in the gap? (census, `l4c/`)

The protocol changed the README record in three ways before any data: - an exact census over all `r − 1` targets replaces 10^5 samples; - the counting unit is fixed: signed multisets summing to `R`; - null 2 is dropped, because its expected root count is below 10^{-2} per cell.

A 20-bit smoke run, before registration, showed that a per-target Poisson null is wrong for a finite factor base. Its triple sums collide only through short additive relations, which come in clumps. So the registered comparison is against null 1 (five random factor bases of the same size), with the primary statistic `triple_unshared`: 6-term relations among the factor base.

- **Panel.** Six prime-order curves (`a = −3` and `a = 0` at 26, 28 and 30 bits), with $\delta \in \{1/24, 1/12, 1/8, 1/6, 1/5, 1/4, 0.3\}$. That is 41 cells (one skipped for memory, as registered), 246 censuses.
- **[measured] Null result.** In all 41 cells, the small-x values of `triple_unshared`, `pair_collisions` and `ge2` lie inside the registered band (null-1 range ± 3σ). No (shape, δ) is outside on the same side at two sizes. The falsifier did not fire.
- **[measured] δ ≤ 1/5 is uninformative.** Every count is zero, for the small-x and the null bases alike, as the protocol's limits anticipated. The informative cells are $\delta = 1/4$ and 0.3 (Figure 2).
- **Reading.** Up to the toy sizes tested, small-abscissa factor bases have no more short additive relations than random factor bases of the same size. There is no gap structure for a non-lattice oracle to target. This supports the Poisson-yield assumption in §2.4 and does not reopen lead 4.




Figure 2: Figure 2. L4-C: 6-term relations among the factor base, small-x vs five null-1 bases (exact census over all targets; band = registered null range ± 3σ). Source: l4c/analysis/l4c.svg.

7b.3 L4-A: measured S_4 lattice reach (`l4a/`)

Symmetric JM lattices for S_4: - $g(e_1, e_2, e_3)$ is found by a 35-unknown linear solve, checked on 6 extra points; - the lattice is triangular in (β_3, β), with g monic in e_3^4; - reduction uses main's `lattice::lll_reduce`.

A timing pilot fixed the scope before registration: - dim 35 takes about 20 ms; - dim 165 takes 116 s, about 200 s under load; - no dim-455 reduction finished in 5 h 10 min (`l4a/pilot/PILOT.md`).

So **$\ell \geq 3$ at m = 3, and all of m = 4, stay `[derived]` only.**

Measured cells: - **$\ell = 1$ (dim 35):** 32, 40 and 48 bits; B $\in$ {1, 2, 3, 4, 6, 8, 12, 16}; 8 planted and 8 unplanted targets per cell. - **$\ell = 2$ (dim 165):** 32 and 40 bits; B $\in$ {1, 2, 3, 4, 6}; 4 planted and 4 unplanted per cell.

[measured] v1: the registered falsifier fired, on a planting defect. At $\ell = 2$ on 40 bits, B = 4 succeeded on 3 of 4 targets and B = 3 on 2 of 4, while the §3.2 bound is B $\approx$ 1.7. - PARI/GP shows that **all 8 distinct successful planted targets** in v1, across every cell, satisfy $S_4(x; z) = 0$ over $\mathbb{Z}$ (`l4a/succ.gp`, `l4a/succ.out`). - They have $z \in \{1, 2, 3, 25\}$, and most are trivial $P - P + Q = Q$ decompositions. - Cause: small x's repeat, and the planting rule accepted any root. Any lattice that contains S_4 finds such roots automatically, which is F6 global lifting, not modular reach. - The v1 data, protocol and fired falsifier are kept (`l4a/data-v1/`, Figure 3b). `AMENDMENT-1.md`, hashed before the rerun, redraws planted roots that vanish modulo the independent prime $2^{61} - 1$ and changes nothing else.

[measured] v2 (amendment 1): no planted success in any cell.

bits	ℓ	dim	planted success, all B	unplanted	measured reach	§3.2 bound B_ℓ	median LLL time
32	1	35	0/64	0/64	$B < 2$	1.11	8–19 ms
40	1	35	0/56 ($B = 1$: no non-global root)	0/64	$B < 2$	1.14	9–22 ms
48	1	35	0/64	0/64	$B < 2$	1.17	10–24 ms
32	2	165	0/20	0/20	$B < 2$	1.50	46–131 s
40	2	165	0/16 ($B = 1$: no non-global root)	0/20	$B < 2$	1.66	66–178 s

Every reduction costs far more than the $(B + 1)^2$ enumeration it would replace: 46–178 s against at most 49 candidates at $\ell = 2$.

Reading. The measured $m = 3$ reach agrees with the §3.2 determinant bounds. They predict $B < 2$ at both levels and all three sizes, and v2 finds no success at $B \geq 2$. There is no slack beyond the bound, unlike the up-to-2.5× slack Probe B saw at $m = 2$. No falsifier fired in v2. The $m = 3$ rows at $\ell = 1$ and 2 are now `[measured]`; $\ell \geq 3$ and all of $m = 4$ remain `[derived]`. The v1 episode is a lesson for any future planted-root test: exclude global relations, or a lattice "succeeds" on them by construction.




Figure 3: Figure 3a. L4-A v2: planted-root success of symmetric JM lattices for S_4 against the root bound B, with global relations excluded (zero in every cell). Source: l4a/analysis/l4a.svg.




Figure 4: Figure 3b. L4-A v1 (superseded): the apparent successes, every one a global relation $S_4(x; z) = 0$ over $\mathbb{Z}$. Source: l4a/analysis/l4a-v1.svg.

7b.4 P3 fallback: our search over triangular shift lattices (`p3/`)

What this is not. P3 names the LP of Ding et al. (ePrint 2026/1027). Its PDF is behind a Cloudflare browser challenge this session does not pass, so that LP is **not implemented here**.

What was run instead. An exact count over an explicit family of triangular shift lattices: - every leading vertex of the Newton polytope; - monomial sets made of rectangles with extended caps, cut by `|β| ≤ s` or by one `{0,1,2}` half-space; - `k(β)` maximal subject to the shifts staying in the set.

The score is the determinant bound `δ = Σk / Σ⟨w,β⟩`, without dimension factors. The code reproduces every §3.2 JM value exactly (`jm_set_delta = jm_formula` in all rows). That covers the m = 4 symmetric values: $1.26 \cdot 10^{-4}$, 1/313 and 1/184.

shape	m	best δ found (dim)	best δ / δ_JM at equal dimension	asymptote	exceeded
simplex (e)	3	0.0326 (3642)	1.078 at dim 56	1/24 = 0.0417	no
box (x)	3	0.0306 (15625)	1.010 at dim 1331	0.0417	no
simplex (e)	4	0.0061 (64065)	1.032 at dim 495	1/80 = 0.0125	no
box (x)	4	0.0048 (390625)	1.002 at dim 83521	0.0125	no

- **[derived by computation] No lattice in the family exceeds the asymptote.** At equal dimension the best gain over the JM curve is **≤ 8 %**, at small dimension only. Above a few hundred it is ≤ 3 %. These are extended-strategy shifts, not a new exponent (Figure 4).
- **Reading.** This is consistent with F1–F4 and with L4-B's conditional closure. It is a finite family, not the LP, so P3 remains formally **open**. The ePrint's formula, read and run, is the only thing that could still change it.




Figure 5: Figure 4. P3 fallback: δ against dimension for our shift-lattice family (lines) and the JM levels (squares), with the asymptotes. Source: p3/analysis/p3.svg.

7b.5 Where lead 4 stands

premise / probe	status
P1, irreducible over F_p	measured: GP and native, 20/20 at m = 3 and 4
P1, absolutely irreducible	not decidable by line restrictions; recalled (Diem)
P2, full box	measured: GP and native, dense, 20/20
P3, automated-Coppersmith LP	open; our finite search finds nothing past the JM asymptote
L4-A, measured reach	measured at m = 3, $\ell \le 2$: no success at $B \ge 2$, matching §3.2 (v1 false positive: global relations, amended)
L4-C, gap structure	measured: null, 41/41 cells inside the band
L4-D	not run

Lead 4 stays **closed for lattice oracles on coordinate factor bases**, conditional only on P3. No measured cost changed, so this is not an improvement under the success criterion, and no dashboard point is added (AGENTS §7a).

8. Sources

Each entry is marked with how far it was read.

- [internal] RESEARCH_PRIME_FIELD_BREAKTHROUGH_PROGRAM.md (2026-10-08): §2.1–2.4 cost model and JM table; Probe B/C tables; §4.6 closures.
- [internal] RESEARCH_IC_LEADS_AND_CONNECTIONS_20261010.md §8–§10 (worktree copy).
- [kb] crypto-autoresearcher frontier rows KR-IC-5931ee and KR-IC-5a43e5; KN-LIT-d8e1b9 (Yokoyama, Yasuda, Takahashi, Kogure, *J. Math. Cryptol.* 14 (2020) 460–485, doi:10.1515/jmc-2019-0029).
- [retrieved, 2-page abstract read] G. McGuire, D. Mueller, "A few more index calculus algorithms for the ECDLP", ePrint 2017/1262 (Caen 2018 abstract): O(p log² p), Gröbner-free. Its reference list confirms Amadori–Pintore–Sala ePrint 2017/609 and Faugère–Huot–Joux–Renault–Vitse, EUROCRYPT 2014, LNCS 8441, 40–57. https://nitaj.users.lmno.cnrs.fr/Caen2018/DanielaMuellerCaen2018.pdf
- [retrieved, abstract] J. Hu, "Implementation analysis of index calculus method on elliptic curves over prime finite fields", ePrint 2024/1923. https://eprint.iacr.org/2024/1923
- [retrieved, abstract] Y. Aono, M. Agrawal, T. Satoh, O. Watanabe, "On the optimality of lattices for the Coppersmith technique", ePrint 2012/108; ACISP 2012; AAECC 29(2) 2018. https://eprint.iacr.org/2012/108
- [retrieved, abstract] J. Xu, S. Sarkar, H. Wang, L. Hu, "Improving bounds on elliptic curve hidden number problem for ECDH key exchange", Asiacrypt 2022, ePrint 2022/1239. https://eprint.iacr.org/2022/1239
- [retrieved, search abstract] K. Meers, J. Nowakowski, automated Coppersmith, Asiacrypt 2023, ePrint 2023/1409. https://eprint.iacr.org/2023/1409
- [retrieved, abstract] Z. Ding, Z. Dai, B. Wu, Y. Zhang, K. Zhang, "Computing asymptotic bounds for the automated Coppersmith method via linear programming", ePrint 2026/1027. https://eprint.iacr.org/2026/1027
- [retrieved, search abstract] M.-C. Chang, J. Cilleruelo, M. Garaev, J. Hernández, I. Shparlinski, A. Zumalacárregui, "Points on curves in small boxes and applications", arXiv:1111.1543. https://arxiv.org/abs/1111.1543
- [recalled] I. Semaev, ePrint 2004/031. E. Jochemsz, A. May, Asiacrypt 2006. D. Coppersmith, J. Cryptology 1997. Bostan–Gaudry–Schost, SIAM J. Comput. 2007. Ostrowski's theorem on Newton polytopes of products. Jacobson–Koblitz–Silverman–Stein–Teske, DCC 2000 (xedni). Petit–Kosters–Messeng, PKC 2016. Gaudry–Thériault–Thomé–Diem (genus-g double large prime).
- **Searches that found nothing beyond the above** (2026-10-28):
  - "summation polynomials prime field … Coppersmith small roots lattice index calculus";
  - "ePrint 2023 2024 2025 index calculus elliptic curves prime field summation polynomial";
  - "elliptic curve hidden number problem x-coordinate Coppersmith 2020–2023".

  None of them turned up a small-root method for summation polynomials past the JM reach, or any sub-$\sqrt{p}$ prime-field algebraic ECDLP result. Hu 2024 states that no subexponential method is known.

9. Files and rendering

- `README.md`: this report.
- `delta_arity_map.svg`: Figure 1, hand-written and editable. Every coordinate follows the formula in its header comment.
- `delta_arity_map.png`: the figure rendered with `rsvg-convert -w 2160 delta_arity_map.svg -o delta_arity_map.png`.
- `report.pdf`: rendered from this file with every figure link pointed at its PNG (`pandoc … --pdf-engine=typst`), after §7b was added.
- `l4b/`, `l4c/`, `l4a/`, `p3/`: one directory per §7b probe. Each holds `PROTOCOL.md` (with `AMENDMENT-1.md` for L4-A), `REGISTRATION.sha256`, `BINARY.sha256` (binary and source hashes), raw `data/`, `derive.mjs` (a thin Node derivation over the raw data; tooling, not research code) and `analysis/` (tables, editable SVG figures, PNG renders made with `rsvg-convert -w 2160`).
- The native code lives outside this directory: `src/cryptanalysis/semaev_prime_poly.rs` and the four `examples/semaev_*.rs` programs named in §7b.