Independent weighted doubling review

No blocking discrepancy was found at the stated field-value scope. The Rust dbl nested operations match the Lean expression, including output z unchanged and output w=$2y$.

Independent derivation. Division of $f(X)-(mX+y-mx)^2$ by $(X-x)^2$ leaves constant residual $f(x)-y^2$ and linear residual $f'(x)-2ym$. The explicit cubic quotient has coefficients $1, 2x, 3x^2+a_3, 4x^3+2a_3x+a_2-m^2$. Valid point input and $m=f'(x)/(2y)$ cancel both residuals. A nonzero derivative is unnecessary. Nonzero weights and y, and characteristic different from2, ensure the denominators and output weights are nonzero.

Dispatch scope. The self-add guard applies to identical weighted representations. At nonzero y the earlier inverse guard is excluded; at zero y it takes precedence. No converse for differently weighted inputs is claimed. The producer $f=X^5$ example is polynomial algebra, not smooth-curve evidence.

Replay. All manifest hashes match. Private module, producer controls and separate axiom readback exit0. Only propext, Classical.choice and Quot.sound appear. Reviewer controls apply the full theorem with all coefficients nonzero, nonunit weights and normalized point $(0,1)$ on $X^5+4X^3+3X^2+2X+1$: slope and intercept are1. Changing the constant to2 makes the required divisibility fail, confirming the point premise is material.

Reviewer negative-control attempts used unavailable evaluation APIs and then needed explicit coefficient simplification. Failures are preserved; taking constant coefficients of the witness equation completed the corrected control. Producer sources were unchanged.

Boundary. Monic quintic with no fourth-degree term; field polynomial validity and self dispatch. No UInt64, counter/compiler, divisor-class equality or whole-walk claim. No scientific trial or cost measurement. Exact receipts and hashes are in review.json and custody.json.