K-233 derived row of the admissible-degree inequality (L8)

Class: derived calculation; no S, no measured cost. Calculation: `examples/k233_admissible.rs`; formulas as in `dreg_score slope` (derived table). Curve sect233k1, r = 0x8000…bdf (log2 r = 231), cofactor 4, log2 T_rho = 110.894.

Formulas. R = 2^l/(2n), A = max(1, m! 2^(n-ml)), B = T_rho/(R A); admissible D is the largest D in 2..40 with w log2 C(N, <=D) <= log2 B, for N = ml + (m-2)n (chained) and N = ml, w in {1, 2.2}.

Stable bases (computed). ord_233(2) = 29: nine 2-cyclotomic cosets, one of size 1 and eight of size 29. Stable dimensions 29a + c, a in 0..8, c in 0..1.

m	l	ml	log2 B	D chained w=1	D chained w=2.2	D elim w=1	D elim w=2.2	D=4 fits
3	58	174	0.2	none	none	none	none	no
3	59	177	2.2	none	none	none	none	no
3	87	261	32.8	4	none	4	none	yes
3	88	264	31.8	4	none	4	none	yes
3	116	348	3.8	none	none	none	none	no
4	58	232	56.2	7	2	9	3	yes
4	59	236	59.2	7	3	10	3	yes
4	87	348	32.8	3	none	4	none	yes
5	29	145	−4.1	none	none	none	none	no
5	30	150	−0.1	none	none	none	none	no
5	58	290	61.8	7	3	10	3	yes
5	59	295	60.8	7	3	10	3	yes
6	29	174	22.3	2	none	3	none	no
6	30	180	27.3	2	none	4	none	yes
6	58	348	61.8	7	2	9	3	yes




Figure 1: Admissible D against stable l for each m (derived, from rows.jsonl). Dashed: D = 4. Hollow marker at 1: none.

Conclusion (derived). First D >= 4 by ml: chained w=1 at m=4, l=58 (ml=232, D=7); N=ml, w=1 at m=6, l=30 (ml=180, D=4); w=2.2: none (maximum 3). Measured closure degree is 4 at ml <= 18 and above 4 at ml = 21; every row here has ml >= 145, so the closure degree there is unmeasured and D = 4 fitting is not sufficient (needs closure <= 7 to 10 at w=1, <= 3 at w=2.2). The inequality counts the relation phase only. Distance to rho remains a distance, not a measurement.