1. Twist symmetry in quartic extension index calculus

22 October 2026. Full source-linked record: `QUARTIC_TWIST24_REPORT.md`.

1.1. Question and protocol

We tested a transported Frobenius on a quadratic twist over F_{p^4}, combined it with the cubic $j = 0$ automorphism, and measured an unchanged physical base, the union of two invariant lines, the full invariant two-dimensional abscissa space, tagged pair indexes, and dense same-residual rank extraction. The line panel has seven positive toy curves and three seeds; the full 2D target solves use $p = 13, 37$. Each run has one public target $[271]G$. The signed four-factor oracle, matrix encodings, and 24-fold rho reference are native Rust. The frozen protocol and raw records are in the same study directory.

The construction uses $F = \frac{F_p[t]}{t^4-d}$ with d the least nonsquare, $E : y^2 = x^3 + b_0 t^3$, $h = t^{\frac{1-p}{2}}$, $\omega^3 = 1$, and $\psi(x, y) = (h^2 x^p, h^3 y^p)$, $\iota(x, y) = (\omega x, y)$. The norm identity $h^{1+p+p^2+p^3} = -1$ gives $\psi^4 = [-1]$. The two maps commute. On the certified prime-order group their checked eigenvalues have orders eight and three, yielding order 24 on signed points and 12 on nonzero abscissae. Thus a stable base with B abscissae has $\frac{B}{12}$ unknown logs, compared with $\frac{B}{3}$ for the cubic automorphism alone.


For the four lines $t^j F_p$, ψ multiplies abscissae by z^{j-1}, $z = t^{p-1}$. Lines 1 and 3 cannot contain a nonzero odd-order point: ψ or ψ^2 would fix its abscissa and hence act as either plus one or minus one, contradicting $\psi^4 = [-1]$. The complete sweep found only lines 0 and 2 usable, with line 0/2 counts shown below.

p	r	line 0	line 2	24-fold columns 0→0+2
13	28,753	0	0	none
37	1,873,633	36	12	3→4
73	28,408,033	48	48	4→8
97	88,538,353	36	36	3→6
157	607,561,153	108	84	9→16
229	2,750,120,881	96	96	8→16
241	3,373,515,841	96	96	8→16
337	12,897,718,513	180	180	15→30

1.2. Fixed physical base: what the fold changes

The same signed points and residual/decomposition stream feed flat, cubic-only and combined matrices. All 63 IC solves and 1,008 rho walks recovered 271. Independent native replay reconstructs 1,821 selected relations, all row encodings and matrix spans. The full A/A rerun matched every non-timing field.

p	cubic/twist queries	cubic/twist row ops	cubic/twist M	twist/rho24 M
37	3.27	10.78	0.998	55.5
73	2.57	11.24	1.49	112.9
97	2.96	12.51	2.66	192.9
157	3.70	11.03	2.49	201.9
229	3.48	14.79	3.18	404.6
241	3.81	13.30	3.80	376.8
337	3.33	14.45	2.99	402.1




M is recorded base-field multiplications on each path with its relevant model, input, eigen, base, map, table, stream and verification work. Row operations are a separate unit. The 64-query batch floor erases the field-multiplication improvement at $p = 37$. The rho comparison is same-target and includes its 24-fold endomorphism; the large gap is a partial-cost diagnostic, not an admitted speed ratio.

Applying an endomorphism to a found relation transforms the target and multiplies both the folded row and its right side by the eigenvalue. The audit verifies this covariance on every selected relation. Orbit images are not independent rows. Therefore the unchanged physical base gains relation **utility**, not raw hit probability.

1.3. Two invariant lines: actual relation yield

For each curve and seed, L_0 and $L_0 \cup L_2$ receive the exact same target stream, with identical L_0 points and signs. The independent native oracle recomputed every one of 69,045 query hit counts using a fully signed two-pair hash, re-added 5,298 selected group relations, and checked all 126 matrix solves. The producer recorded 7,275 decompositions. Both full runs matched on every non-timing field. Medians are over three paired, separate one-target runs.

p	union/single hit yield	single/union queries	single/union row ops	single/union counted M
37	2.04	1.67	0.61	0.57
73	8.82	4.82	0.30	0.26
97	14.60	6.73	0.30	1.01
157	5.68	3.72	0.34	0.63
229	17.55	11.07	0.33	2.48
241	14.85	7.08	0.23	1.49
337	11.28	5.08	0.26	1.24



The union improves same-target hit yield on every curve and lowers query count, but doubles or nearly doubles matrix columns and increases row work. Counted field multiplications improve in all three seeds at $p = 229, 241$; they worsen at $p = 37, 73, 157$ and straddle one at $p = 97, 337$. The figure keeps the row-operation and field-multiplication units distinct. This is a relation-yield result with a cost tradeoff, not a uniform whole-method win.

1.4. Interpretation, limits, and provenance

The combination of Frobenius-invariant factor bases and matrix-orbit savings is prior work (Galbraith–Granger–Merz–Petit). GLS twist endomorphisms are prior work (Galbraith–Lin–Scott). This study makes a specific canonical quartic construction and paired mixed-line measurement explicit. An earlier repository E18 run already used the cubic $j = 0$ fold; this panel provides the same-base cubic control. The earlier repository E2b GLS experiment implemented an S_3 Weil-descent resultant oracle. The main quartic relation panel uses exact four-sum meet-in-the-middle; a separate six-query pilot runs the native S_5 solver.

The experiment is L0 on a shared Intel Xeon 6975P-C host. The main panel does not emit IC-stream group additions, hash/table operations or a calibrated group-addition-equivalent band. Its field-multiplication path projection excludes work done solely for the other two matrix arms. Whole-process wall times were not isolated and are not used for a speed claim. A single-arm sealed run and full polynomial relation stream compared with msolve are needed before an end-to-end cost or asymptotic claim. The four one-dimensional Frobenius eigenspaces were exhausted. A separate full census tested the two-dimensional space $F_p + t^2 F_p$; other invariant subspaces and curve models remain open.

The field/twist proof above is handwritten and computationally checked. `QuarticTwistSymmetry.lean` proves conditional consequences (fourth-to-eighth iterate, composed eigenvalue, row covariance) with an audited Lean kernel. It does not formalize the concrete finite field or twist isomorphism. The full proof map and source links are in `QUARTIC_TWIST24_REPORT.md`.


Evidence: `quartic_twist24_full_v2.ndjson`, `quartic_twist24_native_audit_v2.json`, `quartic_twist_mixed_v1.ndjson`, `quartic_twist_mixed_native_audit_v1.json`, `quartic_twist_mixed_summary_v1.json`, the frozen protocol, and `OPEN_PROBLEMS.md`. Write-once run reports with graphs and PDFs are under `reports/`. All artifacts are in this study directory.

1.5. Polynomial pilot and larger invariant space


The registered native S_5 pilot ran one successful and one unsuccessful line-0 residual at each $p = 37, 73, 97$. All six polynomial outputs exactly matched the signed group oracle; every returned quadruple passed group addition. Each curve's symmetrisation precomputation used 1.226 billion F_p multiplications. The same degree-29 Macaulay matrix shape appeared on all six solves: 41,780 rows and 40,920 columns. The plot shows counted solve multiplications; shared-host L0 solve times were 49–58 seconds and are not isolated speed evidence.

p	hit query / $10^{12}M$	miss query / $10^{12}M$	oracle agreement
37	0 / 1.158	1 / 1.159	2/2
73	25 / 1.203	0 / 1.189	2/2
97	14 / 1.211	0 / 1.197	2/2

The two-dimensional invariant abscissa space $F_p + t^2 F_p$ was exhaustively scanned on all eight curves. Independent norm-character and point-orbit replay checked all 265,008 candidate abscissae. Every nonempty abscissa orbit has size 12. The formerly empty-line $p = 13$ curve has 72 liftable interior abscissae and six folded columns. At $p = 337$, the direct exact-oracle pair construction would generate 3.178 billion signed entries; this count does not measure allocated memory.



On the $p = 13$ two-dimensional base, a separately frozen one-target exact-oracle IC pilot processed seven queries, found 4,041 decompositions, selected seven verified relations, reached rank seven, and recovered 271. Independent replay recomputed all query counts and rows; the full A/A rerun matched every non-timing field. This shows that the larger invariant base revives the toy empty-line case. It is not a polynomial-system or rho speed result.




The complete measurements, per-curve S5 records, two-dimensional census, native audits, p=13 run, figures and write-once run reports are linked from `QUARTIC_TWIST24_REPORT.md`.

1.6. Twist symmetry in the relation-finder pair table

On the same p=13 two-dimensional base, the signed two-factor table has 5,112 entries modulo global sign. The 12 abscissa actions partition it into 426 orbits, each of size 12. The direct quotient stores one pair representative per orbit under 350 canonical sum-abscissa keys. It reconstructs transformed pairs when queried. On every one of the seven previously recorded residuals, its **complete signed four-factor decomposition set** equals that of the full pair table; all 4,041 decompositions and seven selected IC relations remain available. A full A/A rerun matched every non-timing field.

table arm	stored pair records	setup F_pM	seven-query F_pM	setup group ops
full	5,112	495,864	7,029,782	5,112
orbit quotient	426	544,746	96,036,422	426




This exact symmetry quotient reduces stored pair **records** by 12 times and setup group additions by 12 times. It does not measure allocated bytes; action maps and hash buckets also consume memory. Its canonical lookup uses 13.66 times the query base-field multiplications of the full table, while query group-operation counts are equal. Therefore the tested direct quotient preserves relation yield and trades arithmetic for a smaller representative set. It is not a relation-lookup speed improvement on these targets.

Raw records: `quartic_twist_pair_quotient_p13_v1.json` and its A/A copy. Source: `examples/quartic_twist_pair_quotient_p13.rs`; the frozen protocol, full run report and exact operation counts are linked in `QUARTIC_TWIST24_REPORT.md`.

1.7. Action-tagged relation-finder index

The ordered, hash-key and direct-posting variants store one pair tuple per 12-action orbit while indexing each actual transformed pair-sum abscissa. They avoid query-time canonicalization. On the same p=13 seven residuals and p=37 three residuals, every variant returned the **exact same complete signed four-factor decomposition sets** as the full table. A separate group addition audited all 5,112 and 388,752 postings. Six non-timing A/A diffs are empty.

p	full / tagged setup F_pM	full / tagged setup group ops	full / tagged query F_pM	full / direct retained requested MB
13	495,864 / 154,922	5,112 / 426	7,029,782 / 7,029,782	1.273 / 0.682
37	37,708,944 / 7,796,605	388,752 / 32,396	226,616,754 / 226,616,754	91.007 / 47.721




Setup group additions fall 12-fold; counted setup F_p multiplications fall 3.20-fold at p=13 and 4.84-fold at p=37. Query F_p multiplication and group-operation counts are exactly equal. For setup plus these registered queries, direct posting uses 1.047-fold and 1.128-fold fewer counted F_p multiplications. The ordered index has the smallest retained requested payload but high binary-search overhead; the hash and direct variants use more bytes and less query CPU. Shared-host L0 timings do not establish a wall-time gain. Retained requested bytes are allocator payloads, not RSS. A complete one-target polynomial index calculus, rho comparison and isolated native-time result remain open.

The twist action alone does not create new relations on a fixed physical base. Exact decomposition sets are identical pointwise; physical-base expansion, as in the mixed-line and 2D experiments, is what raises hit yield. The tagged index changes the **cost** of finding the same relations. Commands, source digests, all raw counters and write-once run PDFs are linked from `QUARTIC_TWIST24_REPORT.md`.

1.8. Full p=37 two-dimensional rank pilot

The registered full $F_p + t^2 F_p$ base has 624 signed physical points and 52 twist-folded factor-log columns. The first 53 seed-zero residuals on one public target $[271]G$ yielded 2,833,126 exact signed decompositions and 53 selected relations. The matrix reached rank 53 and recovered 271; every factor log was checked by group multiplication. A separate point-keyed four-sign pair oracle enumerated all 1,873,633 subgroup logs, recomputed every hit count, checked selected rows and rank, and recovered the same scalar. The non-timing A/A diff is empty.

phase	F_p multiplications	group operations
model and base	2,568,538	0
input, eigenvalues, orbit map	1,898,403	17,726
full pair table	37,708,944	388,752
53-query stream	4,003,923,826	41,277,322
answer checks	2,004,307	18,854




The two-line shared prefix had 43 hits and 89 decompositions over these 53 residuals; the full 2D base had 53 hits and 2,833,126 decompositions. The larger base also raises the factor-log column count and per-query pair work. Total counted path: 4,048,104,018 F_p multiplications, 41,702,654 group operations and 369,021 modular row operations. This is an exact group-oracle pilot, not an S_5 or calibrated rho result.

1.9. Dense rows from one residual

The previous pilot retained one relation per residual. The new registered extractor kept every rank-raising decomposition of the **first** residual and fixed the public generator's factor log to 1. For a fixed residual, all unanchored row differences annihilate the nonzero factor-log vector and have target coordinate zero, so their rank is at most m for $m+1$ unknowns. That ceiling was attained: 6/7 at p=13 and 52/53 at p=37. After removing the known column, the first residual had full rank on both curves. The independent point-keyed oracle verified all first-query decompositions, selected terms, rank and answer; A/A non-timing diffs are empty.

p	queries old → dense	accepted / candidates scanned	old → dense counted F_pM	old → dense group ops
13	7 → 1	6 / 6	8,096,696 → 2,045,670	81,076 → 18,609
37	53 → 1	52 / 115	4,048,104,018 → 119,740,945	41,702,654 → 1,204,356




The p=37 counted one-target field and group paths fall 33.807-fold and 34.627-fold. This combines dense row retention with a public known-log anchor on the twist-folded 2D base; the reduction is not attributable to twist symmetry alone. The exact group oracle remains far from matched 24-fold rho and does not price polynomial relation finding, hash CPU, allocator work, isolated native time or a GAE band. The derivation, certified signed-count polynomial (`IDC1hf0237ddd316cd9d8`), raw records, independent audits, and limitations are in `QUARTIC_TWIST24_REPORT.md`.

For B signed physical factors, a uniform residual has $16\frac{\binom{B}{4}}{r} \approx \left(\frac{2}{3}\right)\frac{B^4}{r}$ expected decompositions. With $m = \frac{B}{12}$ folded columns, the expected count reaches m near $B^3 \approx \frac{r}{8}$; this is an occupancy heuristic, not a rank guarantee. The explicit pair oracle then builds $\Theta(B^2) = \Theta\left(r^{\frac{2}{3}}\right)$ entries with one group addition each, exceeding constant-automorphism rho's $\Theta\left(r^{\frac{1}{2}}\right)$ exponent. A 12-fold orbit quotient changes only the constant. Polynomial and implicit-table relation finders are outside this projection.

1.10. Full-stream tagged-table boundary

On the **old** 53-residual p=37 full-2D trace, direct-posting and full signed-pair tables returned identical complete sets for all 2,833,126 decompositions. Every tagged posting was checked by a separate group addition; A/A non-timing records matched. Query arithmetic was identical: 4,003,562,862 F_p multiplications and 41,273,856 group operations in each arm.

arm	setup F_pM	setup group ops	retained requested bytes
full	37,708,944	388,752	91,007,248
tagged direct	7,796,605	32,396	47,721,464




Setup plus 53 queries cost 4,041,271,806 versus 4,011,359,467 counted F_p multiplications, a 1.00746-fold reduction. The 4.84-fold table-setup gain is diluted by the query stream. Its L0 wall time is not isolated; the tagged arm was slower in that shared-host observation. Dense extraction reduces the number of quadratic-size queries and gives the larger counted gain. Neither result establishes a polynomial or rho speedup.

1.11. Combined tagged index and dense rows in one producer

The registered p=37 combined producer charged model/base/input/eigen/map, a direct-posting tagged pair index, the first seed-zero residual, anchored rank selection, matrix solve and all answer checks in one exact group-oracle path. It used the same 624-point base and 52 folded columns as the earlier full-table dense arm. The tagged and separately charged full reference returned identical complete sets of 53,551 signed four-distinct-factor decompositions. The first 115 candidates supplied anchored rank 52/52 and the target scalar 271. The independent point-keyed auditor enumerated all 1,873,633 subgroup logs, replayed the complete hit count and selected rows, and independently recovered 271; the final-source A/A and original/final-source non-timing diffs are empty.

one-target counted path	full-table dense	tagged dense	full/tagged
F_p multiplications	119,740,945	89,828,606	1.333
group operations	1,204,356	848,000	1.420
table setup F_pM	37,708,944	7,796,605	4.837
table setup group ops	388,752	32,396	12.000



The total deltas of 29,912,339 F_pM and 356,356 group operations equal the table-setup deltas exactly; all other counted phases match. The tagged table retained 47,721,464 requested bytes and the producer peaked 78,834,528 requested bytes above pre-table allocation. Its first-residual stream recorded 777,503 hash lookups and 324,839 candidate postings. The anchored matrix needed 32,453 modular operations; 17,251,447 additional operations checked an unanchored-rank diagnostic, so the shared-host L0 wall observation of 639.113 ms is not an isolated anchored-path time. Relative to the older 53-query full-table pilot, the combined counted ratios are 45.065-fold F_pM and 49.178-fold group operations. This larger difference also includes generic dense-row extraction and a public known-log anchor, not only twist symmetry.

The exact-oracle path still costs about 257 times the earlier same-target 24-fold rho F_pM count and 309 times its group-operation count, as separate cross-run diagnostics. Polynomial S_5 decomposition enumeration, hash/allocator CPU, isolated timing and a matched in-process 0/0.1/1 GAE band remain unmeasured. The full source-linked report, raw records, run/A-A PDFs, audit and protocol are in the study directory.