A finite exceptional-dispatch witness for the genus-2 weighted law

Source-bound review packet

25 October 2026

Input. Over $\mathbf{F}_{17}$, take $C : y^2 = x^5 + x + 2$ and the reduced divisor with $u = x^2 + 6x + 5 = (x - 12)(x - 16)$, $v = 0$. The source-bound checker recomputes that C is smooth and $u \mid f$. The compiled Rust fixture checks validity and independent Cantor addition.

Path	Observed branch	Result
Author public dispatcher	inverse before double	identity
Rust public dispatcher	same_u_inverse	Identity
Independent Cantor	$D + D$	Identity
Author direct doubling	DBL2B0	$(0, 0, 0, 0, 0, 0)$
Rust direct doubling	b_zero_identity	Identity

The direct author output has $Z = W = 0$ and is not a valid weighted divisor. Two normalization denominators would be zero. The author's public dispatcher checks the inverse condition first and therefore does not reach that subroutine on this input. The public paths have zero undefined inverses for this fixture. This is a subroutine edge, not a public-law discrepancy.

Reproduction. The standardized row is `ROWS.csv`; `run.py` compiles the pinned Rust source plus a temporary test module. The initial intentionally failed assertion and corrected passing rerun are retained as raw logs. `check.py` pins the complete author and Rust source SHA-256 hashes, checks dispatch order, and evaluates the author expressions. The exact command and source URL are in `README.md`.

Boundary. This checks one $\mathbf{F}_{17}$ input. The author Magma source was evaluated, not executed in Magma. It establishes no universal formula parity, Picard-group theorem, or new cost measurement.