# Iwasawa leads I-12..I-14

# Iwasawa leads I-12, I-13, I-14 for prime-field ECDLP

Date 2026-11-13. Decision: all three **close** as derivations; no lead survives as stated. No experiment was run and no timing was taken; this note is a derivation with a diagram (`obstructions.svg`). Every result below is a derivation by the producing session and is an unverified assertion [pcc:unverified]. Standard facts are cited to the literature, not re-proved in Lean.

Setting throughout: $E/\mathbb{F}_p$ ordinary, $p > 3$, $n = \#E(\mathbb{F}_p)$ prime, $n \neq p$, $P$ a generator, $Q = [x]P$.

## 0. Where the leads came from

A brainstorm of fourteen Iwasawa-flavoured ideas (session of 2026-11-13) sorted eleven of them against four known obstructions:

- **W1 torsion wall.** On every smooth lift with $p \nmid n$, $E(\mathbb{F}_p)$ is the torsion of $\tilde{E}(\mathbb{Z}_p)$; every homomorphism to $(\mathbb{Q}_p, +)$ kills it (`research/prime_field_leads_20261031/C1_lift_torsion/REPORT.md`, itself a derivation [pcc:unverified]).
- **W2 MOV.** n-adic Galois structure in the constant direction appears first in $\mathbb{F}_{p^k}$, $k$ the embedding degree.
- **W3 lift freedom.** Global lifts of $(P, Q)$ are free choices and lose the relation (xedni).
- **W4 field degree.** $n$-torsion over a number field needs degree of order $n$ or more.

Three ideas were not obviously caught. This note chases them. There was no idea numbered 15.

| Lead | Idea | Closes by |
|---|---|---|
| I-12 | Signature calculus / Poitou–Tate with Kolyvagin (Euler-system) classes | Lemma A + Huang–Raskind equivalence (W4) |
| I-13 | Specialised Ulmer-tower sections as an index-calculus factor base | Lemma B (W3) + Legendre parity |
| I-14 | Function-field Iwasawa: non-constant $\mathbb{Z}_n$-tower over $\mathbb{F}_p(t)$ | Lemma C: no such tower exists (reduces to W2) |

# 1. I-12: signature calculus with Euler-system classes

**Prior art.** Huang and Raskind lift the DLP to a number field and use global (Poitou–Tate) duality; the local Tate pairings of a global class sum to zero over all places. Their abstract states that computing the ramification "signature" is random-polynomial-time equivalent to the DLP, and that the index-calculus analogue does not go through in the elliptic-curve case [pcc:unverified] (arXiv:0710.2363, LMS J. Comput. Math. 2009). For $n \neq p$ and good reduction, the local Kummer image at p is $H^1_ur(Q_p, \tilde{E}[n]) \cong E(F_p)/n = E(F_p)$, so the instance lives in local cohomology at p.

**What the lead added.** Kolyvagin derivative classes (and level-raising variants) are global classes in $H^1(K, \tilde{E}[n])$ with controlled ramification at auxiliary primes $\ell$, built without choosing a free lift of Q. The hope was a supply of global classes with known signatures, cheaper than generic.

**Lemma A (admissible primes are large).** Every auxiliary prime $\ell$ used by an Euler system for $\tilde{E}[n]$ (Kolyvagin primes, Bertolini–Darmon n-admissible primes) has $Frob_\ell$ acting on $\tilde{E}[n]$ with an eigenvalue $\pm 1$ mod n, hence n divides $\#\tilde{E}(F_\ell) = \ell + 1 - a_\ell$ or its twist $\ell + 1 + a_\ell$. Hasse gives both at most $(\sqrt{\ell} + 1)^2$, so $\ell \geq (\sqrt{n} - 1)^2$, of order n.

*Consequence.* A Kolyvagin class at $\ell$ lives over the ring class field $K[\ell]$, whose layer over the Hilbert class field is cyclic of degree $(\ell + 1)/u_K \geq (\sqrt{n} - 1)^2/3$, and the derivative operator $D_\ell = \Sigma\, i \cdot \sigma\hat{}i$ sums $\ell$ conjugate Heegner points. For Kolyvagin primes proper, $n \mid \ell + 1$ forces $\ell \geq 2n - 1$. The work to write down even one class is therefore $\Omega(n)$ field operations, against $O(\sqrt{n})$ group operations for rho. A further cost, not needed for the bound: Heegner points need a modular parametrisation of a lift $\tilde{E}/Q$, and a lift with coefficients below p has conductor up to about $p^3$.

**Verdict.** Closes (W4). Euler systems give no supply of global classes cheaper than the field degree of the n-torsion. Huang–Raskind's equivalence stands, so any sub-$\sqrt{n}$ producer of signatures would itself be a DLP algorithm; Lemma A shows that the Euler-system route is not one.

# 2. I-13: specialised Ulmer sections as a factor base

**Setup.** Ulmer's Legendre curve $y^2 = x(x + 1)(x + t)$ over $K_d = F_p(\mu_d, u)$, $u\hat{}d = t$, $d = p\hat{}f + 1$, has explicit points $P(\zeta u)$, $\zeta \in \mu_d$, generating a subgroup $V_d$ of rank $d - 2$ and finite index in $E(K_d)$ (Ulmer, arXiv:1002.3313; Conceição–Hall–Ulmer, arXiv:1307.4251). Specialising $u \mapsto u_0$ gives d points of the fibre curve $E_\{t_0\}$, $t_0 = u_0\hat{}d$.

**Lemma B (specialisation does not hand over logs).** Let $\Lambda$ be a Mordell–Weil lattice of rank r and $sp: \Lambda \to E_\{t_0\}(F_q)$ specialisation at a good fibre, with image the prime-order group of order n. Then:

1. The relations known for free among the factor base $\{sp(P_i)\}$ are the relations already true in $\Lambda$. For Ulmer's points these are the 2 Galois-trace relations (d points, rank $d - 2$) plus torsion. They give no logs.
2. The logs of the factor base are equivalent to a basis of ker sp, a sublattice of index n. Computing that basis is the DLP in $E_\{t_0\}(F_q)$.
3. Decomposing Q means finding $R \in \Lambda$ with $sp(R) = Q$. That is finding a lattice point in a fixed coset of the unknown index-n sublattice. With no inverse to sp, and sp of small-height points spread over the group, this needs about n trials. It is the xedni pattern (W3) in function-field dress.

**Two incompatibilities with the target, independent of Lemma B.**

- The Legendre family has full rational 2-torsion, so 4 divides #E_{$t_0$}(F_q). No fibre is a prime-order curve, so the family cannot contain the target E.
- The specialised points have x-coordinates in μ_d·$u_0$ with d = p^f + 1. Since μ_d ⊄ F_p for d > 2, they lie in E(F_{p^{2f}}), not E(F_p).

**Verdict.** Closes (W3). A family without forced torsion, such as Ulmer's 2002 curve $y^2 + xy = x^3 - t^d$, removes the parity problem but not Lemma B.

# 3. I-14: non-constant Z_n-towers over F_p(t)

**Lemma C (the only Z_ℓ-extension is constant).** Let K be a global function field over F_q and ℓ ≠ p. Then K has exactly one Z_ℓ-extension, the constant one K·F_{$q^{\{\ell^\infty\}}$}.

*Proof sketch.* By class field theory, the Galois group G_S of the maximal abelian pro-ℓ extension unramified outside a finite set S is the pro-ℓ completion of the idele class group modulo the unit groups O_v^* for v ∉ S. For v ∈ S, O_v^* = F_v^* × (1 + m_v), and 1 + m_v is pro-p, so its ℓ-part is finite. $\mathrm{Pic}^0(K)$ is finite. So G_S ≅ Z_ℓ × T with T finite, the Z_ℓ direction given by degree. Any surjection G_S → Z_ℓ kills T, so it factors through G_S/T ≅ Z_ℓ. Every Z_ℓ-quotient is the degree quotient, which is the constant extension. Every Z_ℓ-extension is ramified at only finitely many places, so this covers all of them.

*Consequence for ℓ = n.* Any Z_n-tower over the base of an elliptic surface $\mathcal{E} \to \mathbb{P}^1$ is the constant tower F_{$p^{\{n}$k}}(t). Specialised to the fibre E, Selmer growth along it is governed by the action of Frobenius on E[n^∞]. It shows up first where E[n] becomes rational, the embedding-degree extension. That is W2 (MOV).

**The two ways around Lemma C also close.**

- ℓ = p (Artin–Schreier–Witt towers, which are geometric and plentiful): these control the p-primary Selmer group. Specialisation lands in E(F_p)[p^∞] = 0 because p ∤ n. That is W1 in characteristic p.
- The non-abelian n-division tower F_p(t)($\mathcal{E}$[n^k]), geometric part $SL_2(Z_n)$: its first layer is the function field of a modular curve of level n, of genus of order $n^3$ (W4). At a fibre it specialises to the field of definition of E[n], again an embedding-degree extension (W2).

**Verdict.** Closes (reduces to W2). Function-field Iwasawa theory at ℓ = n has no non-constant tower to work with.

# 4. What is left

Across the fourteen ideas, every Iwasawa mechanism falls into one of three kinds:

- p-adic analysis, which is blind to torsion prime to p (W1);
- n-adic Galois action, which costs field degree of order n or the embedding degree (W2, W4);
- a global lift, which is a free choice (W3).

The two new lemmas (A and C) are standard facts applied to this setting. They are recorded so that later leads can cite them. No residual lead is proposed. An Iwasawa-type idea that escaped all three kinds would need a tower whose first layer has degree o(√n), acts on the prime-to-p, n-part of E(F_p), and is computable without a lift. Lemma C rules this out for abelian towers over function fields of characteristic p, and Lemma A rules it out for Euler systems.

# Sources checked

- Huang, Raskind, *Global duality, signature calculus and the discrete logarithm problem*, arXiv:0710.2363 (LMS J. Comput. Math. 2009), abstract and summary only (web search).
- Ulmer, *Explicit points on the Legendre curve*, arXiv:1002.3313; Conceição, Hall, Ulmer, *Explicit points on the Legendre curve II*, arXiv:1307.4251 (web search summaries).
- `research/prime_field_leads_20261031/C1_lift_torsion/REPORT.md` (W1); `docs/guides/P256_PROPERTIES.md` §6 and line 413 (canonical-lift falsification).
- research-kb: not reachable in this session (no MCP tool exposed). The lead atlas was not checked (sibling checkout absent). Both are open before any atlas edit.

