Topological Anti-Gravity: π-Berry Phase Materials as Gravitational Sign Reversers in the Spectral Action Framework

Wright Brothers[1]

[1] *Independent Research*

(Dated: April 7, 2026)

We show that a material possessing a topological π-Berry phase reverses the sign of the effective gravitational coupling in Connes' spectral action framework. The mechanism is elementary: a π-phase shift applied to all vacuum modes transforms the spectral heat kernel from $\zeta(t)$ to $-\zeta(t)$, reversing the Seeley-DeWitt coefficient a_2 and hence $G_{\text{eff}} = -G$. No prime-by-prime Euler factor manipulation is required; a single global topological phase suffices. Candidate materials include π-Josephson junctions, topological insulator surfaces (Bi_2Se_3), and Weyl semimetals—all experimentally characterized. The anti-gravity effect is *passive*: the Berry phase is geometric, requiring no energy input. We propose two experimental tests: (1) a torsion balance measurement of gravitational anomaly near a π-junction cavity, and (2) the Casimir sign test in a π-junction-bounded resonator. If confirmed, π-Berry phase materials constitute the first known class of "anti-gravity matter"—substances that gravitationally repel rather than attract.

INTRODUCTION

All known matter gravitationally attracts. The weak energy condition (WEC) $T_{\mu\nu}u^\mu u^\nu \geq 0$ holds for all classical matter, and violations require exotic quantum states [**?**]. Here we identify a mechanism that reverses the *coupling* $G \to -G$ rather than the energy density $\rho \to -\rho$, achieving gravitational repulsion with ordinary positive-energy matter.

The mechanism arises in Connes' spectral action principle [**?**], where Newton's constant emerges from the spectrum of the Dirac operator D: $G^{-1} \propto a_2(D^2)$, the second Seeley-DeWitt coefficient of the heat kernel $K(t) = \text{Tr}(e^{-tD^2})$. If the spectrum is globally sign-reversed ($K \to -K$), then $a_2 \to -a_2$ and $G \to -G$.

We show that a material with topological π-Berry phase implements precisely this sign reversal.

THE SIGN-REVERSAL THEOREM

Theorem 1 (Topological anti-gravity)**.** *Let (A, H, D) be a spectral triple with heat kernel $K(t) = \text{Tr}(e^{-tD^2}) = \zeta(t)$ (the Bost-Connes identification [**?**]). If a global π-phase is applied to all modes, the heat kernel becomes*

$$K_\pi(t) = \text{Tr}\big((-1)\, e^{-tD^2}\big) = -\zeta(t), \tag{1}$$

and the effective gravitational coupling reverses:

$$\boxed{G_{\text{eff}} = -G.} \tag{2}$$

Proof. The Seeley-DeWitt expansion for $d = 4$: $K(t) = a_0 t^{-2} + a_2 t^{-1} + a_4 + \cdots$. The gravitational coupling is $G^{-1} \propto a_2 \propto K'(0)$.

For the normal case: $K(0) = \zeta(0) = -\frac{1}{2}$, $K'(0) = \zeta'(0) = -\frac{1}{2}\log(2\pi)$.

For the sign-reversed case: $K_\pi(0) = -\zeta(0) = +\frac{1}{2}$, $K'_\pi(0) = -\zeta'(0) = +\frac{1}{2}\log(2\pi)$.

The ratio:

$$\frac{G_{\text{eff}}}{G} = \frac{K'_\pi(0)}{K'(0)} = \frac{-\zeta'(0)}{\zeta'(0)} = -1. \tag{3}$$

□

The result is exact, not perturbative. Unlike the Liouville rotation [**?**] ($\zeta(2t)/\zeta(t)$, giving $G = -2G$ but requiring infinitely many operations), the global sign reversal requires a *single* operation: multiply every mode by (-1).

PHYSICAL MEANING: BOSONIC → FERMIONIC VACUUM

In quantum field theory, bosonic and fermionic fields contribute with opposite signs to the vacuum energy:

$$E_{\text{bos}} = +\tfrac{\hbar}{2}\sum_n \omega_n = +\tfrac{\hbar\omega_0}{2}\,\zeta(-d), \tag{4}$$

$$E_{\text{fer}} = -\tfrac{\hbar}{2}\sum_n \omega_n = -\tfrac{\hbar\omega_0}{2}\,\zeta(-d). \tag{5}$$

The global π-phase transforms every mode from bosonic (+) to fermionic (−) character. This is not a change of spin or statistics in the particle-physics sense; it is a change of the *vacuum energy sign* at the level of the spectral action.

In supersymmetric theories, the exact cancellation $E_{\text{bos}} + E_{\text{fer}} = 0$ solves the cosmological constant problem. Here we go further: a *fully fermionic* vacuum gives $E_{\text{total}} = -E_{\text{bos}}$, reversing the gravitational coupling.

π-BERRY PHASE MATERIALS

A topological π-Berry phase arises when a wave function acquires a geometric phase of exactly π upon traversal of a closed path in parameter space. This is topologically quantized and robust against perturbations.

Material	Mechanism	Ref.
π-Josephson junction	intrinsic π shift	[?]
Bi_2Se_3 surface	Dirac cone Berry phase	[?]
Graphene (K-point)	sublattice Berry phase	[?]
HgTe quantum well	2D TI edge state	[?]
Weyl semimetal	chiral anomaly	[?]

All candidates are experimentally well-characterized. π-Josephson junctions in particular can be fabricated as thin films [?] and integrated into superconducting circuits.

Implementation

A cavity whose boundary is a π-Berry phase material imposes the condition that every mode acquires a π-phase upon reflection from the boundary. The effective boundary condition is:

$$\psi(x_{\text{boundary}}) = -\psi(x_{\text{boundary}}), \tag{6}$$

which is an *anti-periodic* (fermionic) boundary condition.

The cavity spectral zeta function becomes $\zeta_{\text{cavity}}(t) = -\zeta(t) + O(\text{finite-size corrections})$, implementing the sign reversal of Theorem 1.

The effect is passive: the Berry phase is a property of the material's topology, not of an externally applied field. Once fabricated, the π-junction cavity maintains $G_{\text{eff}} = -G$ without energy input.

EXPERIMENTAL TESTS

Test 1: Casimir sign reversal

In a standard (bosonic-boundary) cavity: $E_{\text{Cas}} < 0$ (attractive Casimir force). In a π-junction cavity: $E_{\text{Cas}} > 0$ (repulsive Casimir force).

Setup. Compare the Casimir force between two plates: (A) both plates normal metal (DD, bosonic), (B) one or both plates π-Josephson junction film.

Prediction. $F_B/F_A = -1$ (sign reversal, same magnitude). Measurable with existing Casimir force apparatus [?].

Test 2: Torsion balance gravitational anomaly

A more direct test: measure the gravitational force between a π-junction cavity and a test mass using a torsion balance.

Prediction. If $G_{\text{eff}} = -G$ inside the cavity, the stress-energy tensor inside couples to gravity with reversed sign. The cavity would gravitationally *repel* the test mass.

The expected signal is extremely small ($\Delta F/F \sim \rho_{\text{Cas}}/\rho_{\text{bulk}} \ll 1$), but the *sign* is the key observable.

Test 3: Combined with 2-prime muting

The Euler product test (182:1 ratio in a 2-prime-sieved BAW cavity [?]) validates the spectral action framework. If positive, the π-junction test follows as the next step.

RELATION TO KNOWN PHYSICS

Why this is not trivially ruled out

One might object: a π-phase on cavity modes cannot affect gravity, because gravity couples to the stress-energy tensor $T_{\mu\nu}$, not to wave function phases.

In classical GR, this is correct. However, the spectral action principle *derives* $T_{\mu\nu}$ from the spectrum of D. If the spectrum includes a global π-phase, $T_{\mu\nu}$ itself is modified.

This is the distinction between:

- Standard GR: $T_{\mu\nu}$ is given, gravity follows.
- Spectral gravity: D is given, both $T_{\mu\nu}$ and $g_{\mu\nu}$ follow.

In the spectral framework, modifying D modifies everything simultaneously—matter *and* gravity.

Connection to dark energy

In previous work [?], we showed $\Omega_\Lambda = (d/\text{cd})\,|\xi_{\neg 2}(-1)| = 2\pi/9$, where the sign of dark energy ($\zeta_{\neg 2}(-1) = +1/12 > 0$) follows from $p = 2$ muting of the fermionic vacuum ($\zeta(-1) = -1/12 < 0$).

The π-Berry phase result extends this: *the full fermionic vacuum $-\zeta(t)$ is gravitationally repulsive at all scales*, not just at the cosmological scale. Dark energy ($\Omega_\Lambda > 0$) is the cosmological manifestation of the same sign reversal.

DISCUSSION

Assumptions. The result depends on Connes' spectral action principle applied to the Bost-Connes spectral triple. Both are mathematically rigorous but physically unverified for real gravity. The proposed experiments test this directly.

If confirmed: π-Berry phase materials would constitute the first known "anti-gravity matter"—substances whose topological properties reverse the gravitational coupling. Applications include shielding, propulsion, and the inner structure of warp bubble architectures [**?**].

If refuted: the spectral action framework does not describe physical gravity at the level of vacuum mode phases, and the Bost-Connes identification $K(t) = \zeta(t)$ is mathematical rather than physical.

Either outcome would be a significant result for the foundations of quantum gravity.

We thank the developers of PARI/GP, SageMath, and mpmath.