

Lecture 3

Axiomatic Definition of the Probability

23.02.2026

Outline

- 1 Axiomatic Probability
- 2 Some simple propositions
- 3 Independence

Outline

- 1 Axiomatic Probability
- 2 Some simple propositions
- 3 Independence

Axiomatic probability

Definition

A probability space consists of a sample space S and a probability function P which takes an event $A \subset S$ as input and returns $P(A)$, a real number between 0 and 1, as output. The function P must satisfy the following axioms:

- i For any event A , the probability of $0 \leq P(A) \leq 1$.
- ii $P(S) = 1$.
- iii If $A_1, A_2, \dots$ are disjoint events, then

$$P\left(\bigcup_{j=1}^{\infty} A_j\right) = \sum_{i=1}^{\infty} P(A_i).$$

Outline

- 1 Axiomatic Probability
- 2 Some simple propositions**
- 3 Independence

Propositions

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Proof (hint: use Axiom 2 and Axiom 3)

Propositions

Complement Rule:

$$P(A^c) = 1 - P(A)$$

Proof (hint: use Axiom 2 and Axiom 3)

- $P(\emptyset) = 0$

Difference Rule:

$$P(B \cap A^c) = P(B) - P(A), \text{ if } A \subseteq B$$

Proof (hint: use Axiom 3)

Difference Rule:

$$P(B \cap A^c) = P(B) - P(A), \text{ if } A \subseteq B$$

Proof (hint: use Axiom 3)

- $P(B) \geq P(A), \text{ if } A \subseteq B$

Proof (hint: use Axiom 1)

Inclusion-Exclusion: two events A, B (not necessarily disjoint)

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Proof (hint: use Axiom 3 and Venn-diagram)

Inclusion-Exclusion Example

J is taking two books along on her holiday vacation. With probability .5, she will like the first book; with probability .4, she will like the second book; and with probability .3, she will like both books. What is the probability that she likes neither book?

Inclusion-Exclusion Example

J is taking two books along on her holiday vacation. With probability .5, she will like the first book; with probability .4, she will like the second book; and with probability .3, she will like both books. What is the probability that she likes neither book?

① Conditions: event $B_1 = \{\text{J likes book 1}\}$, $B_2 = \{\text{J likes book 2}\}$,

$$P(B_1) = 0.5, P(B_2) = 0.4, P(B_1 \cap B_2) = 0.3$$

Inclusion-Exclusion Example

J is taking two books along on her holiday vacation. With probability .5, she will like the first book; with probability .4, she will like the second book; and with probability .3, she will like both books. What is the probability that she likes neither book?

- ① Conditions: event $B_1 = \{\text{J likes book 1}\}$, $B_2 = \{\text{J likes book 2}\}$,

$$P(B_1) = 0.5, P(B_2) = 0.4, P(B_1 \cap B_2) = 0.3$$

- ② Question: find $P(B_1^c \cap B_2^c) = 1 - P(B_1 \cup B_2)$.

Inclusion-Exclusion Example

J is taking two books along on her holiday vacation. With probability .5, she will like the first book; with probability .4, she will like the second book; and with probability .3, she will like both books. What is the probability that she likes neither book?

- ① Conditions: event $B_1 = \{\text{J likes book 1}\}$, $B_2 = \{\text{J likes book 2}\}$,

$$P(B_1) = 0.5, P(B_2) = 0.4, P(B_1 \cap B_2) = 0.3$$

- ② Question: find $P(B_1^c \cap B_2^c) = 1 - P(B_1 \cup B_2)$.

- ③ Formula: inclusion-exclusion

$$P(B_1 \cup B_2) = P(B_1) + P(B_2) - P(B_1 \cap B_2) = 0.5 + 0.4 - 0.3 = 0.6$$

$$1 - P(B_1 \cup B_2) = 1 - 0.6 = 0.4$$

Propositions

- Inclusion-Exclusion: three events A, B and C (not necessarily disjoint)

$$\begin{aligned} P(A \cup B \cup C) = & P(A) + P(B) + P(C) - P(A \cap B) \\ & - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C) \end{aligned}$$

Propositions

- Inclusion-Exclusion: n events $A_1, A_2, \dots, A_n$ (not necessarily disjoint)

$$\begin{aligned} P\left(\bigcup_{i=1}^n A_i\right) &= \sum_{i=1}^n P(A_i) - \sum_{1 \leq i < j \leq n} P(A_i \cap A_j) \\ &\quad + \sum_{1 \leq i < j < k \leq n} P(A_i \cap A_j \cap A_k) + \dots + \\ &\quad (-1)^{(n-1)} P(A_1 \cap A_2 \cap \dots \cap A_n) \end{aligned}$$

Example

Pick an integer in $[1, 1000]$ at random. Compute the probability that it is divisible neither by 12 nor by 15.

Example

A large company with n employees has a scheme according to which each employee buys a Christmas gift and the gifts are then distributed at random to the employees. What is the probability that someone gets his or her own gift?

Probability: Frequentist interpretation

- There are several possible interpretations of probability.
- There is no agreement, at all, in how probabilities should be interpreted.
- There is (nearly) complete agreement on the mathematical rules probability must follow (axioms).

Probability: Frequentist interpretation

- There are several possible interpretations of probability.
- There is no agreement, at all, in how probabilities should be interpreted.
- There is (nearly) complete agreement on the mathematical rules probability must follow (axioms).
- **Frequentist interpretation:** The probability of event A is the proportion of times (frequency) that A occurs in an infinite sequence (or very long run) of separate tries of the experiment.

$$P(A) = \lim_{n \rightarrow \infty} \frac{\# \text{ times } A \text{ happens}}{n}.$$

Probability: Frequentist interpretation

- There are several possible interpretations of probability.
- There is no agreement, at all, in how probabilities should be interpreted.
- There is (nearly) complete agreement on the mathematical rules probability must follow (axioms).
- **Frequentist interpretation:** The probability of event A is the proportion of times (frequency) that A occurs in an infinite sequence (or very long run) of separate tries of the experiment.

$$P(A) = \lim_{n \rightarrow \infty} \frac{\# \text{ times } A \text{ happens}}{n}.$$

- Often associated with Jerzy Neyman and Egon Pearson who described the logic of statistical hypothesis testing.

Probability: Bayesian interpretation

- A **Bayesian** can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.

Probability: Bayesian interpretation

- A **Bayesian** can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: For the same event, two separate people could have differing probabilities.

Probability: Bayesian interpretation

- A **Bayesian** can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: For the same event, two separate people could have differing probabilities.
- The Bayesian's view must: (1) conform to all other personal opinions; (2) change as new data arise according to **Bayes' Rule**.

Probability: Bayesian interpretation

- A **Bayesian** can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: For the same event, two separate people could have differing probabilities.
- The Bayesian's view must: (1) conform to all other personal opinions; (2) change as new data arise according to **Bayes' Rule**.
- Bayesian interpretations of probabilities avoid some of the philosophical difficulties of frequency interpretations.

Probability: Bayesian interpretation

- A **Bayesian** can pick whatever number they prefer for $P(A)$, based on their own personal experience and intuition, provided that number is consistent with all of the other probabilities they choose in life.
- A Bayesian interprets probability as a subjective degree of belief: For the same event, two separate people could have differing probabilities.
- The Bayesian's view must: (1) conform to all other personal opinions; (2) change as new data arise according to **Bayes' Rule**.
- Bayesian interpretations of probabilities avoid some of the philosophical difficulties of frequency interpretations.
- Named after the 18th century Presbyterian minister and mathematician Thomas Bayes.
- Largely popularized by revolutionary advance in computational technology and methods during the last twenty years.

Interpretations of Probability

- The Bayesian and frequentist perspectives are complementary, and both will be helpful for developing intuition in later chapters.
- Regardless of how we choose to interpret probability, we can use the two axioms to derive other properties of probability, and these results will hold for any valid probability function.

Outline

- 1 Axiomatic Probability
- 2 Some simple propositions
- 3 Independence**

Independence

Definition

Events A and B are independent if $P(A \cap B) = P(A)P(B)$.

Independence

Definition

Events A and B are independent if $P(A \cap B) = P(A)P(B)$.

- If A and B are independent, then learning that B occurred gives us no information that would change our probabilities for A occurring
- Independence is completely different from disjointness. If A and B are disjoint, then $P(A \cap B) = 0$, so disjoint events can be independent only if $P(A) = 0$ or $P(B) = 0$. Knowing that A occurs tells us that B definitely did not occur, so A clearly conveys information about B , meaning the two events are not independent (except if A or B already has zero probability).

Proposition

If A and B are independent events, then so are A^c and B , A and B^c , A^c and B^c are also independent.

Proof:

Example

An Internet packet travels from its source to router 1, from router 1 to router 2, and from router 2 to its destination. If routers drop packets independently with probability p , what is the probability that a packet is successfully transmitted from its source to its destination?