

Quantifying the Impacts of Cover Crops on Soil Moisture and Temperature

Emily Waring, Linda Schott, Ainis Lagzdins, Carl Pederson, Matt Helmers

Iowa State University

Introduction and Rationale

One of the leading causes of adverse environmental impacts in the Gulf of Mexico is nitrate-N originating from agricultural systems in the Midwest.



In Iowa, cover crops (left) are planted in the fall and grow in the spring before the cash crop is planted. These act as a mitigation strategy for nitrate-N leaching. However, they are a cost to the farmer and a potential decrease in yield because of the shift in water and temperature.

The objective of this study was:

1. To quantify the effects of a winter rye cover crop on soil moisture and temperature in a corn and soybean rotation in a tile drained system in north central Iowa.

Experimental Procedure

Study began in 2010 with 2 treatments (4 plots each):

- Corn and Soybean Rotation with Winter Rye
- Corn and Soybean Rotation without Winter Rye



DECAGON 5TE SOIL MOISTURE AND TEMPERATURE SENSORS

Placed at 4 depths in each plot:

10 cm, 20 cm, 40 cm, 60 cm



Logged and analyzed 5 minute intervals from 2012-2014



This research is part of a regional collaborative project supported by the USDA-NIFA, Award No. 2011-68002-30190 "Cropping Systems Coordinated Agricultural Project (CAP): Climate Change, Mitigation, and Adaptation in Corn-based Cropping Systems" sustainablecorn.org

SUSTAINABLE
CORN.ORG
CROPS, CLIMATE, CULTURE AND CHANGE



United States Department of Agriculture
National Institute of Food and Agriculture


Results and Discussion

When comparing No Till with Rye to No Till without Rye, only 27/ 324 Treatment-Month-Depth Combinations were different ($p < 0.05$). These differences are shown below.




DIFFERENCES IN MONTHLY AVERAGES


PRECIPITATION (mm)




MOISTURE ($\Delta\theta_v$)




TEMPERATURE ($\Delta^{\circ}C$)



2
0
1
3



2
0
1
4




Conclusions and Implications

- In most plots, Winter Rye did not significantly effect temperature or water.
- In plots with significant changes, both soybean and corn plots were well represented across all months except September.
- Soils with a winter rye cover crop were warmer and drier in most plots. Additional analysis is needed to further analyze this trend.

Acknowledgements


We would like to acknowledge Lance Henrichs, Alex Martin, and Quenton Schneider for their excellent field work and data collection!



Quantifying the Impacts of Cover Crops on Soil Moisture and Temperature

Emily Waring, Linda Schott, Ainis Lagzdins, Carl Pederson, Matt Helmers

Iowa State University



Recommend to put your institutional logo here

This research is part of a regional collaborative project supported by the USDA-NIFA, Award No. 2011-68002-30190 "Cropping Systems Coordinated Agricultural Project (CAP): Climate Change, Mitigation, and Adaptation in Corn-based Cropping Systems" sustainablecorn.org

**SUSTAINABLE
CORN.ORG**
CROPS, CLIMATE, CULTURE AND CHANGE

USDA
United States Department of Agriculture
National Institute of Food and Agriculture