

Logistic regression

Using python.

Iván Andrés Trujillo Abella

Facultad de Ingeniería
Pontificia Universidad Javeriana

trujilloiv@javeriana.edu.co
addajaveriana



Introduction

Logistic regression is used broadly in empirical works, it is used in economics, engineering, epidemiology and clinical research. In a simplified way the logistic regression it is used to binary problems.





If we take a point near to the change in concavity the method could produce divergence.

Bolzano theorem



illustration





$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} \quad (1)$$

Thus in i iteration we have:

$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)} \quad (2)$$

We select the point A the tangent line cross the x-axis in x_0 and again in $f(x_0)$ and again the tangent line of this point (B) cross the x-axis in x_1 each step is also known as a iteration. The algorithm converge to the the value x^* think that this could be used to *optimization problems*.



Newton-raphson program

```
def quadratic(a,b,c, x):  
    return a*x**2 + b*x + c  
def dfQuadratic(a,b,x):  
    return a*x**2 + b*x  
a,b,c ,x0 = 1,-3,-4,8  
def raphsonQuadratic(a,b,c,x0, error_max=0.0000015,  
    iteration_max=100):  
    xi = x0  
    iter, error = 0, 100  
    data = []  
    while (iter < iteration_max) and (error > error_max):  
        xj = xi - quadratic(a,b,c,xi) / dfQuadratic(a,b,xi)  
        error = abs(xj - xi)  
        iter += 1  
        data.append((xj,xi,error,iter))  
        xi = xj  
    return data  
raphsonQuadratic(a,b,c,4)[-1]
```



second order approximation



taylor series



Insights about classic and bayesian

Uses likelihood function estimation pretend assume that

$$P(A \mid B) = P(B \mid A)$$



$P(\Theta)$ prior distribution, posterior distribution $P(\Theta | X)$. likelihood $P(X | \Theta)$. Prior the belief before seen the data.



Conjugate prior



Gamma to poison

$$x \sim \text{Poisson}(\lambda)$$



Max a posteriori

A set of features $X = \{x_1, \dots, x_n\}$ assuming a distribution $P(X, \Theta)$ where Θ is a parameter (a random variable).

$$\Theta_{ma} = \max P(\Theta | X) \quad (3)$$

the last equation must be compared regarding the maximum likelihood estimation. that establish the $\max P(X | \Theta)$.



Bayesian continuous

$$f_{X|Y}(x | y) = \frac{f_{Y|X}(y | x)f_X(x)}{f_Y(y)} = \frac{f_{Y|X}(y | x)f_X(x)}{\int f_{Y|X}(y | x)f_X(x)dx} \quad (4)$$



Maximun Likelihood Estimation

MLE

Suppose that you data it is generated by a theoretical distribution, the inverse problem is determine the most probable parameter that generate the data.



Problem

you have n balls in a bag where there are j reds and k black thus $n = j + k$. However you do not know the really proportion of each one color. if you draw balls and both are different what is the proportion of black balls Θ .

MLE insight

Then we choose a Θ among all posible values that maximize the probability of seen the data.



Problem

In a formal way we can assume that $x \sim B(n, \Theta)$ and we have seen the data results $\{x_1, x_2, x_3, \dots, x_n\}$

$$P(X = x_1 \cap X = x_2, \dots, \cap X = x_n) \quad (5)$$

The variables are i.i.d and therefore the joint probability is the result of multiply the marginal probabilities.

$$P(X = x_1 \cap X = x_2, \dots, \cap X = x_n) = \prod_{i=1}^n P(x_i) \quad (6)$$



Problem

$$\begin{aligned}\prod_{i=1}^n P(x_i) &= \prod_{i=1}^n \binom{m}{x_i} \Theta^{x_i} (1 - \Theta)^{m-x_i} \\ &= \prod \binom{m}{x_i} \prod \Theta^{x_i} \prod (1 - \Theta)^{m-x_i} \\ &= \prod \binom{m}{x_i} \Theta^{\sum x_i} (1 - \Theta)^{\sum (m-x_i)}\end{aligned}\tag{7}$$

$$\ln(\prod P(x_i)) = \ln\left(\binom{m}{x_i}\right) + \sum x_i \ln(\Theta) + (nm - \sum x_i) \ln(1 - \Theta)$$



Limit bound Cramer-Rao

a bound to limit



Fisher information

Measure the amount of information that have a random variable about a parameter.



concepts about estimators

The quality of estimators:

- **Unbiased estimator** : The mean of the estimator is equal to the mean of parameter.
- **Variance**: the dispersion of the estimations regarding the mean value of the same.
- **quadratic value mean**: offer information about another two.



Topics to research

Teorema de Blackwell Teorema de Basu Teorema de Lehman-Scheffé
propiedad de invarianza? teorema de Zenha



Compare estimators



the lake is shallow



Score Statistics



Suppose that do you have a function $f(x, y) = x^2 + y^2$, then we can define the gradient as follow:

$$\nabla f(x, y) = \begin{bmatrix} \frac{df}{dx} \\ \frac{df}{dy} \end{bmatrix} \quad (9)$$



Notes about gradient

The gradient is perpendicular to each point in the level curve to any surface or level curve.



important resource

<https://www.youtube.com/watch?v=rB83DpBJQsE>



Escalar fields


it is a function that $f : R^m \rightarrow R$




Vector field

it is a function $\vec{F} = R^n \rightarrow R^n$ that associated a vector to each point in the euclidean space. you can use the gradient to construct the vector field.

$$x \exp(-x^2 - y^2)$$



Logistic equation



logistic equation

$$f(x) = \frac{\kappa}{1 + e^{-\alpha(x-x_0)}} \quad (10)$$

Where κ it is the maximun value.



logistic equation

Population growth

$$\frac{dy}{dt} = ry\left(1 - \frac{y}{k}\right) \quad (11)$$



python implementation

Statsmodels

```
formula = 'died~studytime + C(drug) ' # logit died studytime
        i.drug
model = smf.logit(formula= formula, data=df)
results = model.fit()
print(results.summary())
coefs = pd.DataFrame({
    'coef': results.params.values,
    'odds ratio': np.exp(results.params.values),
    'name': results.params.index
})
coefs
```



Prediction score

The usual way



Talleres

1. structure problems 2. loop problems 3. Combinatorial problems 4. derivatives systems 5. matrix algorithm 6. matrix algorithm 7. Economic growth modeling 8. Analytic problem 9. Analytic problem 7 k-means algorithm 8. perceptron algorithm 9. World bank data 10. Icfes data 11. Mortality Data 12. review project



Solve

with newton method solve the following:

$$e^x = 4 - x^2 \quad (12)$$

las dos soluciones son: $x_1 = 1.05$ $x_2 = -1.96$



Entropy



scoring

