

Entanglement in Featureless Mott Insulators

Brayden Ware

September 23th 2014

Outline

1 Motivation

- Featureless Insulators
- Topological Band Insulators
- Bosonic Band Insulators?

2 Featureless Boson Mott Insulators

- Honeycomb Featureless Insulator Proposal

3 Entanglement Spectra

- CFT Identification

Outline

1 Motivation

- Featureless Insulators
- Topological Band Insulators
- Bosonic Band Insulators?

2 Featureless Boson Mott Insulators

- Honeycomb Featureless Insulator Proposal

3 Entanglement Spectra


- CFT Identification

Featureless Insulators


Definition of 'Featureless Insulator'

- Gapped
- Symmetric
- No topological order
- Integer charge per unit cell
- Unique ground state with P.B.C.

Examples:



Band Insulator



Bosonic Mott insulator
with integer filling



Heisenberg AF Spin-1 chain

Free Fermion Band Insulators

- Crystalline, 0T insulators (including semiconductors)
- Tight-binding Hamiltonian

$$\mathcal{H}_{FF} = \sum_{\langle ij \rangle} \sum_{\alpha, \beta} -t_{\alpha, \beta} c_i^{\alpha \dagger} c_j^{\beta} - \mu \sum_{i, \alpha} N_i^{\alpha}$$

- Bloch Wavefunctions $|u_{\mathbf{k}}^{\alpha}\rangle$
- Massive Dirac Hamiltonian

$$\mathcal{H}_D(\mathbf{k}) = \mathbf{k}_x \sigma_x + \mathbf{k}_y \sigma_y + m_* \sigma_z$$

- Atomic-insulating like Wannier basis



Semiconductor GaN

Free Fermion Band Insulators

- Crystalline, 0T insulators (including semiconductors)
- Tight-binding Hamiltonian

$$\mathcal{H}_{FF} = \sum_{\langle ij \rangle} \sum_{\alpha, \beta} -t_{\alpha, \beta} c_i^{\alpha \dagger} c_j^{\beta} - \mu \sum_{i, \alpha} N_i^{\alpha}$$

- Bloch Wavefunctions $|u_{\mathbf{k}}^{\alpha}\rangle$
- Massive Dirac Hamiltonian

$$\mathcal{H}_D(\mathbf{k}) = \mathbf{k}_x \sigma_x + \mathbf{k}_y \sigma_y + m_* \sigma_z$$

- Atomic-insulating like Wannier basis



Band Theory

Free Fermion Band Insulators

- Crystalline, 0T insulators (including semiconductors)
- Tight-binding Hamiltonian

$$\mathcal{H}_{FF} = \sum_{\langle ij \rangle} \sum_{\alpha, \beta} -t_{\alpha, \beta} c_i^{\alpha \dagger} c_j^{\beta} - \mu \sum_{i, \alpha} N_i^{\alpha}$$

- Bloch Wavefunctions $|u_{\mathbf{k}}^{\alpha}\rangle$
- Massive Dirac Hamiltonian

$$\mathcal{H}_D(\mathbf{k}) = \mathbf{k}_x \sigma_x + \mathbf{k}_y \sigma_y + m_* \sigma_z$$

- Atomic-insulating like Wannier basis



Dirac Band Theory

Free Fermion Band Insulators

- Crystalline, 0T insulators (including semiconductors)
- Tight-binding Hamiltonian

$$\mathcal{H}_{FF} = \sum_{\langle ij \rangle} \sum_{\alpha, \beta} -t_{\alpha, \beta} c_i^{\alpha \dagger} c_j^{\beta} - \mu \sum_{i, \alpha} N_i^{\alpha}$$

- Bloch Wavefunctions $|u_{\mathbf{k}}^{\alpha}\rangle$
- Massive Dirac Hamiltonian

$$\mathcal{H}_D(\mathbf{k}) = \mathbf{k}_x \sigma_x + \mathbf{k}_y \sigma_y + m_* \sigma_z$$

- Atomic-insulating like Wannier basis



Wannier Function

Motivating Questions

Existence

Are there **constraints** on the existence of featureless insulators?
Can we **construct** featureless insulators when possible?

Given a lattice Λ , and an integer m , is there a featureless insulating phase of matter with m particles per unit cell?

Uniqueness

How can we **distinguish** different classes of featureless insulators?
Can we **enumerate** all such classes?

Frame Title

Frame Title

Outline

1 Motivation

- Featureless Insulators
- Topological Band Insulators
- Bosonic Band Insulators?

2 Featureless Boson Mott Insulators

- Honeycomb Featureless Insulator Proposal


3 Entanglement Spectra

- CFT Identification

Frame Title


Honeycomb Featureless Boson Insulator

Proposed Wavefunction




Honeycomb Featureless Boson Insulator

Proposed Wavefunction




Honeycomb Featureless Boson Insulator

Proposed Wavefunction




Honeycomb Featureless Boson Insulator

Proposed Wavefunction



Honeycomb Featureless Boson Insulator

A wavefunction written as a product of local operators acting on a product state can simply be turned into a tensor network.



Outline

1 Motivation

- Featureless Insulators
- Topological Band Insulators
- Bosonic Band Insulators?


2 Featureless Boson Mott Insulators

- Honeycomb Featureless Insulator Proposal

3 Entanglement Spectra

- CFT Identification

Entanglement Spectra




Finite Size Analysis of Spectra


- Topological entanglement entropy is 0
- Low energy modes show gapless $1/L$ behavior

Finite Size Analysis of Spectra


- Topological entanglement entropy is 0
- Low energy modes show gapless $1/L$ behavior




Identifying CFTs: Measuring c




Conformal Tower




Conformal Tower



Conformal Tower



Conformal Tower



Level identification in CFT spectra

To make a precise comparison with the free-boson CFT, we'll need to solve for (or look up) the solution of this model.

The free-boson CFT is created from the Lagrangian

$$\mathcal{L} = \frac{g}{2} \int dt \int_0^L dx \left(\frac{1}{v^2} (\partial_t \phi)^2 - (\partial_x \phi)^2 \right)$$

and with the compactified field identification

$$\phi \equiv \phi + 2\pi R$$

and placed on the circle of circumference L with periodic boundary conditions


$$\phi(x) \equiv \phi(x + L).$$

Level identification in CFT spectra


\mathbf{L}_0	$2\pi g \left(\frac{e}{4\pi g R} + \frac{mR}{2} \right)^2 + n$
$\bar{\mathbf{L}}_0$	$2\pi g \left(\frac{e}{4\pi g R} - \frac{mR}{2} \right)^2 + \bar{n}$
$\mathbf{P} = \frac{2\pi v}{L} (\mathbf{L}_0 - \bar{\mathbf{L}}_0)$	$\frac{2\pi v}{L} (em + n - \bar{n})$
$\mathbf{H} = \frac{2\pi v}{L} (\mathbf{L}_0 + \bar{\mathbf{L}}_0)$	$\frac{2\pi v}{L} \left(\frac{e^2}{4\pi g R^2} + \pi g m^2 R^2 + n + \bar{n} \right)$
$\tilde{\mathbf{H}} = \frac{L}{2\pi v \kappa} \mathbf{H}$	$e^2 + \frac{m^2}{4\kappa^2} + \frac{1}{\kappa} (n + \bar{n})$

Eigenvalues of states $|e, m\rangle_{n, \bar{n}}$. The rescaled Hamiltonian $\tilde{\mathbf{H}}$ has eigenvalues that depend on only one free-parameter, $\kappa = 1/(4\pi g R^2)$. (Note: A common convention is to set $g = 1/4\pi$ and describe the system using $R = \sqrt{1/\kappa}$.)

Level identification in CFT spectra



Level identification in CFT spectra



Open questions and speculation

Correspondence between edge and bulk physics In a RG-fixed point tensor network state (such as toric code): Can

Resources



A Practical Introduction to Tensor Networks

Orus, R. A Practical Introduction to Tensor Networks: Matrix Product States and Projected Entangled Pair States. arXiv [cond-mat.str-el] (2013). at <http://arxiv.org/abs/1306.2164>

Questions?

Brayden Ware

brayden@physics.ucsb.edu

Bonus slides

Bonus slides