
Changed significant GO Biological Process pathways
Both contrasts adjusted P <= 0.1; shown if NES sign switches or |delta NES| >= 1
−log10 adjusted P
5
10
15
20
Gene set size
100
200
300
young CD vs old CD
old ID
Switched: young−down, ID−up
humoral immune response
cell killing
antigen processing and presentation of peptide antigen
adaptive immune response
humoral immune response mediated by circulating immunoglobulin
complement activation
leukocyte mediated immunity
complement activation, classical pathway
myeloid leukocyte migration
leukocyte chemotaxis
adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin superfamily domains
granulocyte migration
antigen processing and presentation
defense response to Gram−positive bacterium
B cell mediated immunity
myeloid leukocyte activation
immunoglobulin mediated immune response
lymphocyte mediated immunity
antigen processing and presentation of exogenous peptide antigen
leukocyte migration
killing of cells of another organism
disruption of cell in another organism
neutrophil migration
antigen processing and presentation of peptide antigen via MHC class I
myeloid leukocyte mediated immunity
antigen processing and presentation of exogenous antigen
antimicrobial humoral immune response mediated by antimicrobial peptide
disruption of anatomical structure in another organism
granulocyte chemotaxis
mononuclear cell migration
−2
0
2
Normalized enrichment score