

**Olin College of Engineering**  
**ENGR2410 – Signals and Systems**

**Assignment 9**

**Problem 1** (2 points) Show Parseval's Theorem

$$\int_{t=-\infty}^{t=\infty} [x(t)]^2 dt = \int_{\omega=-\infty}^{\omega=\infty} |X(j\omega)|^2 \frac{d\omega}{2\pi}$$

where

$$|X(j\omega)|^2 = X(j\omega)X^*(j\omega)$$

$x(t)$  is real, and  $X^*(j\omega)$  is the complex conjugate of  $X(j\omega)$ .

*Hint:*


$$X^*(j\omega) = \int_{t=-\infty}^{t=\infty} x(t)e^{j\omega t} dt$$

**Problem 2** (4 points) A non-ideal filter or communication channel  $H(j\omega)$  with finite *transition band*  $\Delta f$  between the *passband* (where  $H(j\omega) \neq 0$ ) and *stopband* (where  $H(j\omega) = 0$ ) can be modeled as  $H(j\omega) = H_0(j\omega) * H_\Delta(j\omega)$  where

$$H_0(j\omega) = \begin{cases} 1 & -2\pi f_0 < \omega < 2\pi f_0 \\ 0 & \text{otherwise} \end{cases} \quad H_\Delta(j\omega) = \begin{cases} \frac{1}{2\pi\Delta f} & -2\pi\Delta f/2 < \omega < 2\pi\Delta f/2 \\ 0 & \text{otherwise} \end{cases}$$

- A. Find an expression for  $h_0(t)$ , the impulse response of the ideal channel.
- B. Find an expression for  $H(j\omega)$  and sketch it.
- C. Find an expression for the impulse response  $h(t)$  of the non-ideal channel.
- D. Plot  $h(t)$  and  $h_0(t)$  on the same axes when  $f_0 = 10$  kHz and  $\Delta f = 2$  kHz and sketch the associated  $H(j\omega)$ . Repeat the plot for the same values when  $\Delta f = 5$  kHz. Compare the plots. For what values of  $\Delta f$  is the effect of the non-ideal transition band noticeable? Is the effect what you would expect? Explain.

**Problem 3** (4 points) The system shown below represents a basic communication system where two messages  $x_1(t)$  and  $x_2(t)$  share a common communication channel. Signals  $x_1(t)$  and  $x_2(t)$  are bandlimited to  $f_B$  and have a frequency content as shown below. The receiver has an ideal low-pass filter  $H(j\omega)$  with a cutoff frequency of  $f_B$  as shown below.



- What would happen if  $\omega_1 = \omega_2 = 0$ ? Find  $y(t)$  in terms of  $x_1(t)$  and/or  $x_2(t)$ , and show its frequency content.
- Find constraints on  $\omega_1$  and  $\omega_2$  such that there is no frequency interference (aliasing). Show the frequency content of  $m(t)$  and  $d(t)$  under these constraints. *Note: There may be multiple solutions; just find one that works.*
- Show the frequency content and find an algebraic expression for  $y(t)$  in terms of  $x_1(t)$  and/or  $x_2(t)$  assuming the constraints of part B.