

# Hercules-Aquila and Virgo Clouds with Gaia DR2. Evidence for a common origin

Keith T. Smith,<sup>1</sup><sup>★</sup> A. N. Other,<sup>2</sup> Third Author<sup>2,3</sup> and Fourth Author<sup>3</sup>

<sup>1</sup>*Royal Astronomical Society, Burlington House, Piccadilly, London W1J 0BQ, UK*

<sup>2</sup>*Department, Institution, Street Address, City Postal Code, Country*

<sup>3</sup>*Another Department, Different Institution, Street Address, City Postal Code, Country*

Accepted XXX. Received YYY; in original form ZZZ

## ABSTRACT

200 words for Letters. No references should appear in the abstract.

**Key words:** keyword1 – keyword2 – keyword3

## 1 INTRODUCTION

How do you hide the evidence for a massive impact event that caused the extinction of most of the dinosaurs as well as 75% of all species on Earth? You bury it deep under the sea, covered with a layer of sediment taller than the Empire State Building (Hildebrand et al. 1991). Without the discovery of the giant Chicxulub crater, the meteorite impact hypothesis would remain a neat theory supported by striking but indirect evidence. A hypothesis of an ancient dramatic collision between the Milky Way and an unidentified massive dwarf galaxy was put forward by Deason et al. (2013) to explain a particular feature in the overall stellar halo density profile (Watkins et al. 2009; Deason et al. 2011). Most recently, through a study of a portion of the nearby stellar halo, Belokurov et al. (2018b) demonstrated that the great impactor must have collided with the young Milky Way on a nearly radial orbit, thus swamping the inner stellar halo with metal-rich material with orbital anisotropy (see Binney & Tremaine 2008) close to unity. Merger events like this tend to leave behind a battery of debris clouds and shells (see Amorisco 2015; Hendel & Johnston 2015), which - akin to the peak rings of impact craters (see e.g. Morgan et al. 2016) - if discovered could help to reconstruct the collision as well as pin down the properties of the progenitor (e.g. Sanderson & Helmi 2013; Johnston 2016).

Before the Data Release 2 (Gaia Collaboration et al. 2018) of the ESA’s Gaia mission (Gaia Collaboration et al. 2016), five large and diffuse cloud-like structures had been discovered in the Galaxy’s halo. These include: the Virgo Over-Density (VOD, Vivas et al. 2001; Newberg et al. 2002; Duffau et al. 2006; Jurić et al. 2008; Bonaca et al. 2012), the Hercules-Aquila Cloud (HAC, Belokurov et al. 2007; Simion et al. 2014), the Tringulum-Andromeda structure (Tri-And, Rocha-Pinto et al. 2004; Majewski et al. 2004; Deason et al. 2014), the Pisces Over-density (Sesar et al.

2007; Watkins et al. 2009; Nie et al. 2015) and the Eridanus-Phoenix over-density (Eri-Pho, Li et al. 2016). According to the most recent investigations, Tri-And likely comprises of Galactic disc stars kicked out of the plane in a recent interaction with a dwarf galaxy, probably the Sagittarius dSph (e.g. Price-Whelan et al. 2015; Bergemann et al. 2018; Hayes et al. 2018). Of the remaining four, the Pisces overdensity clearly stands out as it reaches much larger Galacto-centric distances. On the other hand, the VOD, HAC and Eri-Pho structures occupy a very similar range of distances, between 10 and 20 kpc from the Galactic center. This lead Li et al. (2016) to suggest that these three Clouds could all be part of one merger event, a galaxy accreted onto the Milky Way on a polar orbit (see also Jurić et al. 2008).


As demonstrated by the recent re-interpretation of the Monoceros Ring (and the associated sub-structures) and the Tri-And, deciphering the nature of halo over-densities is often impossible without either high-resolution spectroscopy (e.g. Bergemann et al. 2018) or accurate astrometry (e.g. de Boer et al. 2018; Deason et al. 2018a). In this Letter, we look for clues to the formation of the Hercules-Aquila and Virgo Clouds using proper motions provided as part of the Gaia DR2. At our disposal are highly pure samples of members of each Cloud, namely the RR Lyrae stars that i) are co-spatial with HAC and VOD in 3-D and ii) that have their line-of-sight velocities measured. By complementing the publicly available 4-D data with the GDR2 proper motions, we build a large tracer set with complete 6-D phase space information and study the make-up of each structure using the orbital properties of the constituent stars.

## 2 DATA AND ANALYSIS

### 2.1 4-D data

The Hercules-Aquila and Virgo Clouds are revealed unambiguously as overdensities of RR Lyrae in the halo, which peak at similar heliocentric distances, ~17 kpc (HAC: Simion

<sup>★</sup> E-mail: mn@ras.org.uk (KTS)



**Figure 1.** Spatial distribution of the RR Lyrae used in this work with full 6-D phase space measurements, in Galactic coordinates (left panel) and in the  $x-y$  (middle) and  $y-z$  (right) planes. The HAC field contains 44 RR Lyrae which likely belong to the Cloud with measured line-of-sight velocities (Simion et al. 2018) and Gaia DR2 proper motions. The VOD field contains 411 RRL which belong to several halo associations, including the Sagittarius stream and the VOD, with line-of-sight velocities provided by Vivas et al. 2016 and proper motions from Gaia DR2. In particular we mark group 2, a ‘high significance’ kinematical group, which contains 18 stars (green circles). The semi-circles are centred on the Sun’s position and have radius of 10 and 20 kpc. The Sun (yellow star) is located at  $(x_{\odot}, y_{\odot}, z_{\odot}) = (0, -8, 0)$  kpc and the Galactic centre at  $(0, 0, 0)$  - black circle.


et al. 2014) and  $\sim 19$  kpc (VOD: Vivas & Zinn 2006; Duffau et al. 2014; Vivas et al. 2016) respectively, but in opposite quadrants (e.g. their distribution in Galactic coordinates in Fig. 1). Other tracers (e.g. BHBs, MSTO, K giants) have been used to pin down the morphology of the Clouds with slightly inconsistent results between them, caused by poor distance determination, scarcity or simply trace the structures differently; therefore, the most recent kinematic follow-ups of the clouds have used RR Lyrae selected in the vicinity of the overdensities peak:

- Simion et al. (2018) provide a table of 46 RRL with radial velocity measurements (45 observed with MDM and 1 from SDSS) with heliocentric distances between 15 and 18 kpc;
- Vivas et al. (2016) compiled a catalog of 412 RRL in the region of the sky covered by the VOD with distances between 4 and 75 kpc from the Sun with radial velocity measurements from La Silla-QUEST, QUEST, CRTS and LINEAR.

## 2.2 6-D data: velocity distributions

From the two tables of RR Lyrae with 4D measurements, 44 stars in the HAC field and 411 in the VOD field have matches in GDR2 within  $2''$  and proper motion measurements. The only star in the VOD region without proper motion belongs to a ‘high-significance’ kinematical group (group 1), likely the Sagittarius stream, identified by Vivas et al. (2016). This group contains 113 stars (112 with proper motions) which we remove from the VOD sample as a major contaminant of the field. The spatial distribution of the remaining stars (44 from Simion et al. 2018 and 299 from Vivas et al. 2016) with full 6-D phase space measurements is illustrated in Figure 1, in Galactic coordinates (left panel) and in the  $(x-y)$  Galactic plane and the  $(x-z)$  plane, perpendicular to  $(x-y)$ . We adopted a left-handed Galactic Cartesian coordinates with the Sun located at  $(x_{\odot}, y_{\odot}, z_{\odot}) = (-8, 0, 0)$  kpc, the  $x$ -axis positive in the direction of the Galactic center,  $y$ -axis oriented along the Galactic rotation and the  $z$ -axis directed towards the north Galactic pole.


While Vivas et al. 2016 identify 6 significant kinematical groups in the VOD field (their table 5) only groups 1 (Sagittarius stream) and 2 (likely members of the VOD, with



**Figure 2.** RRL velocity distribution in spherical polar coordinates ( $v_r, v_{\theta}, v_{\phi}$  are the radial, azimuthal and polar components respectively) in the HAC field (top row) and the VOD field (middle and bottom rows). The error on the velocity components of each star  $i$ ,  $[\sigma_{v_r}^i, \sigma_{v_{\theta}}^i, \sigma_{v_{\phi}}^i]$ , has been propagated by randomly drawing 1000 stars from a multivariate Gaussian distribution with mean the measurement ( $ra^i, dec^i, d^i, pmra^i, pmdec^i, v_h^i$ ) and full covariance matrix (takes into account the covariances between  $ra, dec$  and proper motions). The orbital anisotropy, is highly radial in the HAC field ( $\beta = 0.91 \pm 0.03$ ) where the stars are most likely members of the Cloud and mildly radial in the VOD field ( $\beta = 0.74 \pm 0.04$ ) in which stars span a much wider range of distances (see Fig. 1).

$\langle v_{GSR} \rangle = 135$  km/s) contain more than 10 stars. We mark group 2 with green circles in Figure 1. We also mark with red circles the location of a group of stars with galactocentric distances similar to the HAC sample,  $11 < r_{GC}/kpc < 16$  as we intend to follow their velocity distribution and orbital properties.

The velocity distribution in the VOD and HAC fields is shown in Fig. 2 in spherical polar coordinates, where  $v_r, v_{\theta}, v_{\phi}$  are the radial, azimuthal and polar components respectively. ‘Group 2’ stars (lime) form a group at  $v_r = 135$  km/s as expected, while the stars at intermediate  $r_{GC}$  seem to have a velocity distribution very similar to HAC, shown in the top panels. To estimate the error on each velocity



**Figure 3.** Radial versus azimuthal velocity in the VOD field, in three distance ranges above the Galactic plane. The fraction of RR Lyrae of Oosterhoff type I is reported in each panel.

component we resample the data 1000 times from a multivariate Gaussian distribution with mean the measurement  $\{\alpha, \delta, d_{\text{helio}}, \mu_{\alpha \cos(\delta)}, \mu_{\delta}, v_{\text{helio}}\}_i$  and full covariance matrix which takes into account the covariances between the right ascension  $\alpha$ , declination  $\delta$  and proper motions, provided by GDR2. We take the standard deviation of the resulting  $\{v_r, v_\theta, v_\phi\}$  distributions as the upper limit of the velocity uncertainties and show them in Fig. 2.


We calculate the velocity anisotropy in the two fields, keeping in mind the two samples span very different distance ranges. We compute the median and standard deviation over 500 non parametric bootstrap resampling trials, where each trial was modelled with a velocity ellipsoid using the Extreme Deconvolution module implemented in astroML (Vanderplas et al. 2012). We find the HAC stars have radially biased orbits ( $\beta = 0.91 \pm 0.03$ ) while the VOD stars have slightly less radial orbits ( $\beta = 0.74 \pm 0.04$ ).

We fit a model with two multivariate Gaussians to the VOD velocity ellipsoid using Extreme Deconvolution and obtain  $\beta_1 = 0.96^{+0.02}_{-0.44}$  (marked in magenta in Fig. 2) and  $\beta_2 = 0.44^{+0.45}_{-0.20}$  (blue).

Fig. 3 shows the behaviour of the VOD azimuthal  $v_\theta$  and radial  $v_r$  velocity distributions in 3 distances ranges above the Galactic plane. In each slice we have calculated the fraction of Oosterhoff type I (Oo I) RR Lyrae, using equations 1 and 2 in Belokurov et al. (2018a) to classify the RRL into two types. According to this classification, Oosterhoff type II (Oo II) RR Lyrae will include both Oo II and Intermediate objects. In the  $10 < z/\text{kpc} < 20$  range, where the velocity anisotropy is the highest ( $\beta = 0.84 \pm 0.03$ ) approaching the value in the HAC field, the Oo I type dominates (77%). In the same slice, 73% of the stars belong to the more squashed (or ‘sausage’ looking) velocity ellipsoid. The same behaviour but less accentuated can be noticed in the  $0 < z/\text{kpc} < 10$  slice where the anisotropy  $\beta = 0.7 \pm 0.1$  is lower but the fraction of Oo I stars decreases dramatically to 57%. Further from the plane, at  $z > 20$  kpc, the velocity ellipsoid is almost isotropic with  $\beta = -0.1 \pm 0.2$ . Here, the  $\beta$  value may still be affected by the presence of the Sagittarius stream in the field. Group 2 and red points are all located at  $z < 20$  kpc, with the red points following an anisotropic velocity distribution.

### 2.3 Orbital properties of the HAC and VOD

For orbit integration we use the galpy package Bovy (2015) in the recommended MWPotential2014 model for the Galactic potential which is composed of a Miyamoto-Nagai disc, a bulge with a power-law density profile that is exponen-



**Figure 4.** Orbital properties of the stars in the HAC and VOD fields. ‘group 2’ has similar orbital properties to the HAC, however it does not display a sausage velocity distribution (see middle row figure 2) - they are concentrated at  $v_r = 135 \text{ km/s}$  as calculated by Vivas et al. 2016.


tially cut-off, and a dark matter halo described by a NFW potential. The parameters are given in Bovy (2015), table 1. The resulting orbital parameters of the HAC and VOD are given in Fig. 4. To compute the errors (not shown for VOD to simply the figure) we integrated 500 orbits for each star where the orbits were initialised on parameters resampled from data, as in the previous section. The pdfs of the eccentricities, apocentres and pericentres are also shown. If the HAC orbits are clearly highly eccentric, we find that the majority of stars in VOD are also on quite eccentric orbits: in particular, the stars with  $11 < r_{\text{GC}}/\text{kpc} < 16$  (in red), follow closely the HAC orbital properties.

### 3 ARE THE VOD AND THE HAC RELATED?


In this section we discuss the possible connection between the two clouds. We use the 6-D phase space measurements as initial conditions for backward orbit integration. We integrate the orbits for 8 Gyrs and show them in Fig. 5.

### 4 CONCLUSIONS

Using a sample of  $\sim 500$  RR Lyrae with full 6-D phase space information, we have studied the orbital properties of the Hercules-Aquila and Virgo Clouds. Both Clouds appear dominated by stars on highly eccentric orbits. Assuming that the kinematics of each structure is well described by a single Gaussian, the orbital anisotropy of the HAC is  $\beta = 0.91$  and for the VOD,  $\beta = 0.74$ . Note, however that the original criteria applied to the RR Lyrae stars to select targets for spectroscopic follow-up differ drastically between



**Figure 5.** The backward orbit integration for HAC (top panels) and VOD (bottom panels) for 8 Gyrs look back time. We use  $M_{vir} = 0.8 \times 10^{12} M_{\odot}$ , the default galpy value.  $\log(N)$  shown, notice the change in colour scale between top and bottom rows. The present day loci of HAC and VOD are marked with gray rectangles. The initial conditions of 44 stars with heliocentric distances between 15 and 18 kpc were used for the HAC backward orbit integration and of 299 stars with heliocentric distances between 4 and 75 kpc for the VOD orbit integration.



**Figure 6.** EXTRA: angular momentum - I can combine these two and also add another panel with energy vs total  $L$  when I figure out the energy.

the HAC and the VOD datasets. The HAC sample covers a very limited region in the of  $l, b, D$  space, while the VOD dataset spans a wide range of longitudes, latitudes and heliocentric distances. It is therefore likely that the VOD dataset contains a mixture of several halo sub-structures (see [Vivas et al. 2016](#), for a detailed discussion). For the entirety of the analysis described here, we made sure to cull the probable Sgr stream members. Additionally, we explore how the VOD's make-up changes with Galactic height and demonstrate that for  $|z| < 20$  kpc, the VOD orbital anisotropy is  $\beta \sim 0.84$ , while above this threshold, it quickly changes to  $\beta \sim 0$ . We conclude therefore, that an assumption of a sin-

gle Gaussian for the entire VOD sample is not appropriate. Modeling the kinematics of the VOD stars with a mixture of 2 multi-variate Gaussians, we show that two thirds of the VOD sample are the stars with  $\beta = 0.96$ , while the remainder has  $\beta = 0.44$ , in good agreement with the local measurement presented in [Belokurov et al. \(2018b\)](#).

As revealed by Gaia, the two structures are composed of stars on nearly radial orbits, with peaks in the eccentricity distribution at 0.9 (0.8) for the HAC (VOD). The distributions of the peri-centric and apo-centric distances also match: the stars in the Clouds turn around at 1 – 3 and 15 – 30 kpc. Not only the HAC and the VOD look alike kinematically, their orbital composition is in perfect agreement with the stellar halo properties as analysed locally by [Belokurov et al. \(2018b\)](#) and globally (out to 40 kpc) by [Deason et al. \(2018b\)](#). As these authors demonstrate, the inner halo is dominated by metal-rich debris from an old and massive accretion event. In particular, [Belokurov et al. \(2018b\)](#) use Cosmological simulations of Milky Way halo formation, to bracket the time of the merger - between 8 and 11 Gyr ago - and its mass, which they show to be in excess of  $10^{10} M_{\odot}$ . The tell-tale sign of this dramatic head-on collision is the particular shape of the corresponding stellar velocity ellipsoid, which is stretched so much in the radial direction compared to the tangential ones, that it resembles a sausage.



An alternative view of this merger can be found in [Myeong et al. \(2018a\)](#), where the local stellar halo is mapped out in the action space. Here, the metal-rich stars are shown have extended radial action distribution in addition to a prominent spray of material on retrograde orbits. The high mass of the progenitor is evidenced not only by the metallicity distribution of its likely member or the numerical simulations of halo formation, but also by a sizeable number of Globular Clusters that could be attributed to the same event (see [Myeong et al. 2018b](#)).

## ACKNOWLEDGEMENTS

The Acknowledgements section is not numbered. Here you can thank helpful colleagues, acknowledge funding agencies, telescopes and facilities used etc. Try to keep it short.

## REFERENCES

- Amorisco N. C., 2015, *MNRAS*, 450, 575
- Belokurov V., Deason A. J., Koposov S. E., Catelan M., Erkal D., Drake A. J., Evans N. W., 2018a, *MNRAS*, 477, 1472
- Belokurov V., Erkal D., Evans N. W., Koposov S. E., Deason A. J., 2018b, *MNRAS*, 478, 611
- Belokurov V., Evans N. W., Bell E. F., Irwin M. J. et al., 2007, *ApJ*, 657, L89
- Bergemann M., Sesar B., Cohen J. G., Serenelli A. M. et al., 2018, *Nature*, 555, 334
- Binney J., Tremaine S., 2008, *Galactic Dynamics: Second Edition*. Princeton University Press
- Bonaca A., Jurić M., Ivezić Ž., Bizyaev D. et al., 2012, *AJ*, 143, 105
- Bovy J., 2015, *ApJS*, 216, 29
- de Boer T. J. L., Belokurov V., Koposov S. E., 2018, *MNRAS*, 473, 647
- Deason A. J., Belokurov V., Evans N. W., 2011, *MNRAS*, 416, 2903
- Deason A. J., Belokurov V., Evans N. W., Johnston K. V., 2013, *ApJ*, 763, 113
- Deason A. J., Belokurov V., Hamren K. M., Koposov S. E. et al., 2014, *MNRAS*, 444, 3975
- Deason A. J., Belokurov V., Koposov S. E., 2018a, *MNRAS*, 473, 2428
- Deason A. J., Belokurov V., Koposov S. E., Lancaster L., 2018b, *ArXiv e-prints*
- Duffau S., Vivas A. K., Zinn R., Méndez R. A., Ruiz M. T., 2014, *A&A*, 566, A118
- Duffau S., Zinn R., Vivas A. K., Carraro G., Méndez R. A., Winnick R., Gallart C., 2006, *ApJ*, 636, L97
- Gaia Collaboration, Brown A. G. A., Vallenari A., Prusti T., de Bruijne J. H. J., Babusiaux C., Bailer-Jones C. A. L., 2018, *ArXiv e-prints*
- Gaia Collaboration, Prusti T., de Bruijne J. H. J., Brown A. G. A. et al., 2016, *A&A*, 595, A1
- Hayes C. R., Majewski S. R., Hasselquist S., Beaton R. L. et al., 2018, *ApJ*, 859, L8
- Hendel D., Johnston K. V., 2015, *MNRAS*, 454, 2472
- Hildebrand A. R., Penfield G. T., Kring D. A., Pilkington M., Camargo Z. A., Jacobsen S. B., Boynton W. V., 1991, *Geology*, 19, 867
- Johnston K. V., 2016, in *Astrophysics and Space Science Library*, Vol. 420, *Tidal Streams in the Local Group and Beyond*, Newberg H. J., Carlin J. L., eds., p. 141
- Jurić M., Ivezić Ž., Brooks A., Lupton R. H. et al., 2008, *ApJ*, 673, 864

- Li T. S., Balbinot E., Mondrik N., Marshall J. L. et al., 2016, *ApJ*, 817, 135
- Majewski S. R., Ostheimer J. C., Rocha-Pinto H. J., Patterson R. J., Guhathakurta P., Reitzel D., 2004, *ApJ*, 615, 738
- Morgan J. V., Gulick S. P. S., Bralower T., Chenot E. et al., 2016, *Science*, 354, 878
- Myeong G. C., Evans N. W., Belokurov V., Sanders J. L., Koposov S. E., 2018a, *ApJ*, 856, L26
- Myeong G. C., Evans N. W., Belokurov V., Sanders J. L., Koposov S. E., 2018b, *ArXiv e-prints*
- Newberg H. J., Yanny B., Rockosi C., Grebel E. K. et al., 2002, *ApJ*, 569, 245
- Nie J. D., Smith M. C., Belokurov V., Fan X. H. et al., 2015, *ApJ*, 810, 153
- Price-Whelan A. M., Johnston K. V., Sheffield A. A., Laporte C. F. P., Sesar B., 2015, *MNRAS*, 452, 676
- Rocha-Pinto H. J., Majewski S. R., Skrutskie M. F., Crane J. D., Patterson R. J., 2004, *ApJ*, 615, 732
- Sanderson R. E., Helmi A., 2013, *MNRAS*, 435, 378
- Sesar B., Ivezić Ž., Lupton R. H., Jurić M. et al., 2007, *AJ*, 134, 2236
- Simion I. T., Belokurov V., Irwin M., Koposov S. E., 2014, *MNRAS*, 440, 161
- Simion I. T., Belokurov V., Koposov S. E., Sheffield A., Johnston K. V., 2018, *MNRAS*, 476, 3913
- Vanderplas J., Connolly A., Ivezić Ž., Gray A., 2012, in *Conference on Intelligent Data Understanding (CIDU)*, pp. 47–54
- Vivas A. K., Zinn R., 2006, *AJ*, 132, 714
- Vivas A. K., Zinn R., Andrews P., Baily C. et al., 2001, *ApJ*, 554, L33
- Vivas A. K., Zinn R., Farmer J., Duffau S., Ping Y., 2016, *ApJ*, 831, 165
- Watkins L. L., Evans N. W., Belokurov V., Smith M. C. et al., 2009, *MNRAS*, 398, 1757

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.