

Common origin for Hercules-Aquila and Virgo Clouds in Gaia DR2

Iulia T. Simion^{1★}, Vasily Belokurov^{2,3} and Sergey E. Koposov^{2,4}

¹*Key Laboratory for Research in Galaxies and Cosmology, Shanghai Astronomical Observatory, 80 Nandan Road, Shanghai 200030, China*

²*Institute of Astronomy, Madingley Rd, Cambridge, CB3 0HA*

³*Center for Computational Astrophysics, Flatiron Institute, 162 5th Avenue, New York, NY 10010, USA*

⁴*Department of Physics, McWilliams Center for Cosmology, Carnegie Mellon University, 5000 Forbes Avenue, Pittsburgh, PA 15213, USA*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We use a sample of ~500 RR Lyrae stars with radial velocities and Gaia DR2 proper motions to study orbital properties of the Hercules-Aquila Cloud (HAC) and Virgo Over-density (VOD). We demonstrate that both structures are dominated by stars on highly eccentric orbits, with pericentres around ~ 1 kpc and apocentres between 15 and 25 kpc from the Galactic centre. Given that the stars in the HAC and the VOD occupy very similar regions in the space spanned by integrals of motion, we conclude that these diffuse debris clouds are part of the same accretion event. More precisely, these inner halo sub-structures likely represent two complementary not-fully-mixed portions of an ancient massive merger, also known as the “sausage” event.

Key words: Galaxy: structure – Galaxy : formation – galaxies: individual: Milky Way.

1 INTRODUCTION

How do you hide the evidence for a massive impact event that caused the extinction of most of the dinosaurs as well as 75% of all species on Earth? You bury it deep under the sea, covered with a layer of sediment taller than the Empire State Building (Hildebrand et al. 1991). Without the discovery of the giant Chicxulub crater, the meteorite impact hypothesis would remain a neat theory supported by striking but indirect evidence. A hypothesis of an ancient dramatic collision between the Milky Way and an unidentified massive dwarf galaxy was put forward by Deason et al. (2013) to explain a particular feature in the overall stellar halo density profile (Watkins et al. 2009; Sesar et al. 2011; Deason et al. 2011). Most recently, through a study of a portion of the nearby stellar halo, Belokurov et al. (2018b) demonstrated that the great impactor must have collided with the young Milky Way on a nearly radial orbit, thus swamping the inner stellar halo with metal-rich material with orbital anisotropy (see Binney & Tremaine 2008) close to unity. Merger events like this tend to leave behind a battery of debris clouds and shells (see e.g. Johnston et al. 2008; Amorisco 2015; Hendel & Johnston 2015), which - akin to the peak rings of impact craters (see e.g. Morgan et al. 2016) - if discovered could help to reconstruct the collision as well as pin down the properties of the progenitor (e.g. Sanderson & Helmi 2013; Johnston

2016). Before the Data Release 2 (Gaia Collaboration et al. 2018) of the ESA’s Gaia mission (Gaia Collaboration et al. 2016), five large and diffuse cloud-like structures had been discovered in the Galaxy’s halo. These include: the Virgo Over-Density (VOD, Vivas et al. 2001; Newberg et al. 2002; Duffau et al. 2006; Jurić et al. 2008; Bonaca et al. 2012), the Hercules-Aquila Cloud (HAC, Belokurov et al. 2007; Simion et al. 2014), the Tringulum-Andromeda structure (Tri-And, Rocha-Pinto et al. 2004; Majewski et al. 2004; Deason et al. 2014), the Pisces Over-density (Sesar et al. 2007; Watkins et al. 2009; Nie et al. 2015) and the Eridanus-Phoenix over-density (Eri-Pho, Li et al. 2016). According to the most recent investigations, Tri-And likely comprises of Galactic disc stars kicked out of the plane in a recent interaction with a dwarf galaxy, probably the Sagittarius dSph (e.g. Price-Whelan et al. 2015; Bergemann et al. 2018; Hayes et al. 2018). Of the remaining four, the Pisces overdensity clearly stands out as it reaches much larger Galacto-centric distances. On the other hand, the VOD, HAC and Eri-Pho structures occupy a very similar range of distances, between 10 and 20 kpc from the Galactic center. This led Li et al. (2016) to suggest that these three Clouds could all be part of one merger event, a galaxy accreted onto the Milky Way on a polar orbit (see also Jurić et al. 2008). As demonstrated by the recent re-interpretation of the Monoceros Ring (and the associated sub-structures) and the Tri-And, deciphering the nature of halo over-densities is often impossible without either high-resolution spectroscopy (e.g. Bergemann et al.

★ E-mail: isimion@shao.ac.cn



Figure 1. Spatial distribution of ~ 350 RR Lyrae with 6-D phase space information in the HAC and VOD fields, in Galactic coordinates (left panel) and in the x-y (middle) and y-z (right) planes. In green we mark a significant kinematical group identified by Vivas et al. (2016) (group 2 in their table 5) and in red a sub-sample of the VOD RRL with galactocentric distances similar to the HAC sample, $11 < r_{GC} < 16$. The semi-circles are centred on the Sun's position and have cylindrical radius of 10 and 20 kpc. The Sun (yellow star) is located at $(x_{\odot}, y_{\odot}, z_{\odot}) = (-8, 0, 0)$ kpc and the Galactic centre at $(0, 0, 0)$ kpc.

2018) or accurate astrometry (e.g. de Boer et al. 2018; Deason et al. 2018a). In this Letter, we look for clues to the formation of the Hercules-Aquila and Virgo Clouds using proper motions provided as part of the Gaia DR2. At our disposal are highly pure samples of members of each Cloud, namely the RR Lyrae stars that i) are co-spatial with HAC and VOD in 3-D and ii) that have their line-of-sight velocities measured. By complementing the publicly available 4-D data with the GDR2 proper motions, we build a large tracer set with complete 6-D phase space information and study the make-up of each structure using the orbital properties of the constituent stars.

2 DATA AND ANALYSIS

2.1 4-D RR Lyrae data

The Hercules-Aquila and Virgo Clouds are diffuse stellar over-densities in the inner stellar halo, located on the opposite sides of the Galaxy (see Figure 1). At high latitudes, these are detected as peaks in RR Lyrae number counts - curiously - at similar heliocentric distances, i.e. ~ 17 kpc (HAC: Watkins et al. 2009; Simion et al. 2014) and ~ 19 kpc (VOD: Vivas & Zinn 2006; Duffau et al. 2014; Vivas et al. 2016). Note that other tracers (e.g. BHBs, MSTO and K and M giants) have also been used to pin down the morphology of the Clouds (see e.g. Belokurov et al. 2007; Jurić et al. 2008; Sharma et al. 2010; Bonaca et al. 2012; Conroy et al. 2018). The RR Lyrae, however offer the clearest view of these halo sub-structures thanks to their associated accurate distances and minuscule Galactic foreground contamination. Therefore, in this work, we have focused on the two recently published samples of RR Lyrae towards the Clouds, where each star has a well-measured line-of-sight velocity.

- Simion et al. (2018) provide a table of 46 RRL with radial velocity measurements (45 observed with MDM and 1 from SDSS) with heliocentric distances between 15 and 18 kpc;
- Vivas et al. (2016) compiled a catalog of 412 RRL in the region of the sky covered by the VOD with distances between 4 and 75 kpc from the Sun with radial velocity measure-

ments of stars from La Silla-QUEST, QUEST, CRTS and LINEAR.

2.2 From 4-D to 6-D. Velocity distributions

By cross-matching to the GDR2 data with an aperture of $2''$, we have found Gaia counterparts to 44 HAC stars and 411 VOD. From the VOD sample, we remove 112 stars likely belonging to the Sgr stream as identified by (Vivas et al. 2016, their group 1). The spatial distribution of the remaining stars (44 from Simion et al. 2018 and 299 from Vivas et al. 2016) with full 6-D phase space measurements is given in Figure 1, in Galactic coordinates in the left panel and in the x-y (x-z) Galactic plane in the middle (right) panel. We adopt a left-handed Galactic Cartesian coordinates with the Sun located at $(x_{\odot}, y_{\odot}, z_{\odot}) = (-8, 0, 0)$ kpc, the x-axis positive in the direction of the Galactic center, y-axis oriented along the Galactic rotation and the z-axis directed towards the north Galactic pole. While Vivas et al. 2016 identify 6 significant kinematical groups in the VOD region (their table 5), only groups 1 (Sagittarius stream) and 2 (likely members of the VOD, with $\langle v_{GSR} \rangle = 135$ km/s) contain more than 10 stars. We mark group 2 with green circles in Figure 1. We also mark with red circles the location of a group of stars selected to have galactocentric distances similar to the HAC sample, i.e. $11 < r_{GC} < 16$ to facilitate a fair comparison of their velocities and orbits. Figure 2 shows the spherical polar components of the velocities (radial v_r , azimuthal v_{θ} and polar v_{ϕ}) of stars in the HAC (top row) and VOD (bottom row) fields. Group 2 stars (green) can be seen clustering at $v_r = 135$ km/s (by design), while the stars at intermediate r_{GC} (red) seem to have a velocity distribution very similar to those in the HAC, shown in the top row. To estimate the uncertainty in each velocity component we propagate the measured proper motion and line-of-sight velocity errors using Monte-Carlo re-sampling, where we take into account the covariances between the measurements of the Right Ascension and Declination components of proper motion as provided in GDR2. We use the standard deviations of the resulting $\{v_r, v_{\theta}, v_{\phi}\}$ distributions as the upper limits of the velocity uncertainties; these are shown in Fig. 2. As evident from the Figure, the velocity distributions are highly anisotropic, with the dispersion in the radial component dominating the tangential ones, especially in the HAC data. To describe the shape of the velocity distributions, we model each stellar sample as a single-component multivariate Gaussian using the Extreme Deconvolution module implemented in astroML (Vanderplas et al. 2012). The resulting parameters and the associated uncertainties of the velocity ellipsoids are taken to be the median and the standard deviation values of 500 bootstrap re-sampling trials. The velocity ellipsoid shape can be summarized using the anisotropy parameter $\beta = 1 - (\sigma_{\theta}^2 + \sigma_{\phi}^2)/2\sigma_r^2$ (see Binney & Tremaine 2008). We find the HAC stars have radially biased orbits with $\beta = 0.91 \pm 0.03$ while for the whole of the VOD sample, $\beta = 0.74 \pm 0.04$. Note, however, that the two samples span very different ranges in Galactic l, b and distances. According to Belokurov et al. (2018b), the inner stellar halo can be viewed as a mixture of two debris components with distinct properties. Accordingly, we fit a model with two multivariate Gaussians to the VOD velocity data




Figure 2. Velocity distribution in spherical polar coordinates (v_r , v_θ , v_ϕ are the radial, azimuthal and polar components respectively, in km/s) of the HAC (top panels) and VOD (bottom panels) RR Lyrae. The label for the y-axis is reported in the upper-left corner of each panel. The velocity uncertainties were estimated through error propagation. Both fields have radially biased orbits with anisotropy parameters $\beta = 0.91 \pm 0.03$ (HAC) and $\beta = 0.74 \pm 0.04$ (VOD). A mixture of two multivariate Gaussians was fitted to the VOD velocity ellipsoid: 62% of stars seem to belong to the $\beta_1 = 0.96^{+0.02}_{-0.44}$ (blue) component and 38% to the $\beta_2 = 0.44^{+0.45}_{-0.20}$ (magenta) component. The Vivas Gr. 2 and VOD (11-16) groups are also shown in lime and red, as in Fig.1.

using Extreme Deconvolution (see Bovy et al. 2011). The VOD sample appears to be composed of roughly two thirds of stars with $\beta_1 = 0.96^{+0.02}_{-0.44}$ (marked in blue in Fig. 2) and a third with $\beta_2 = 0.44^{+0.45}_{-0.20}$ (magenta) in good agreement with the results for the local halo presented in Belokurov et al. (2018b). Using virial theorem, Myeong et al. (2018b) concluded that the radially anisotropic component of the stellar halo is also significantly flattened vertically. To test this hypothesis, we split the VOD sample into 3 groups according to their distance from the Galactic plane and show the behaviour of the azimuthal v_θ and radial v_r velocity distributions in Figure 3. Additionally, for each z slice we have calculated the fraction of Oosterhoff type I (Oo I) RR Lyrae, using equations 1 and 2 in Belokurov et al. (2018a) to explore the changes in the RRL mixture. In the $10 < z/\text{kpc} < 20$ range, where the velocity anisotropy is the highest ($\beta = 0.84 \pm 0.03$) approaching the value in the HAC field, the Oo I type dominates with $(77 \pm 13)\%$. In the same slice, 73% of the stars belong to the more squashed (or ‘sausage’ looking) velocity ellipsoid. The same behaviour but less accentuated can be noticed in the $0 < z/\text{kpc} < 10$ slice where the anisotropy $\beta = 0.7 \pm 0.1$ is lower but the fraction of Oo I stars is somewhat lower at $(57 \pm 21)\%$. Further from the plane, at $z > 20$ kpc, the velocity ellipsoid changes dramatically to almost isotropic with $\beta = -0.1 \pm 0.2$ and Oo I type stars fraction of $(64 \pm 13)\%$. Please note that in this particular z bin, the β value may still be affected by the presence of the Sagittarius stream. Group 2 and the stars sharing the same galactocentric distance range with the HAC, are all located at $z < 20$ kpc. Interestingly, the HAC counterparts in the VOD (red points) are all clearly part of the anisotropic component. To summarize, it is clear that the velocity ellipsoid shape does




Figure 3. Velocity distribution of the VOD RR Lyrae, in three distance slices above the Galactic plane (see also the bottom left panel of Fig. 2). The fraction of Oosterhoff type I RR Lyrae is reported for each slice, along with the orbital anisotropy β . Note that in the middle panel where $10 < z (\text{kpc}) < 20$, the anisotropy parameter is $\beta = 0.84 \pm 0.03$, value close to the one reported for HAC (top panels of Fig. 2); in the same field the large majority of RRL is of Oosterhoff type I.

change noticeably with vertical height. However, given the limited range of Galactic l and b in the VOD sample, it is not possible to conclude whether the structure is flattened or simply ends around $z \sim 20$ kpc.

2.3 Orbital Properties

We use the 6-D measurements described above to initialise the RR Lyrae orbits in the HAC and VOD fields. The orbits are integrated using the galpy package (Bovy 2015) in the recommended Galactic potential model for the Milky Way, MWPotential2014 composed of a Miyamoto-Nagai disc, a bulge with a power-law density profile that is exponentially cut-off and a dark matter halo described by a NFW potential, with parameters given in Table 1 of Bovy (2015). The orbital properties, i.e. the peri- and apo-centric radii, eccentricity, and the maximal height above the disc plane, are shown in Figure 4. To compute the errors (not shown for VOD to simplify the figure) we integrate 500 orbits for each star where the initial conditions are re-sampled from the observations according to the associated uncertainties. The probability distributions of the eccentricities, apo-centres and peri-centres are shown in the bottom row of the Figure. Note that the default Milky Way mass in galpy, $M_{\text{vir}} = 0.8 \times 10^{12} M_\odot$, is somewhat lower than suggested by recent measurements. We have checked however that the distributions of the orbital parameters displayed in Figure 4 are minimally affected if we increase the Galaxy’s mass.

The stars in the HAC field typically travel as high as ~ 15 kpc above the Galactic plane. The apo-centres bunch up around 15-20 kpc from the Galactic center, with a tail to higher values. The distribution of the peri-centric distances of the HAC stars peaks sharply around ~ 1 kpc with most stars having their per-centers within 5 kpc from the Galactic center. This naturally implies highly eccentric orbits (with eccentricity close to 1) for the vast majority of the HAC stars. Compared to the HAC, the VOD stars explore broader range of apo-centres, with the distribution of the peri-centres much less strongly peaked. As a result, the typical eccentricity for a VOD star is around ~ 0.8 . Note however, that a sample of VOD stars comprised of Group 2 (shown in green) and stars that share instantaneous Galactocentric distances with objects in the HAC field appears to have orbital properties much closer to that of HAC. In particular, once the




Figure 4. Orbital properties of the RR Lyrae orbits in the HAC and VOD. In the top and middle rows we show the eccentricity, maximal height above the orbit plane, apocentre and Galactocentric radius for the HAC (top) and VOD (middle). In the bottom row we show the probability distributions of the eccentricities, pericentres and apocentres for both fields, including for the VOD subsample with Galactocentric distances $11 < r_{\text{GC}}(\text{kpc}) < 16$ (in red).




Figure 5. Distribution of the L_{\perp} and L_z components of the angular momentum for the HAC (left panel) and VOD (middle panel). The right panel shows the distribution of the total angular momentum as a function of energy for all the stars in the HAC and VOD fields.

stars with similar distances are selected, the apo-centre and peri-center distance distributions in the two fields look very similar. The similarity of orbital properties of the HAC and VOD stars is emphasized in Figure 5. Here, we first show the current Galacto-centric radii and speeds of the stars in both fields (left panel). Middle panel gives the distribution of total angular momentum as a function of energy for stars in the HAC (black) and stars in the VOD with matching distances ($11 < r_{\text{GC}}/\text{kpc} < 16$, red). The bulk of the stellar debris in both fields are on high energy, low angular momentum orbits. Finally, the right panel of the Figure presents the distribution of the components of the angular momentum L_z and L_{\perp} . As evident from this panel, while stars in both HAC and VOD occupy highly radial orbits, both prograde and retrograde objects exist in both fields, with slight prevalence of the retrograde ones (see also Myeong et al. 2018b,a; Helmi et al. 2018). Figure 6 provides an alternative view of the orbital properties of the HAC and VOD stars. Here we have used the stellar 6-D phase-space measurements as initial condi-




Figure 6. Backward orbit integration for HAC (top panels) and VOD (middle panels) for 8 Gyrs. We show a density plot of the test particle positions along their individual orbits across the past 8 Gyrs, in Galactic coordinates (left panels) and in the x-y and y-z Galactic planes. In the bottom panels we show the VOD orbits only for the ‘sausage’ component. It is useful to compare this figure obtained from actual observations to a simulated accretion event shown in figure 7 of Simion et al. (2018): they look very similar.

tions and integrated the orbits back in time for 8 Gyrs. The Figure shows the density of the test particle positions along each orbits for all orbits across the entire temporal range of the integration. The HAC stars occupy a slightly flattened (in the vertical direction) cross-like (in the y-z plane and its projection on the sky - l, b plane) structure with a density peak at the Galactic centre. Note that some of the appearance of the debris distribution could be due to the limited initial conditions of the stars with available radial velocity measurements. The VOD stars typically move through very similar regions of the Galaxy (especially the distance-matched sub-sample) shown in the bottom row. While, overall, the match between the stellar debris distributions in Figure 6 is striking, the HAC and VOD stars studied here do not have identical orbital properties. For example, the VOD stars travel further above the Galactic disc, as exhibited by much rounder, or perhaps even vertically stretched, y-z distribution (right panels). Notwithstanding possible selection effects present in the data, the stellar density distribution shown in Figure 6 looks staggeringly similar to the debris distribution of the simulated accretion event presented in Figure 7 of Simion et al. (2018). This particular merger happened some 11 Gyr ago according to the suite of numerical stellar halo formation models of Bullock & Johnston (2005). Giving the conspicuous similarity in the appearance of the orbital density distributions of the HAC and the VOD stars and the simulated merger example, we conclude that the Hercules-Aquila Cloud and the Virgo Overdensity are both parts of one ancient massive head-on collision.

3 CONCLUSIONS

Using a sample of ~ 500 RR Lyrae with full 6-D phase space information, we have studied the orbital properties of the Hercules-Aquila and Virgo Clouds. Both Clouds appear

dominated by stars on highly eccentric orbits. Assuming that the kinematics of each structure is well described by a single Gaussian, the orbital anisotropy of the HAC is $\beta = 0.91$ and for the VOD, $\beta = 0.74$. Note, however that the original criteria applied to the CRTS RR Lyrae dataset to select targets for spectroscopic follow-up differ drastically between the HAC and the VOD datasets. The HAC sample covers a very limited region in the of l, b, D space, while the VOD dataset spans a wide range of longitudes, latitudes and heliocentric distances. It is therefore likely that the VOD dataset contains a mixture of several halo sub-structures (see Vivas et al. 2016, for a detailed discussion). For the entirety of the analysis described here, we made sure to cull the probable Sgr stream members. Additionally, we explore how the VOD’s make-up changes with Galactic height and demonstrate that for $|z| < 20$ kpc, the VOD orbital anisotropy is $\beta \sim 0.8$, while above this threshold, it quickly changes to $\beta \sim 0$. We conclude therefore, that an assumption of a single Gaussian for the entire VOD sample is likely not appropriate. Modeling the kinematics of the VOD stars with a mixture of 2 multi-variate Gaussians, we show that two thirds of the VOD sample are the stars with $\beta = 0.96$, while the remainder has $\beta = 0.44$, in good agreement with the local measurement presented in Belokurov et al. (2018b).

As revealed by Gaia, the two structures are composed of stars on nearly radial orbits, with peaks in the eccentricity distribution at 0.95 (0.8) for the HAC (VOD). The distributions of the peri-centric and apo-centric distances also match: the stars in the Clouds turn around at 1 – 2 and 15 – 25 kpc. Not only the HAC and the VOD look alike kinematically, their orbital composition is in perfect agreement with the stellar halo properties as analysed locally by Belokurov et al. (2018b) and globally (out to 40 kpc) by Deason et al. (2018b). As these authors demonstrate, the inner halo is dominated by metal-rich debris from an old and massive accretion event. In particular, Belokurov et al. (2018b) use Cosmological simulations of the Milky Way halo formation, to bracket the time of the merger - between 8 and 11 Gyr ago - and its mass, which they show to be in excess of $10^{10} M_{\odot}$. The tell-tale sign of this dramatic head-on collision is the particular shape of the corresponding stellar velocity ellipsoid, which is stretched so much in the radial direction (compared to the tangential ones), that it resembles a sausage. An alternative view of this merger can be found in Myeong et al. (2018b), where the local stellar halo is mapped out in the action space. Here, the metal-rich stars are shown to have an extended radial action distribution in addition to a prominent spray of material on retrograde orbits. The high mass of the progenitor is evidenced not only by the metallicity distribution of its likely members or the numerical simulations of halo formation, but also by a sizeable number of Globular Clusters that could be attributed to the same event (see Myeong et al. 2018c; Kruijssen et al. 2018).

Our interpretation of the nature of the HAC and VOD is in broad agreement with the earlier studies of Johnston et al. (2012) and Carlin et al. (2012) whereby each group of authors have singled out a progenitor on a high eccentricity orbit as a culprit for the production of HAC and VOD respectively. However, for the first time, we connect both of the discussed debris Clouds to a single event with yet higher eccentricity and yet larger mass. Additionally, rather than

being a recent accretion, the “sausage” merger likely happened around the epoch of the Galactic disc formation, i.e. 8 to 11 Gyrs ago (see Belokurov et al. 2018b).

ACKNOWLEDGEMENTS

The authors wish to thank Alis Deason for insightful comments that helped to improve the quality of the manuscript. The research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement n. 308024.

REFERENCES

- Amorisco N. C., 2015, MNRAS, 450, 575
- Belokurov V., Deason A. J., Koposov S. E., Catelan M., Erkal D., Drake A. J., Evans N. W., 2018a, MNRAS, 477, 1472
- Belokurov V., Erkal D., Evans N. W., Koposov S. E., Deason A. J., 2018b, MNRAS, 478, 611
- Belokurov V., Evans N. W., Bell E. F., Irwin M. J. et al., 2007, ApJ, 657, L89
- Bergemann M., Sesar B., Cohen J. G., Serenelli A. M. et al., 2018, Nature, 555, 334
- Binney J., Tremaine S., 2008, Galactic Dynamics: Second Edition. Princeton University Press
- Bonaca A., Jurić M., Ivezić Ž., Bizyaev D. et al., 2012, AJ, 143, 105
- Bovy J., 2015, ApJS, 216, 29
- Bovy J., Hogg D. W., Roweis S. T., 2011, Annals of Applied Statistics, 5
- Bullock J. S., Johnston K. V., 2005, ApJ, 635, 931
- Carlin J. L., Yam W., Casetti-Dinescu D. I., Willett B. A., Newberg H. J., Majewski S. R., Girard T. M., 2012, ApJ, 753, 145
- Conroy C., Bonaca A., Naidu R. P., Eisenstein D. J., Johnson B. D., Dotter A., Finkbeiner D. P., 2018, ArXiv e-prints
- de Boer T. J. L., Belokurov V., Koposov S. E., 2018, MNRAS, 473, 647
- Deason A. J., Belokurov V., Evans N. W., 2011, MNRAS, 416, 2903
- Deason A. J., Belokurov V., Evans N. W., Johnston K. V., 2013, ApJ, 763, 113
- Deason A. J., Belokurov V., Hamren K. M., Koposov S. E. et al., 2014, MNRAS, 444, 3975
- Deason A. J., Belokurov V., Koposov S. E., 2018a, MNRAS, 473, 2428
- Deason A. J., Belokurov V., Koposov S. E., Lancaster L., 2018b, ArXiv e-prints
- Duffau S., Vivas A. K., Zinn R., Méndez R. A., Ruiz M. T., 2014, A&A, 566, A118
- Duffau S., Zinn R., Vivas A. K., Carraro G., Méndez R. A., Winnick R., Gallart C., 2006, ApJ, 636, L97
- Gaia Collaboration, Brown A. G. A., Vallenari A., Prusti T., de Bruijne J. H. J., Babusiaux C., Bailer-Jones C. A. L., 2018, ArXiv e-prints
- Gaia Collaboration, Prusti T., de Bruijne J. H. J., Brown A. G. A. et al., 2016, A&A, 595, A1
- Hayes C. R., Majewski S. R., Hasselquist S., Beaton R. L. et al., 2018, ApJ, 859, L8
- Helmi A., Babusiaux C., Koppelman H. H., Massari D., Veljanoski J., Brown A. G. A., 2018, ArXiv e-prints
- Hendel D., Johnston K. V., 2015, MNRAS, 454, 2472

- Hildebrand A. R., Penfield G. T., Kring D. A., Pilkington M., Camargo Z. A., Jacobsen S. B., Boynton W. V., 1991, *Geology*, 19, 867
- Johnston K. V., 2016, in *Astrophysics and Space Science Library*, Vol. 420, *Tidal Streams in the Local Group and Beyond*, Newberg H. J., Carlin J. L., eds., p. 141
- Johnston K. V., Bullock J. S., Sharma S., Font A., Robertson B. E., Leitner S. N., 2008, *ApJ*, 689, 936
- Johnston K. V., Sheffield A. A., Majewski S. R., Sharma S., Rocha-Pinto H. J., 2012, *ApJ*, 760, 95
- Jurić M., Ivezić Ž., Brooks A., Lupton R. H. et al., 2008, *ApJ*, 673, 864
- Kruijssen J. M. D., Pfeffer J. L., Reina-Campos M., Crain R. A., Bastian N., 2018, *MNRAS*
- Li T. S., Balbinot E., Mondrik N., Marshall J. L. et al., 2016, *ApJ*, 817, 135
- Majewski S. R., Ostheimer J. C., Rocha-Pinto H. J., Patterson R. J., Guhathakurta P., Reitzel D., 2004, *ApJ*, 615, 738
- Morgan J. V., Gulick S. P. S., Bralower T., Chenot E. et al., 2016, *Science*, 354, 878
- Myeong G. C., Evans N. W., Belokurov V., Sanders J. L., Koposov S. E., 2018a, *MNRAS*
- Myeong G. C., Evans N. W., Belokurov V., Sanders J. L., Koposov S. E., 2018b, *ApJ*, 856, L26
- Myeong G. C., Evans N. W., Belokurov V., Sanders J. L., Koposov S. E., 2018c, *ArXiv e-prints*
- Newberg H. J., Yanny B., Rockosi C., Grebel E. K. et al., 2002, *ApJ*, 569, 245
- Nie J. D., Smith M. C., Belokurov V., Fan X. H. et al., 2015, *ApJ*, 810, 153
- Price-Whelan A. M., Johnston K. V., Sheffield A. A., Laporte C. F. P., Sesar B., 2015, *MNRAS*, 452, 676
- Rocha-Pinto H. J., Majewski S. R., Skrutskie M. F., Crane J. D., Patterson R. J., 2004, *ApJ*, 615, 732
- Sanderson R. E., Helmi A., 2013, *MNRAS*, 435, 378
- Sesar B., Ivezić Ž., Lupton R. H., Jurić M. et al., 2007, *AJ*, 134, 2236
- Sesar B., Jurić M., Ivezić Ž., 2011, *ApJ*, 731, 4
- Sharma S., Johnston K. V., Majewski S. R., Muñoz R. R., Carlberg J. K., Bullock J., 2010, *ApJ*, 722, 750
- Simion I. T., Belokurov V., Irwin M., Koposov S. E., 2014, *MNRAS*, 440, 161
- Simion I. T., Belokurov V., Koposov S. E., Sheffield A., Johnston K. V., 2018, *MNRAS*, 476, 3913
- Vanderplas J., Connolly A., Ivezić Ž., Gray A., 2012, in *Conference on Intelligent Data Understanding (CIDU)*, pp. 47–54
- Vivas A. K., Zinn R., 2006, *AJ*, 132, 714
- Vivas A. K., Zinn R., Andrews P., Baily C. et al., 2001, *ApJ*, 554, L33
- Vivas A. K., Zinn R., Farmer J., Duffau S., Ping Y., 2016, *ApJ*, 831, 165
- Watkins L. L., Evans N. W., Belokurov V., Smith M. C. et al., 2009, *MNRAS*, 398, 1757

This paper has been typeset from a \LaTeX file prepared by the author.