

Appendix 3: FKB327-003 protocol outline

CLINICAL PROTOCOL

**An Open-label Extension Study to Compare the Long-term Efficacy, Safety,
Immunogenicity and Pharmacokinetics of FKB327 and Humira® in Patients with
Rheumatoid Arthritis on Concomitant Methotrexate (ARABESC-OLE)**

PROTOCOL SUMMARY

Study Title:

An Open-label Extension Study to Compare the Long-term Efficacy, Safety, Immunogenicity and Pharmacokinetics of FKB327 and Humira® in Patients with Rheumatoid Arthritis on Concomitant Methotrexate (ARABESC-OLE)

Study Number:

FKB327-003

Protocol Acronym:

ARABESC-OLE

Name of Investigational Product:

FKB327

Name of Active Ingredient:

adalimumab

Study Phase:

Phase 3

Number of Patients Planned:

up to 600 patients

Number of Sites Planned:

approximately 120 - the sites will be the same as for study FKB327-002

Study Population:

Only patients who completed the double-blind study FKB327-002 will be permitted to continue into this extension study. These will be patients with rheumatoid arthritis (RA) who are taking concomitant methotrexate (MTX) and folate

Primary Objective:

The primary objective of this study is to compare the safety of long-term treatment with FKB327 and Humira in patients with RA

Secondary Objectives: The secondary objectives of this study are as follows:

- To compare the efficacy of long-term treatment with FKB327 and Humira in patients with RA
- To compare the proportion of patients developing anti-drug antibodies (ADAs) on long-term treatment with FKB327 and Humira in patients with RA
- To compare the pharmacokinetics (PK) of long-term treatment with FKB327 and Humira in patients with RA
- To evaluate safety, changes in efficacy, and changes in PK and immunogenicity in patients who are switched from Humira in the preceding FKB327-002 double-blind study to FKB327 in the FKB327-003 OLE study, and of patients who are switched from FKB327 to Humira, respectively
- To evaluate safety, changes in efficacy, and changes in PK and immunogenicity in patients who are switched from FKB327 in the preceding FKB327-002 double-blind study to

Humira in the FKB327-003 OLE study, and then switch back to FKB327 in the second part of the FKB327-003 OLE study

Key Inclusion Criteria:

1. Patient has completed the Week 24 visit procedures of Study FKB327-002 and are continuing with methotrexate
2. In the investigator's opinion, the patient showed a clinical response to treatment during Study FKB327-002

Key Exclusion Criteria:

1. Patient has evidence of a serious adverse event (SAE) ongoing from Study FKB327-002
2. Patient has presence of active and/or untreated latent tuberculosis (TB)

Study Drug:

Patients will receive subcutaneous (sc) FKB327 40 mg every other week (eow) or sc Humira 40 mg eow from Week 0 to Week 28; after this, all patients will receive FKB327 40 mg eow from Week 30 to Week 76. Patients will continue to take the same stable dose of MTX (10-25 mg/week) throughout this study as taken in Study FKB327-002, with folic/folinic acid at a dose of at least 5 mg/week.

Blinding and Randomisation:

This extension study is open-label throughout. Patients from Study FKB327-002 will be re-randomised to receive either FKB327 40 mg eow or Humira 40 mg eow from Week 0 to Week 28. Patients who received FKB327 in Study FKB327-002 will be re-randomised to receive FKB327 or Humira in a 2:1 ratio. Patients who received Humira in Study FKB327-002 will be re-randomised to receive Humira or FKB327 in a 2:1 ratio. From Week 30 to Week 76, all patients will receive FKB327 40 mg eow in an open fashion.

Study Design and Methodology:

The first part of this study is an open-label, randomised, comparative, multi-centre, 2-arm extension Phase 3 study in patients with RA who are taking MTX who will continue from the preceding Study FKB327-002. Patients who complete Study FKB327-002 and consent to this open-label extension study will be re-randomised to receive treatment with either FKB327 40 mg eow or Humira 40 mg eow from Week 0 to Week 28. The second part of the study is an open-label, multi-centre, single arm extension in which all patients from the first part may receive prolonged FKB327 treatment from Week 30 to Week 76, followed by a 4-week follow-up period. Scheduled clinic visits will be conducted at Weeks 0, 2, 4, 8, 12, 24, 30, 32, 34, 42, 54, 66, 76, and 80. The patient or carer will administer interim doses of study drug at home (eow) between clinic visits.

Primary Efficacy Endpoint:

- As the primary endpoint is safety, there is no primary efficacy endpoint in this study

Secondary Efficacy Endpoint:

- ACR20, ACR50 and ACR70 response rates from baseline (Week 0 from Study FKB327-002)
- Changes in DAS28-CRP score compared to baseline (Week 0 from Study FKB327-002)

Other Endpoints:

- Immunogenicity.
- PK

Safety Endpoints:

- Adverse events (AEs).
- SAEs, including serious infections and malignancies.
- Vital signs.
- Clinical laboratory tests.

Assessment**Assessment of Efficacy**

- ACR20, ACR50 and ACR70
- DAS28

Assessment of Safety

- Clinical Laboratory Tests
- Physical Examination
- Vital Signs
- Chest X-ray
- Injection Site Assessment
- Immunogenicity Assessments
- Pharmacokinetic Assessments

Statistical Methods for primary and secondary endpoints:

Safety endpoints: AEs will be summarised by System Organ Class (SOC) and Preferred Term (PT) using the Medical Dictionary for Regulatory Activities (MedDRA). AEs will be tabulated by treatment sequence and relationship to treatment. SAEs, serious infections, and malignancies will be summarised by treatment sequence. Safety data including vital signs, haematology, serum chemistry and urinalysis will be summarised by treatment sequence for each visit.

The category and size of any injection site reaction will be summarised by open-label treatment and visit using descriptive statistics.

Efficacy endpoints: ACR20, ACR50 and ACR70 response rates from Baseline (Week 0) in Study FKB327-002 will be summarised by treatment sequence and visit using observed cases. DAS28 score and the change from Baseline (Week 0 in the FKB327-002 study) in DAS28 score will be summarised by treatment sequence and visit using observed cases.

SCHEDULE OF EVENTS

Table 1: Study Schedule - Randomised Treatment Period

Study Visit	Treatment Period									
Week	0	2	4	6	8	10	12	14-22	24	26 and 28
Informed consent	•									
Medical history	•									
Re-randomisation	•									
Study medication administration (mg)	40	40	40	40	40	40	40	40 (eow)	40	40 (eow)
Concomitant MTX and folate										
Physical examination, Vital signs, ECG, Chest X-ray, Laboratory testing including CRP, ESR are assessed at the schedule defined by protocol										
Serum drug concentration	(•)						•		•	
ADA test	(•)						•		•	
Injection site assessment (including injection site pain VAS)	•									
Tender/Swollen Joint Counts	(•)		•		•		•		•	
Patient global assessment	(•)		•		•		•		•	
Patient pain VAS	(•)		•		•		•		•	
HAQ-DI	(•)		•		•		•		•	
Physician global assessment	(•)		•		•		•		•	
Adverse events										

Table 2: Study Schedule - Follow-up Treatment Period

Study Visit	Treatment Period										
Week	30	32 & 34	36-40	42	44-52	54	56-64	66	68-74	76	80 or EOS
FKB327 administration (mg)	40	40	40 (eow)	40	40 (eow)	40	40 (eow)	40	40 (eow)	40	
Concomitant MTX and folate											→
Physical examination, Vital signs, ECG, Chest X-ray, Laboratory testing including CRP, ESR assessed at schedule defined by protocol											→
Serum drug concentration	•					•				•	•
ADA test	•					•				•	•
Injection site assessment (including injection site pain VAS)	•										
Assessment of tender/swollen joints (68/66 joints)	•			•		•		•		•	•
Patient's global assessment	•			•		•		•		•	•
Patient's pain VAS	•			•		•		•		•	•
HAQ-DI	•			•		•		•		•	•
Physician's global assessment	•			•		•		•		•	•
Adverse events											→