

DEVELOPMENT OF PAPILLOMAVIRUS INFECTION IN MEN



Islamov Nurali Hikmatovich Samarkand State Medical University
Department of Skin and Genital Organ Diseases

Jalilov Asliddin Xolmatovich
Samarkand State Medical University Department of Propriedevology of
Childhood Diseases

Achilova Feruza Axtamovna
Samarkand State Medical University Department of Propriedevology of
Childhood Diseases

Annotation: This article summarizes a study of 38 couples who continued uninterrupted sexual life for at least 3 years, in which partners were carriers of oncogenic types of HPV. DNA of oncogenic HPV strains in the studied samples was detected by real-time PCR. work consisted in determining the frequency of infection and describing the clinical manifestations of PVI in men from married couples with an infected partner.

Keywords: papillomavirus infection, couples, clinical manifestations

Introduction: Human papillomaviruses (HPV), or human papillomaviruses (HPV, English Human papillomavirus, HPV) are a group of viruses from the papillomavirus family, including 27 species from 5 genera (Alphapapillomavirus, Betapapillomavirus, Gammapapillomavirus, Mupapillomavirus and Nupapillomavirus) and more than 170 types (strains).

About 80% of the sexually active population becomes infected with HPV during their lifetime. 660 million people in the world are already infected with HPV (12% of the world's population). 750 thousand cases of HPV-associated cancers and 32 million cases of urogenital warts are registered annually in the world. More than 300,000 people die every year from cancers caused by HPV. HPV is the most common sexually transmitted disease in the United States, Russia and many other countries.

In itself, HPV infection does not mean evidence of sexual contact, since the standard life cycle of the virus is designed for infection through microcracks in the skin. Infection through the mucous membranes is only more effective and therefore more

common. Infection with the virus is possible through shared objects or even a handshake. Infection through objects and the skin of other people contributes to the extreme resistance of HPV to antiseptics.

Papillomavirus infection (PVI) is one of the most common viral infections. Its high oncogenic risk strains are currently attracting considerable attention. Their frequency and clinical manifestations are well studied in women, but are rarely described in men. One of the reasons for insufficient attention to them may be the latent, subclinical nature of the course of infection, when pathological changes are present and located in the thickness of the epithelium, however, it may be impossible to visualize them with the naked eye. At the same time, the man is a source of infection for the partner and remains at risk of malignancy of the process.

The aim of the work was to determine the frequency of infection and describe the clinical manifestations of PVI in men from married couples with an infected partner.

Materials and methods. The object of the study were 38 married couples who continued uninterrupted sexual life for at least 3 years, in which partners were carriers of oncogenic types of HPV. DNA of oncogenic HPV strains in the studied samples was detected by real-time PCR. The study of the clinical manifestations of PVI caused by oncogenic HPV types in men, in addition to visual examination of the anogenital region, body skin, glans penis and urethral sponges, included simple and extended peniscopy of the glans skin and coronal sulcus. Statistical data analysis was performed using the Statistica for Windows 6.0 program.

Results. Oncogenic risk HPVs were detected in all spouses of infected women, with genotypes matching in 18 out of 38 couples (47.37%). The number of types of viruses detected in women was more significant and averaged 2.13, while in men it was 1.68. In women from married couples, HPV type 16 was most often detected (42.11%), HPV type 51 was detected in 28.95% of cases. Among men, HPV genotype 16 was the most common, found in 28.95% of cases, with a frequency of 21.05%, genotypes 31 and 52 were found. During a standard visual examination, no pathology was detected in any of the men. After the use of peniscopy in 13 (31.6%)

patients, certain clinical manifestations were visualized. So, in 8 of them, flat areas of thin acetowhite epithelium with uneven, indistinct edges were detected, in 3 - limited pearly papules in the region of the coronal sulcus and in 2 cases - in the region of the glans penis. With this subclinical form of infection, the most frequently determined HPV genotype was 31, detected in 9 people, while 16 genotype was found only in 4.

Conclusions: 1. Infection with oncogenic HPV types of sexual partners in married couples who continued uninterrupted sexual life for at least 3 years is 100%. 2. The most common HPV genotypes in this situation are HPV 16, 31 and 52. 3. When using subclinical changes in PVI, 31.6% of those examined are found. Stress is a non-specific reaction of the body to short-term or long-term exposure to various damaging factors that threaten the constancy of its internal environment. This reaction is accompanied by physiological changes involving the endocrine system of the hypothalamic-pituitary system, adrenal glands, the reaction of the sympathetic division of the autonomic nervous system. All this affects the function of the skin. In a holistic organism, its general stress state is measured by the physiological reactions of the nervous, endocrine, immune, and vascular systems, which cannot but directly affect the skin as an organ. As a result, rashes of a different nature may appear on the skin, due to a decrease in immunity and a weakening of its barrier function, infectious-allergic acute inflammatory processes, chronic dermatosis, hair loss. What are the most common causes of stress? Stress is characterized by the development of a general adaptation syndrome - these are nonspecific changes in the functioning of the nervous and endocrine systems, aimed at maintaining the internal environment (homeostasis) of the body. It goes through three stages in its development:

- mobilization phase,
- phase of resistance,
- phase of decompensation.

Exposure to stress factors leads to the destruction of protective factors, resulting in dehydration of the skin under stress. In modern dermatology, there are two main types of stress that have a negative impact: Oxidative, or "physical". Psychological, or nervous. A sudden, acute neuropsychic factor leads to a reaction of the endocrine system. The endocrine glands secrete a significant amount of biologically active substances into the blood - stress hormones, primarily adrenaline and norepinephrine (secreted by the adrenal cortex). The adrenal glands also produce the glucocorticoid cortisol, which, in addition to its anti-stress effect, leads to various changes in the skin.

Literature:

1. Davlatovich A. X. et al. PAPULYOZ DERMATOZINI DAVOLASH UCHUN PIMICRALIMUS BILAN MAXALLIY DAVOLASHNI QO'LLASH //IQRO. – 2023. – T. 2. – №. 2. – C. 79-82.
2. Ilyaevna P. U. et al. MODERN TRENDS IN TREATMENT SICK MICROSPORIA //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 98-106.
3. Jo'raqulovich U. A. et al. ADVANTAN FOYDALANISH ALLERGIY VA YALIGLI DERMATOZLARNI TASHQI TERAPIYA //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 56-66.
4. Bahodirovich T. I. et al. DERMATOSCOPIC PAINTING ACUTE AND CHRONIC INFLAMMATORY AND ALLERGIC DERMATOSES //Journal of new century innovations. – 2023. – T. 31. – №. 3. – C. 63-72.
5. Ilyaevna P. U. et al. DERMATOSCOPIC PAINTING SOME COMMON DERMATOSES //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 107-120.
6. Davlatovich A. S. et al. SKINOREN GEL PAPULYOZI BO 'LGAN KAMMALARNI KOMPLEKS TERAPIYADA PUSTULAR FORMASI ROSACEA //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 83-90.

7. Abdusalamovich R. I. et al. ROLE OUTDOOR THERAPIES IN COMPLEX TREATMENT SICK WITH VULGAR AND ACNE ROSE //Journal of new century innovations. – 2023. – T. 31. – №. 3. – C. 73-83.
8. Kholikovich E. S. et al. APPLICATION OF" BETASALIK" OINTMENT IN THE THERAPY OF CHRONIC INFLAMMATORY AND ALLERGIC DERMATOSES //Journal of new century innovations. – 2023. – T. 31. – №. 3. – C. 56-62.
9. Xolikovich E. S. et al. TAJRIBASI FOYDALANISH PANAVIDA IN DAVOLASH KASAL PAPILLOMAVIRUS INFEKTSION ANOGENITAL HUDUDLAR //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 91-97.
10. Jo'raqulovich U. A. et al. TAJRIBASI FOYDALANISH TERBIZILA IN DAVOLASH MIKROSPORADA BOLALAR VA KATTALAR //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 67-72.
11. Abdusalamovich R. I. et al. AZELAY KISLOTA PAYLASHMALARI-PAPULOZ-PUSTULOZ VA ROSE AKNELARINI DAVOLASH //World scientific research journal. – 2023. – T. 16. – №. 2. – C. 73-79.
12. Jorakulovich U. A. et al. External Therapy of Allergic and Inflammatory Dermatoses Using Advantan. – 2023.
13. Davlatovich A. X. et al. Using Dermatoscopy to Identify and Differentiate Between Acute and Chronic Inflammatory and Allergic Skin Conditions. – 2023.
14. Abdusalamovich R. I. et al. The Significance of Incorporating Outdoor Therapies into a Comprehensive Treatment Plan for Individuals with Common Acne Vulgaris and Rosacea. – 2023.
15. Жалилов А. Х., Ачилова Ф. А. ЖЕЛЕЗОДЕФИЦИТНАЯ АНЕМИЯ У ДЕТЕЙ И ЕЁ ЛЕЧЕНИЕ //E Conference Zone. – 2022. – С. 123-127.
16. Ачилова Ф., Ибатова Ш., Абдукадирова А. EXOKARDIYOGRAFIYA BO'YICHA BOLALARDA KICHIK YURAK ANOMALIYALARINING

- TARQALISHI //Международный журнал научной педиатрии. – 2022. – №. 5. – С. 11-15.
17. Жалилов А., Ачилова Ф. Эффективность комплексной терапии железодефицитной анемии у детей //Журнал проблемы биологии и медицины. – 2016. – №. 3 (89). – С. 25-28.
18. Ачилова Ф. А., Жалилов А. Х. ДАННЫЕ ЭХОКАРДИОГРАФИИ ПРИ МАЛЫХ АНОМАЛИЙ СЕРДЦА У ДЕТЕЙ //E Conference Zone. – 2022. – С. 136-140.
19. Кодирова М., Хайдарова С., Ачилова Ф. КЛИНИКА И ОСНОВНЫЕ ПРИЗНАКИ НЕВРОТИЧЕСКОГО МИОКАРДИТА НА ЭКГ У ДЕТЕЙ РАННЕГО ВОЗРАСТА //Журнал гепато-гастроэнтерологических исследований. – 2021. – Т. 2. – №. 3.1. – С. 30-32.
20. Эрметов А. Т. и др. Случай пункционного метода лечения абсцесса печени при деструктивном аппендиците, осложненном пилефлебитом //Вестник экстренной медицины. – 2013. – №. 2. – С. 72-74.
21. Жалилов А., Ачилова Ф. Эффективность комплексной терапии железодефицитной анемии у детей //Журнал проблемы биологии и медицины. – 2016. – №. 3 (89). – С. 25-28.
22. Ачилова Ф., Жалилов А. Показатели эхокардиографии при малых аномалиях сердца у детей //Журнал проблемы биологии и медицины. – 2017. – №. 1 (93). – С. 33-35.
23. Абдурахманова С. Современные подходы к диагностике и лечению сочетанных форм миомы матки и аденомиоза у женщин репродуктивного возраста //in Library. – 2020. – Т. 20. – №. 1. – С. 1-141.
24. Kh J. A., Sh M. U., Achilova F. A. The state of the erythron system in acute pneumonia in children. – 2022.
25. Нажимов Ш. Р. У. и др. Особенности показателей перекисного окисления липидов при хронических гематурических нефритах у детей //Достижения науки и образования. – 2019. – №. 12 (53). – С. 56-58.

26. Пирманова Ш. С. и др. Характеристика остеопении при тубулоинтерстициальных заболеваниях почек у детей //Достижения науки и образования. – 2019. – №. 12 (53). – С. 53-56.
27. Уралов Ш. и др. ABOUT MODERN METHODS OF TREATMENT OF ACUTE STENOSING LARYNGOTRACHEITIS IN CHILDREN //Международный журнал научной педиатрии. – 2022. – №. 5. – С. 25-31.
28. Ахматова Ю. и др. ХРОНИЧЕСКИЙ ТУБУЛОИНТЕРСТИЦИАЛЬНЫЙ НЕФРИТ (ХТИН) У ДЕТЕЙ НА ФОНЕ УРАТУРИИ: КЛИНИКО-ЛАБОРАТОРНАЯ И ФУНКЦИОНАЛЬНАЯ ХАРАКТЕРИСТИКА //InterConf. – 2020.
29. Ахматов А., Ахматова Ю., Сирожова Н. ПИЕЛОНЕФРИТ У ДЕТЕЙ: СОВРЕМЕННЫЕ ОСОБЕННОСТИ ДИАГНОСТИКИ И ЛЕЧЕНИЯ //InterConf. – 2020.
30. Рахмонов З. М. и др. Компенсаторно-приспособительная реакция сосудистого русла внутренних органов при резекции желудка //International Scientific and Practical Conference World science. – ROST, 2017. – Т. 5. – №. 6. – С. 42-44.
31. Хусанов Э. У., Коржавов Ш. О., Сулейманов Р. И. Исследование экскреции лактата кожей в зависимости от различных факторов //Re-health journal. – 2019. – №. 2. – С. 72-75.
32. Oblakulovich K. S. et al. The Thymus gland (Thymus) aspects in children (review of literature) //Central Asian Journal of Medical and Natural Science. – 2022. – Т. 3. – №. 2. – С. 9-16.
33. Хусанов Э. У., Коржавов Ш. О., Ортикбаева Н. Т. Морфологическая картина дегрануляции апудоцитов гастродуоденальной зоны при экспериментальном голодании //International Scientific and Practical Conference World science. – ROST, 2017. – Т. 5. – №. 5. – С. 59-61.

34. Коржавов Ш. О. и др. Развитие микроциркуляторного русла аффлекторов кожи в постнатальном онтогенезе //International Scientific and Practical Conference World science. ROST. – 2017. – Т. 5. – №. 5. – С. 41-43.
35. Негматова Г. Ш. и др. Особенности аутоиммунный тиреоидит в условиях йодного дефицитного региона //Scientific progress. – 2022. – Т. 3. – №. 1. – С. 356-359.
36. Каттаходжаева Д., Муминов О. Морфологическое изменения надпочечников на фоне сахарного диабета : дис. – ТГМУ имени абу Али ибн сино, 2022.
37. Курбонова З. Ч., Бабаджанова Ш. А., Муминов О. А. Лабораторный мониторинг патологии коагуляционного гемостаза у больных COVID-19. – 2021.
38. Курбонова З. Ч. и др. Характеристика функции тромбоцитов при COVID-19. – 2021.
39. Бабаджанова Ш. А., Курбонова З. Ч., Муминов О. А. Частота тромбоэмболических осложнений у больных с коронавирусной инфекцией. – 2021.