

Western blot analysis

Marzia Ognibene¹, Davide Cangelosi¹, Martina Morini¹, Daniela Segalerba¹, Maria Carla Bosco¹, Angela Rita Sementa², Alessandra Eva¹, Luigi Varesio¹

¹ Laboratory of Molecular Biology, Giannina Gaslini Institute, Genova, Italy

² Department of Pathology, Giannina Gaslini Institute, Genova, Italy

Cells are washed in cold PBS and lysed on ice in a buffer containing 1,6mM NaH₂PO₄, 8,6mM Na₂HPO₄, 1% Triton X-100, 0,1% SDS, 0,1% NaN₃, 0,1M NaCl, 0,5% NaDoc, 2mM AEBSF, and 20mg/ml each of aprotinin and leupeptin.

Cell lysates are centrifuged in a microfuge at maximum speed for 10 min to eliminate debris.

Protein content is evaluated by bicinchoninic acid assay (BCA Protein assay kit by Thermo Fisher) in a spectrophotometer.

Lysates (100 µg each) are subjected to 8% SDS-PAGE electrophoresis and transferred to PVDF membrane.

PVDF membrane is incubated in blocking buffer (3% nonfat milk in T-TBS -Tris Buffer Solution 1x with 0,05%Tween) for 2 hr at room temperature.

Membrane is incubated in primary antibody diluted in 0,5%BSA in T-TBS, overnight at 4°C.

Membrane is washed in T-TBS, 5 times, 5 min each.

Membrane is incubated in HRP-conjugated secondary antibody diluted in 0,5%BSA in T-TBS, 2 hr, at room temperature.

Membrane is washed in T-TBS, 5 times, 5 min each.

Proteins are visualized by enhanced chemiluminescence (ECL) detection, through X-ray films exposition or using the Bio-Rad ChemiDoc instrument.