

Partial Differential Equations

HHH

November 15, 2025

Contents

I	Four Important Linear PDE	1
I	Transport Equation	2
II	Laplace Equation	3
§1	Fundmental Solution	3
§2	Solving Poisson's equation	4
§3	Harmonic Function	5
§3.1	Mean-value and Smooth properties	5
§3.2	Maximum Principle	7
§3.3	Local Estimates on Derivatives and Analyticity	8
§4	Subharmonic function	10
§5	Green's funcion	11
§5.1	(Upper) Half-Space	13
§5.2	Ball	14
§6	Energy Methods	15
III	Heat Equation	17
§1	Fundmental Solution	17
§2	Mean-value formula	19
§3	Properties of solution	21
§4	Cauchy Problem	23
§5	Energy Methods	23
IV	Wave Equation	26
§1	Solution for $n=1$, d'Alembert's formula	26
§2	27
§3	Solution for even $n=2k+1$	28
§4	Nonhomogenous Problem	30
§5	Energy Methods	31

II	33
V Nonlinear First-Order PDE	34
§1 Complete integrals, Envelopes	34
§1.1 Complete integrals	34
§2 Characteristics	36
§2.1 Derivation of Characteristic ODE	36
§2.2 Boundary Condition	38
§2.3 Local Solution	40
§3 Hamilton-Jacobi Equation	41
§3.1	41
§3.2 Legendre Transform, Hopf-Lax Formula	44
III Sobolev Space	45
VI Sobolev Space	46
§1 Hölder spaces $C^{k,\gamma}$	46
§1.1 Preliminaries and notation	46
§1.2 The space C^m	47
§1.3 Hölder spaces $C^{k,\gamma}(\bar{\Omega})$	47
§2 Positive Sobolev Spaces $W^{m,p}$	48
§3 Negative Sobolev Spaces $W^{-m,p'}(\Omega)$	49
§4 Extensions	50
§5 Density Theorem	51
§5.1 Approximation by Functions in $C^\infty(\Omega)$	51
§5.2 Approximation by functions in $C^\infty(\bar{\Omega})$	52
§6 Trace	54
§7 The space $W^{k,2}$	55
§7.1 $W^{k,2}(\mathbb{R}^n)$	55
§7.2 Gelfand triple	57
VII Sobolev Inequality; Imbedding Theorem	58
§1 Sobolev inequalities	58
§1.1 Proof of Sobolev inequalities	58
§2 Morrey's inequality	62
§2.1 Proof of Morrey's inequality	62
§3 Poincaré's inequality	65
§4 Compactness	66
§5 Difference quotients and $W^{1,p}$	69
§5.1 Lipschitz functions and $W^{1,\infty}$	71

IV Second Order Elliptic Equatons 73

VII Second Order Elliptic Equations 73

§1	Elliptic Operator and Weak Solution	73
§2	Existence of Weak Solutions	75
§2.1	First Existence Theorem	75
§2.2	Second Existence Theorem	78
§2.3	Third Existence Theorem	80
§3	Regularity	81
§3.1	Interior regularity	81
§3.2	Boundary regularity	87
§4	Maximum principle	88

IX Linear Evolution Equations 93

§1	Second-order parabolic equations	93
§1.1	Weak solution	93
§1.2	Existence of weak solution	95
	Galerkin approximations	95
	Energy estimates	96
	Existence and uniqueness	96
§1.3	Regularity	96

Part I

Four Important Linear PDE

Part II

Part III

Sobolev Space

Chapter VI

Sobolev Space

Contents

§1	Hölder spaces $C^{k,\gamma}$	46
§1.1	Preliminaries and notation	46
§1.2	The space C^m	47
§1.3	Hölder spaces $C^{k,\gamma}(\bar{\Omega})$	47
§2	Positive Sobolev Spaces $W^{m,p}$	48
§3	Negative Sobolev Spaces $W^{-m,p'}(\Omega)$	49
§4	Extensions	50
§5	Density Theorem	51
§5.1	Approximation by Functions in $C^\infty(\Omega)$	51
§5.2	Approximation by functions in $C^\infty(\bar{\Omega})$	52
§6	Trace	54
§7	The space $W^{k,2}$	55
§7.1	$W^{k,2}(\mathbb{R}^n)$	55
§7.2	Gelfand triple	57

§1 Hölder spaces $C^{k,\gamma}$

§1.1 Preliminaries and notation

Throughout the text X denotes a subset of \mathbb{R}^n and $\Omega \subset \mathbb{R}^n$ an open set. We use $C(X)$ for the space of real-valued continuous functions on X (when X is equipped with the subspace topology of \mathbb{R}^n). For a multi-index $\alpha \in \mathbb{N}^n$ we write $|\alpha| = \alpha_1 + \cdots + \alpha_n$ and $\partial^\alpha = \partial_{x_1}^{\alpha_1} \cdots \partial_{x_n}^{\alpha_n}$.

§1.2 The space C^m

Lemma 1.1. *Let X be subset of \mathbb{R}^n .*

1. *Then*

$$C_b(X) = \left\{ f \in C(X) : \sup_{x \in X} |f(x)| < \infty \right\}$$

is a Banach space with the norm $\|f\|_{C_b(X)} := \sup_{x \in X} |f(x)|$.

Remark. *Noted that if X is compact, then $C_b(X) = C(X)$.*

2. *One can view $C(\overline{X})$ as the set*

$$\left\{ f \in C(X) : \exists \text{ continuous } \tilde{f} : \overline{X} \rightarrow \mathbb{R} \text{ s.t. } \tilde{f}|_X = f \right\}$$

Remark. *The extension \tilde{f} is unique if it exists thus we identify f with \tilde{f} if f can be extended to \overline{X} .*

Definition 1.2. *Let Ω be a open set in \mathbb{R}^n , $m \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$ and $0 < \gamma \leq 1$*

1. *The space*

$$C^m(\Omega) := \{u : \Omega \rightarrow \mathbb{R} \mid \partial^\alpha u \text{ exists and is continuous for all } |\alpha| \leq m\}$$

2. *The space*

$$C_b^m(\Omega) := \{u \in C^m(\Omega) \mid \partial^\alpha u \text{ bounded for all } |\alpha| \leq m\}$$

is a Banach space with the norm $\|u\|_{C_b^m(\Omega)} := \sum_{|\alpha| \leq m} \sup_{x \in \Omega} |\partial^\alpha u(x)|$

3. *The space*

$$C^m(\bar{\Omega}) := \{u \in C^m(\Omega) \mid \partial^\alpha u \text{ admits a continuous extension to } \bar{\Omega} \text{ for all } |\alpha| \leq m\}$$

4. *The space*

$$C_b^m(\bar{\Omega}) := C^m(\bar{\Omega}) \cap C_b^m(\Omega)$$

Remark. *Sometimes, we write $(C^m(\Omega), \|\cdot\|_{C^m(\Omega)})$ to denote $(C_b^m(\Omega), \|\cdot\|_{C_b^m(\Omega)})$, and $(C^m(\bar{\Omega}), \|\cdot\|_{C^m(\bar{\Omega})})$ to denote $(C_b^m(\bar{\Omega}), \|\cdot\|_{C_b^m(\bar{\Omega})})$ for simplicity.*

§1.3 Hölder spaces $C^{k,\gamma}(\bar{\Omega})$

Definition 1.3. *Let Ω be a open set in \mathbb{R}^n , $m \in \mathbb{Z}_{\geq 0}$ and $0 < \gamma \leq 1$. If $u : \Omega \rightarrow \mathbb{R}$, we define*

1. The γ^{th} -Hölder seminorm of u is

$$[u]_{C^{0,\gamma}(\Omega)} := \sup_{\substack{x,y \in \Omega \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\gamma}$$

2. The Hölder space

$$C^{k,\gamma}(\Omega)$$

consists of all functions $u \in C^k(\bar{\Omega})$ for which the norm

$$\|u\|_{C^{k,\gamma}(\Omega)} := \sum_{|\alpha| \leq k} \|D^\alpha u\|_{C(\Omega)} + \sum_{|\alpha|=k} [D^\alpha u]_{C^{0,\gamma}(\Omega)}$$

is finite.

Remark. Noted that $C^{k,\gamma}(\Omega) \subset C^k(\bar{\Omega})$ since u^β is uniformly continuous for all $|\beta| = k$. And if $u \in C^{k,\gamma}(\Omega)$ is extended to $\bar{\Omega}$, then the extension is also in $C^{k,\gamma}(\bar{\Omega})$. Thus

$$C^{k,\gamma}(\Omega) = C^{k,\gamma}(\bar{\Omega}) := \left\{ u \in C^k(\bar{\Omega}) : \|D^\alpha u\|_{C(\bar{\Omega})}, [u]_{C^{0,\gamma}(\bar{\Omega})} < \infty \right\}$$

and the norms are equal. We use $C^{k,\gamma}(\Omega)$ and $C^{k,\gamma}(\bar{\Omega})$ interchangeably, but always mean the latter.

Theorem 1.4. The space $C^{k,\gamma}(\bar{\Omega})$ is a Banach space

§2 Positive Sobolev Spaces $W^{m,p}$

Definition 2.1. Let Ω be an open set in \mathbb{R}^n , $1 \leq p \leq \infty$ and $k \in \mathbb{Z}_{\geq 0}$.

1. The Sobolev space

$$W^{m,p}(\Omega) = \{f \in \mathcal{D}'(\Omega) : D^\alpha f \in L^p(\Omega), 0 \leq |\alpha| \leq k\}$$

with the norm

$$\|u\|_{W^{m,p}(\Omega)} := \begin{cases} \left(\sum_{|\alpha| \leq k} \int_{\Omega} |D^\alpha u|^p dx \right)^{1/p} & (1 \leq p < \infty) \\ \sum_{|\alpha| \leq k} \text{ess sup}_{\Omega} |D^\alpha u| & (p = \infty) \end{cases}$$

2. $H^{m,p}(\Omega) \equiv$ the completion of $\{u \in C^m(\Omega) : \|u\|_{m,p} < \infty\}$ with respect to the norm $\|\cdot\|_{m,p}$

3. $W_0^{m,p}(\Omega) \equiv$ the closure of $C_0^\infty(\Omega)$ in the space $W^{m,p}(\Omega)$.

Theorem 2.2. Let $k \in \mathbb{Z}_{\geq 0}$ and $p \in [1, \infty]$, then

1. $W^{m,p}(\Omega)$ is a Banach space.
2. $W^{m,p}(\Omega)$ is separable if $1 \leq p < \infty$.
3. $W^{m,p}(\Omega)$ is uniformly convex and reflexive if $1 < p < \infty$.
4. In particular, $W^{m,2}(\Omega)$ is a separable Hilbert space with inner product

$$(u, v)_{W^{m,2}(\Omega)} = \sum_{0 \leq |\alpha| \leq m} (D^\alpha u, D^\alpha v)_{L^2} \quad (1)$$

Proof. Step 1. Let $\{f_n\}$ be a Cauchy sequence in $W^{m,p}(\Omega)$, then $\{D^\alpha f_n\}$ are Cauchy sequence in $L^p(\Omega)$ for all index $|\alpha| \leq k$. By the completeness of L^p , there exist f_α such that

$$D^\alpha f_n \rightarrow f_\alpha \quad \text{in } L^p(\Omega)$$

Considering in $\mathcal{D}(\Omega)'$, $f_n \rightarrow f$ in $\mathcal{D}(\Omega)'$, thus $D^\alpha f_n \rightarrow D^\alpha f$ in $\mathcal{D}(\Omega)'$. Then we have $D^\alpha f = f_\alpha$ and

$$f_n \rightarrow f \quad \text{in } W^{m,p}(\Omega)$$

Step 2. Suppose that $p < \infty$. Let $N = \sum_{|\alpha| \leq k} 1$ be the number of all distinct index $|\alpha| \leq k$, define the map

$$u \mapsto \{D^\alpha u\}_{|\alpha| \leq m}$$

from $W^{m,p}(\Omega)$ to $(L^p(\Omega))^N$. Thus $W^{m,p}(\Omega)$ can be viewed as a closed subspace of $(L^p(\Omega))^N$ which is reflexive, then so $W^{m,p}(\Omega)$. \square

§3 Negative Sobolev Spaces $W^{-m,p'}(\Omega)$

Definition 3.1. Let Ω be an open region in \mathbb{R}^n , $1 \leq p < \infty$ and $m \in \mathbb{Z}_{\geq 0}$. We define the dual space of $W_0^{m,p}(\Omega)$,

$$W^{-m,p'}(\Omega) := (W_0^{m,p}(\Omega))^*$$

where $\frac{1}{p'} + \frac{1}{p} = 1$. Noted that $p' \in (1, \infty]$

Theorem 3.2. Let Ω an open set, $k \in \mathbb{Z}_{\geq 0}$, $p' \in (1, \infty]$, we have

$$W^{-m,p'}(\Omega) = \left\{ f = \sum_{|\alpha| \leq k} (-1)^{|\alpha|} D^\alpha f_\alpha : f_\alpha \in L^{p'}(\Omega) \right\}$$

where the functional f acts on $\varphi \in \mathcal{D}(\Omega)$ by

$$\langle f, \varphi \rangle = \sum_{|\alpha| \leq m} \int_{\Omega} f_\alpha D^\alpha \varphi \, dx$$

and extend to $W_0^{m,p'}(\Omega)$ by density. Moreover, the norm of f in $W^{-m,p'}(\Omega)$ is equivalent to

$$\inf \left\{ \left(\sum_{|\alpha| \leq m} \|g_\alpha\|_{L^{p'}(\Omega)}^{p'} \right)^{1/p'} : f = g \text{ in } W^{-m,p'}(\Omega) \right\}$$

§4 Extensions

Theorem 4.1 (Extension Theorem). *Let Ω be a bounded Lipschitz open set; $1 \leq p \leq \infty$. Select a bounded open set V such that $\Omega \subset\subset V$. Then there exists a bounded linear operator*

$$E : W^{k,p}(\Omega) \longrightarrow W^{k,p}(\mathbb{R}^n)$$

such that for each $u \in W^{k,p}(\Omega)$:

- (i) $Eu|_\Omega = u$ in $W^{k,p}(\Omega)$,
- (ii) Eu has support within V ,
- (iii) E is bounded: $\|Eu\|_{W^{k,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{k,p}(\Omega)}$ where the constant $C = C(n, p, \Omega, V)$ depends only on p, Ω , and V .

We call Eu an **extension** of u to \mathbb{R}^n .

Proof. Step 1: Local flattening of the boundary. Since Ω is Lipschitz, for each point $x_i^0 \in \partial\Omega$, there exists a radius $r_i > 0$ and a Lipschitz function $\gamma_i : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ such that, upon a suitable rotation of coordinates,

$$\Omega \cap B(x_i^0, r_i) = \{x = (x', x_n) \in B(x_i^0, r_i) : x_n > \gamma_i(x')\}.$$

By compactness of $\partial\Omega$, finitely many such neighborhoods $B(x_i^0, r_i)$ cover $\partial\Omega$.

Step 2: Local extensions. On each neighborhood $B(x_i^0, r_i)$, define a local extension \tilde{u}_i of u to the whole $B(x_i^0, r_i)$ (or a slightly larger set) by reflection across the Lipschitz boundary:

$$\tilde{u}_i(x', x_n) := \begin{cases} u(x', x_n), & x_n > \gamma_i(x'), \\ u(x', 2\gamma_i(x') - x_n), & x_n \leq \gamma_i(x'). \end{cases}$$

Standard estimates show that

$$\|\tilde{u}_i\|_{W^{k,p}(B(x_i^0, r_i))} \leq C \|u\|_{W^{k,p}(\Omega \cap B(x_i^0, r_i))}.$$

Step 3: Partition of unity. Take smooth functions $\{\phi_i\}_{i=0}^N \subset C_c^\infty(\mathbb{R}^n)$ forming a partition of

unity subordinate to $\{B(x_i^0, r_i)\}_{i=1}^N$ and an interior set $B_0 \subset \Omega$, such that

$$\phi_0 + \sum_{i=1}^N \phi_i = 1 \quad \text{on } \overline{\Omega}.$$

Step 4: Global extension. Define

$$Eu := \phi_0 u + \sum_{i=1}^N \phi_i \tilde{u}_i.$$

By construction:

- (a) $(Eu)|_{\Omega} = u$, since on $\Omega \cap B(x_i^0, r_i)$ we have $\tilde{u}_i = u$,
- (b) $\text{supp}(Eu) \subset V$, by choosing $B(x_i^0, r_i) \subset V$ and $\text{supp } \phi_i \subset B(x_i^0, r_i)$,
- (c) $\|Eu\|_{W^{k,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{k,p}(\Omega)}$, by Leibniz formula and finite overlap of supports.

Therefore, E is a bounded linear extension operator satisfying all required properties. \square

§5 Density Theorem

§5.1 Approximation by Functions in $C^\infty(\Omega)$

Lemma 5.1 (Local approximation by smooth functions). *Let Ω be an open set, $1 \leq p < \infty$. Then for every $u \in W^{k,p}(\Omega)$ there exist local smooth function u_n such that*

$$u_m \rightarrow u \quad \text{in } W_{\text{loc}}^{k,p}(\Omega)$$

Proof. Define the

$$u^\varepsilon = \eta_\varepsilon * u \quad \text{in } \Omega_\varepsilon$$

Then we have $u^\varepsilon \in C^\infty(\Omega_\varepsilon)$ and $D^\alpha(\eta_\varepsilon * u) = \eta_\varepsilon * D^\alpha u$ in Ω_ε for all $|\alpha| \leq k$ and $\varepsilon > 0$. By the consequence of mollifier,

$$D^\alpha u^\varepsilon = (D^\alpha u)^\varepsilon \rightarrow D^\alpha u \quad \text{in } L_{\text{loc}}^p(\Omega)$$

if we choose any open set $V \subset\subset \Omega$, then

$$\|u^\varepsilon - u\|_{W^{k,p}(V)}^p = \sum_{|\alpha| \leq k} \|D^\alpha(u^\varepsilon - u)\|_{L^p(V)}^p = \sum_{|\alpha| \leq k} \|(D^\alpha u)^\varepsilon - D^\alpha u\|_{L^p(V)}^p \rightarrow 0$$

as $\varepsilon \rightarrow 0$. This proves that $u_m \rightarrow u$ in $W_{\text{loc}}^{k,p}(\Omega)$. \square

Theorem 5.2 ($H = W$). *Let Ω be an open set and $1 \leq p < \infty$. Then for every $u \in W^{m,p}(\Omega)$ there exist $u_m \in H^{m,p} = C^\infty(\Omega) \cap W^{m,p}(\Omega)$ such that*

$$u_m \rightarrow u \quad \text{in } W^{m,p}(\Omega)$$

Proof. Step 1. Let $\{V_i\}_{i=1}^\infty$ be an exhaustion by open sets of Ω and $\{\zeta_i\}_{i=1}^\infty$ be a smooth partition of unity subordinate to $\{V_i\}_{i=1}^\infty$. Next, choose any function $u \in W^{m,p}(\Omega)$. We have $\text{supp}(\zeta_i u) \subset \text{supp} \zeta_i \subset V_i$ is compact, $\zeta_i u \in W^{m,p}(\Omega)$, and

$$u = \sum_{i=1}^{\infty} u^i$$

where $u^i := \zeta_i u$

Step 2. Fix $\delta > 0$. For every i , choose $\varepsilon_i > 0$ small that $\text{supp } u^i \subset V_i^{\varepsilon_i}$ ($\text{dist}(\text{supp } u^i, \partial V_i) > 0$) and v^i such that

$$\|v^i - u^i\|_{W^{m,p}(\Omega)} = \|v^i - u^i\|_{W^{m,p}(V_i)} \leq \frac{\delta}{2^{i+1}}$$

Write $v := \sum_{i=1}^{\infty} v^i$. This function is well defined and belongs to $C^\infty(\Omega)$ since $\text{supp } u^i \subset \text{supp } \zeta_i$ and $\{\text{supp } u^i\}$ is locally finite. Then for any compact set $\Omega' \subset \Omega$, we have $\Omega' \subset \bigcup_{i=1}^N V_i$ for some N depending on Ω' and

$$\begin{aligned} \|v - u\|_{W^{k,p}(\Omega')} &= \left\| \sum_{i=1}^N v^i - \sum_{i=1}^N u^i \right\|_{W^{k,p}(\Omega')} \\ &\leq \sum_{i=1}^{\infty} \|v^i - u^i\|_{W^{k,p}(\Omega)} \\ &\leq \delta \sum_{i=1}^{\infty} \frac{1}{2^i} \\ &= \delta \end{aligned}$$

Take the supremum over compact sets $\Omega' \subset \Omega$, to conclude $\|v - u\|_{W^{k,p}(\Omega)} \leq \delta$ by Fatou's lemma. \square

§5.2 Approximation by functions in $C^\infty(\bar{\Omega})$

Theorem 5.3. *Let Ω be a bounded Lipschitz open set; $1 \leq p < \infty$. Then for every $u \in W^{k,p}(\Omega)$, then there exist $u_m \in C^\infty(\bar{\Omega})$ such that*

$$u_m \rightarrow u \quad \text{in } W^{k,p}(\Omega)$$

Proof. **Step 1: Extension to the whole space.** Since Ω is Lipschitz, there exists a bounded linear extension operator

$$E : W^{k,p}(\Omega) \rightarrow W^{k,p}(\mathbb{R}^n)$$

such that $(Eu)|_\Omega = u$ and $\|Eu\|_{W^{k,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{k,p}(\Omega)}$. Set $v := Eu$.

Step 2: Mollification in \mathbb{R}^n . By [theorem 5.2](#), there exist $v_\varepsilon \in C^\infty(\mathbb{R}^n)$ such that

$$\|v_\varepsilon - v\|_{W^{k,p}(\mathbb{R}^n)} \longrightarrow 0$$

Step 3: Restriction back to Ω . Set $u_\varepsilon := v_\varepsilon|_\Omega$. Since v_ε is C^∞ on \mathbb{R}^n , we have $u_\varepsilon \in C^\infty(\overline{\Omega})$. Moreover,

$$\|u_\varepsilon - u\|_{W^{k,p}(\Omega)} = \|v_\varepsilon - v\|_{W^{k,p}(\Omega)} \leq \|v_\varepsilon - v\|_{W^{k,p}(\mathbb{R}^n)} \longrightarrow 0.$$

Choosing a sequence $\varepsilon_m \downarrow 0$ and writing $u_m := u_{\varepsilon_m}$ yields the desired approximating sequence. \square

Proof. 1. Fix any point $x^0 \in \partial\Omega$. As $\partial\Omega$ is C^1 , there exist, a radius $r > 0$ and a C^1 function $\gamma : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ such that-upon relabeling the coordinate axes if necessary, we have

$$\Omega \cap B(x^0, r) = \{x \in B(x^0, r) \mid x_n > \gamma(x_1, \dots, x_{n-1})\}.$$

Set $V := \Omega \cap B(x^0, r/2)$.

2. Define the shifted point

$$x^\varepsilon := x + \varepsilon \lambda e_n \quad (x \in V, \varepsilon > 0),$$

and observe that for some fixed, sufficiently large number $\lambda > 0$ the ball $B(x^\varepsilon, \varepsilon) \subset \Omega \cap B(x^0, r)$ for all $x \in V$ and all small $\varepsilon > 0$.

Now define $u_\varepsilon(x) := u(x^\varepsilon) = u(x + \varepsilon \lambda e_n)$ $x \in V$. Next write $v^\varepsilon = \eta_\varepsilon * u_\varepsilon$. Clearly $v^\varepsilon \in C^\infty(\bar{V})$.

3. We now claim

$$v^\varepsilon \rightarrow u \quad \text{in } W^{k,p}(V)$$

To confirm this, take α to be any multiindex with $|\alpha| \leq k$. Then

$$\begin{aligned} \|D^\alpha v^\varepsilon - D^\alpha u\|_{L^p(V)} &= \|\eta_\varepsilon * (D^\alpha u_\varepsilon - D^\alpha u) + \eta_\varepsilon * D^\alpha u - D^\alpha u\|_{L^p(V)} \\ &\leq \|\eta_\varepsilon * (D^\alpha u_\varepsilon - D^\alpha u)\|_{L^p(V)} + \|\eta_\varepsilon * D^\alpha u - D^\alpha u\|_{L^p(V)} \\ &\leq \|\eta_\varepsilon\| \cdot \|(D^\alpha u)_\varepsilon - D^\alpha u\| + \|\eta_\varepsilon * D^\alpha u - D^\alpha u\| \end{aligned}$$

4. Select $\delta > 0$. Since $\partial\Omega$ is compact, we can find finitely many points $x_i^0 \in \partial\Omega$, radius $r_i > 0$, corresponding sets $V_i = \Omega \cap B^0(x_i^0, \frac{r_i}{2})$, and functions $v_i \in C^\infty(\bar{V}_i)$ ($i = 1, \dots, N$) such that $\partial\Omega \subset \bigcup_{i=1}^N B^0(x_i^0, \frac{r_i}{2})$ and

$$\|v_i - u\|_{W^{k,p}(V_i)} \leq \delta$$

Take an open set

$$V_0 = \Omega_\delta \subset\subset V'_0 = \Omega_{\delta'} \subset\subset \Omega$$

such that $\Omega \subset \bigcup_{i=0}^N V_i$ ($0 < \delta' < \delta < \min\{\frac{r_i}{2}\}$) and select, using Theorem 1, a function $v_0 \in C^\infty(V'_0) \subset C^\infty(\bar{V}_0)$ satisfying

$$\|v_0 - u\|_{W^{k,p}(V_0)} \leq \delta$$

5. Now let $\{\zeta_i\}_{i=0}^N$ be a smooth partition of unity on $\bar{\Omega}$, subordinate to the open sets $\{V_0, B^0(x_1^0, \frac{r_1}{2}), \dots, B^0(x_N^0, \frac{r_N}{2})\}$. Define $v := \sum_{i=0}^N \zeta_i v_i$. Then clearly $v \in C^\infty(\bar{\Omega})$. In addition, since $u = \sum_{i=0}^N \zeta_i u$, we see using Leibniz formula that for each $|\alpha| \leq k$

$$\begin{aligned} \|D^\alpha v - D^\alpha u\|_{L^p(\Omega)} &\leq \sum_{i=0}^N \|D^\alpha (\zeta_i v_i) - D^\alpha (\zeta_i u)\|_{L^p(V_i)} \\ &\leq C_{\zeta, \alpha} \sum_{i=0}^N \|v_i - u\|_{W^{k,p}(V_i)} = C(N+1)\delta \end{aligned}$$

□

§6 Trace

Theorem 6.1 (Trace Theorem). *Let Ω be a bounded Lipschitz open set; $1 \leq p \leq \infty$; $1 \leq p < \infty$. Then there exists a bounded linear operator*

$$T : W^{1,p}(\Omega) \rightarrow L^p(\partial\Omega)$$

such that

$$(i) \quad Tu = u|_{\partial\Omega} \text{ if } u \in C(\bar{\Omega}) \cap W^{1,p}(\Omega)$$

$$(ii) \quad T \text{ is bounded: } \|Tu\|_{L^p(\partial\Omega)} \leq C \|u\|_{W^{1,p}(\Omega)} \text{ with the constant } C \text{ depending only on } p \text{ and } \Omega.$$

We call Tu the **trace** of u on $\partial\Omega$.

Remark. By the [theorem 5.3](#), the trace operator is uniquely determined by property (i).

Proof. Step 1: Local flattening of the boundary. Since Ω is Lipschitz, for each point $x_i^0 \in \partial\Omega$, there exists a neighborhood U_i and a Lipschitz function $\gamma_i : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ such that, after a suitable rotation of coordinates,

$$\Omega \cap U_i = \{x = (x', x_n) \in U_i : x_n > \gamma_i(x')\}.$$

By compactness of $\partial\Omega$, finitely many such neighborhoods U_i cover $\partial\Omega$.

Step 2: Reduction to the upper half-space. Consider the local change of coordinates

$$\Phi_i : U_i \rightarrow \tilde{U}_i \subset \mathbb{R}^n, \quad \Phi_i(x', x_n) = (x', x_n - \gamma_i(x')).$$

Then locally $\Phi_i(\Omega \cap U_i) \subset \mathbb{R}_+^n := \{(x', x_n) : x_n > 0\}$.

For $u \in W^{1,p}(\Omega)$, define $v_i = u \circ \Phi_i^{-1}$ in \mathbb{R}_+^n . Classical one-dimensional argument (Hardy inequality) shows

$$\|v_i(\cdot, 0)\|_{L^p(\mathbb{R}^{n-1})} \leq C \|v_i\|_{W^{1,p}(\mathbb{R}_+^n)}.$$

Step 3: Partition of unity. Take a smooth partition of unity $\{\phi_i\}_{i=1}^N$ subordinate to $\{U_i\}$, such that $\sum_i \phi_i = 1$ near $\partial\Omega$.

Define the local traces:

$$T_i u := v_i(\cdot, 0) = (u \circ \Phi_i^{-1})(\cdot, 0) \in L^p(\mathbb{R}^{n-1}).$$

Then the global trace is

$$Tu := \sum_{i=1}^N (\phi_i T_i u) \circ \Phi_i \in L^p(\partial\Omega).$$

Step 4: Verification.

(a) Linearity and boundedness:

$$\|Tu\|_{L^p(\partial\Omega)} \leq C \sum_i \|T_i u\|_{L^p(\mathbb{R}^{n-1})} \leq C \sum_i \|v_i\|_{W^{1,p}(\mathbb{R}_+^n)} \leq C \|u\|_{W^{1,p}(\Omega)}.$$

(b) Consistency with smooth functions: If $u \in C(\bar{\Omega}) \cap W^{1,p}(\Omega)$, then locally $T_i u = u|_{\partial\Omega \cap U_i}$, so globally

$$Tu = u|_{\partial\Omega}.$$

Hence T defines a bounded linear trace operator on $W^{1,p}(\Omega)$. □

Theorem 6.2 (Trace-zero functions in $W^{1,p}$). *Let Ω be a bounded Lipschitz open set; $1 \leq p < \infty$. Suppose furthermore that $u \in W^{1,p}(\Omega)$. Then*

$$u \in W_0^{1,p}(\Omega) \quad \text{iff} \quad Tu = 0 \text{ on } \partial\Omega.$$

§7 The space $W^{k,2}$

§7.1 $W^{k,2}(\mathbb{R}^n)$

Lemma 7.1. *Supposes $u \in L_{\text{loc}}(\mathbb{R}^n)$ has a weak derivative $D^\alpha u \in L^1(\mathbb{R}^n)$ or $L^2(\mathbb{R}^n)$, then*

$$\mathcal{F}D^\alpha u = (iy)^\alpha \hat{u}$$

Theorem 7.2 (Characterization of $W^{k,2}(\mathbb{R}^n)$ by Fourier transform). *Let k be a nonnegative integer.*

A function $u \in L^2(\mathbb{R}^n)$ belongs to $W^{k,2}(\mathbb{R}^n)$ if and only if

$$(1 + |y|^k) \hat{u} \in L^2(\mathbb{R}^n)$$

Remark. In addition, we have the inequalities

$$\|(1 + |y|^k) \hat{u}\|_{L^2(\mathbb{R}^n)} \leq \sqrt{2} \|u\|_{W^{k,2}(\mathbb{R}^n)}$$

for each $u \in W^{k,2}(\mathbb{R}^n)$. Thus the norm $\|\cdot\| := \|(1 + |y|^k) \mathcal{F}\cdot\|_{L^2(\mathbb{R}^n)}$ is equivalent to $\|\cdot\|_{W^{k,2}(\mathbb{R}^n)}$ by the norm equivalent theorem.

Proof. Assume first $u \in W^{k,2}(\mathbb{R}^n)$. Then for each multiindex $|\alpha| \leq k$, we have $D^\alpha u \in L^2(\mathbb{R}^n)$ and

$$\mathcal{F}(D^\alpha u) = (iy)^\alpha \hat{u}$$

belongs to $L^2(\mathbb{R}^n)$. Also,

$$\|D^\alpha u\|_{L^2(\mathbb{R}^n)} = \|(iy)^\alpha \hat{u}\|_{L^2(\mathbb{R}^n)}$$

Thus

$$\begin{aligned} \|(1 + |y|^k) \hat{u}\|_{L^2(\mathbb{R}^n)} &= \left(\int_{\mathbb{R}^n} (1 + 2|y|^k + |y|^{2k}) \hat{u}^2 dy \right)^{\frac{1}{2}} \\ &\leq \left(\int_{\mathbb{R}^n} 2(1 + |y|^{2k}) \hat{u}^2 dy \right)^{\frac{1}{2}} \\ &\leq \sqrt{2} \left(\|\hat{u}\|_{L^2(\mathbb{R})} + \sum_{|\alpha|=k} \|y^\alpha \hat{u}\|_{L^2(\mathbb{R})} \right) \\ &= \sqrt{2} \left(\|u\|_{L^2(\mathbb{R})} + \sum_{|\alpha|=k} \|D^\alpha \hat{u}\|_{L^2(\mathbb{R})} \right) \\ &\leq \sqrt{2} \|u\|_{H^k(\mathbb{R}^n)} \end{aligned}$$

belongs to $L^2(\mathbb{R}^n)$ and

2. Suppose conversely $(1 + |y|^k) \hat{u} \in L^2(\mathbb{R}^n)$ and $|\alpha| \leq k$. Then

$$\|(iy)^\alpha \hat{u}\|_{L^2(\mathbb{R}^n)}^2 \leq \int_{\mathbb{R}^n} |y|^{2|\alpha|} |\hat{u}|^2 dy \leq C_\alpha \|(1 + |y|^k) \hat{u}\|_{L^2(\mathbb{R}^n)}^2 < \infty$$

thus $(iy)^\alpha \hat{u} \in L^2(\mathbb{R}^n)$. Set

$$u_\alpha := \mathcal{F}^{-1}((iy)^\alpha \hat{u})$$

Then for each $\phi \in C_c^\infty(\mathbb{R}^n)$

$$(D^\alpha \phi, u) = (\mathcal{F} D^\alpha \phi, \hat{u}) = ((iy)^\alpha \hat{\phi}, \hat{u}) = (-1)^{|\alpha|} (\hat{\phi}, (iy)^\alpha \hat{u}) = (-1)^{|\alpha|} (\phi, u_\alpha)$$

Thus $u_\alpha = D^\alpha u$ in the weak sense and $D^\alpha u \in L^2(\mathbb{R}^n)$. Hence $u \in H^k(\Omega)$, as required. \square

Definition 7.3. Let $s > 0$ be a noninteger real number. We define the fractional Sobolev space $H^s(\mathbb{R}^n)$ as follows:

$$H^s(\mathbb{R}^n) := \{u \in L^2(\mathbb{R}^n) : (1 + |y|^s) \hat{u} \in L^2(\mathbb{R}^n)\}$$

with the norm

$$\|u\|_{H^s(\mathbb{R}^n)} := \|(1 + |y|^s) \hat{u}\|_{L^2(\mathbb{R}^n)}$$

§7.2 Gelfand triple

Theorem 7.4. Let $\Omega \subset \mathbb{R}^n$ be a open set. Then the following **Gelfand triple** holds:

$$H_0^k(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow H^{-k}(\Omega)$$

where $H^{-k}(\Omega)$ denotes the dual space of $H_0^k(\Omega)$. Moreover, both embeddings are continuous and dense.

Chapter VII

Sobolev Inequality; Imbedding Theorem

Contents

§1	Sobolev inequalities	58
§1.1	Proof of Sobolev inequalities	58
§2	Morrey's inequality	62
§2.1	Proof of Morrey's inequality	62
§3	Poincaré's inequality	65
§4	Compactness	66
§5	Difference quotients and $W^{1,p}$	69
§5.1	Lipschitz functions and $W^{1,\infty}$	71

§1 Sobolev inequalities

Theorem 1.1. *Let Ω be a bounded Lipschitz domain in \mathbb{R}^n . If $u \in W_0^{k,p}(\Omega)$ and $kp < n$, then there exists a constant $C = C(k, p, n, \Omega)$ such that*

$$\|u\|_{L^{p^*}(\Omega)} \leq C \|u\|_{W^{k,p}(\Omega)}$$

where $p^* = \frac{np}{n-kp}$ ($\frac{1}{p^*} = \frac{1}{p} - \frac{k}{n}$) is the Sobolev conjugate of p .

§1.1 Proof of Sobolev inequalities

Definition 1.2. *Let n -multiindex \mathbf{p}_i , θ_i and \mathbf{p} be given such that*

$$\frac{\theta}{\mathbf{p}} = \sum_{i=1}^k \frac{\theta_i}{\mathbf{p}_i}$$

meaning that $\frac{\theta_j}{\mathbf{p}^j} = \sum_{i=1}^k \frac{\theta_i^j}{\mathbf{p}_i^j}$ for each component $j = 1, \dots, n$, and let f_i

$$\left\| \prod_{i=1}^k f_i \right\|_{L^{\mathbf{p}}} \leq \prod_{i=1}^k \|f_i\|_{L^{\mathbf{p}_i}}^{\theta_i}$$

where

$$\|u\|_{L^{\mathbf{p}}}^{\theta} := \left\| \cdots \left\| \|f\|_{L^{\mathbf{p}^1}(dx_1)}^{\theta^1} \right\|_{L^{\mathbf{p}^2}(dx_2)}^{\theta^2} \cdots \right\|_{L^{\mathbf{p}^n}(dx_n)}^{\theta^n}$$

Definition 1.3 (Multi-index Hölder inequality). Let n -multiindices \mathbf{p}_i, θ_i and \mathbf{p} be given such that

$$\frac{\theta}{\mathbf{p}} = \sum_{i=1}^k \frac{\theta_i}{\mathbf{p}_i},$$

meaning that

$$\frac{\theta_j}{\mathbf{p}^j} = \sum_{i=1}^k \frac{\theta_i^j}{\mathbf{p}_i^j}, \quad j = 1, \dots, n.$$

For functions f_i , define the nested $L^{\mathbf{p}}$ -norm with weights θ as

$$\|f\|_{L^{\mathbf{p}}}^{\theta} := \left\| \cdots \left\| \|f\|_{L^{\mathbf{p}^1}(dx_1)}^{\theta^1} \right\|_{L^{\mathbf{p}^2}(dx_2)}^{\theta^2} \cdots \right\|_{L^{\mathbf{p}^n}(dx_n)}^{\theta^n}.$$

Then the multi-index Hölder inequality states that

$$\left\| \prod_{i=1}^k f_i \right\|_{L^{\mathbf{p}}} \leq \prod_{i=1}^k \|f_i\|_{L^{\mathbf{p}_i}}^{\theta_i}.$$

Theorem 1.4 (Estimates for $C_c^1(\mathbb{R}^n)$). Assume $1 \leq p < n$. There exists a constant $C = \frac{p(n-1)}{n-p}$, such that

$$\|u\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du\|_{L^p(\mathbb{R}^n)}$$

for all $u \in C_c^1(\mathbb{R}^n)$.

Proof. 1. First Assume $p = 1$. Since u has compact support, for each $i = 1, \dots, n$ and $x \in \mathbb{R}^n$ we have

$$u(x) = \int_{-\infty}^{x_i} u_{x_i}(x_1, \dots, x_{i-1}, y_i, x_{i+1}, \dots, x_n) \, dy_i$$

and so

$$|u(x)| \leq \int_{-\infty}^{\infty} |Du(x_1, \dots, y_i, \dots, x_n)| \, dy_i$$

Consequently

$$|u(x)|^{\frac{n}{n-1}} \leq \prod_{i=1}^n \left(\int_{-\infty}^{\infty} |Du(x_1, \dots, y_i, \dots, x_n)| \, dy_i \right)^{\frac{1}{n-1}} \quad (1)$$

Integrate with respect to x_1 :

$$\begin{aligned}
 \int_{-\infty}^{\infty} |u|^{\frac{n}{n-1}} dx_1 &\leq \int_{-\infty}^{\infty} \prod_{i=1}^n \left(\int_{-\infty}^{\infty} |Du| dy_i \right)^{\frac{1}{n-1}} dx_1 \\
 &= \left(\int_{-\infty}^{\infty} |Du| dy_1 \right)^{\frac{1}{n-1}} \int_{-\infty}^{\infty} \prod_{i=2}^n \left(\int_{-\infty}^{\infty} |Du| dy_i \right)^{\frac{1}{n-1}} dx_1 \\
 &\leq \left(\int_{-\infty}^{\infty} |Du| dy_1 \right)^{\frac{1}{n-1}} \prod_{i=2}^n \left(\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |Du| dx_1 dy_i \right)^{\frac{1}{n-1}}
 \end{aligned}$$

Then integrate this inequality with respect to x_2 :

We continue by integrating with respect to x_3, \dots, x_n , eventually to find

$$\begin{aligned}
 \int_{\mathbb{R}^n} |u|^{\frac{n}{n-1}} dx &\leq \prod_{i=1}^n \left(\int_{\mathbb{R}^n} |Du| dx_1 \dots dy_i \dots dx_n \right)^{\frac{1}{n-1}} \\
 &= \left(\int_{\mathbb{R}^n} |Du| dx \right)^{\frac{n}{n-1}}
 \end{aligned}$$

This is estimate for $p = 1$.

2. Consider now the case that $1 < p < n$. We apply previous estimate to $v := |u|^\gamma$, where $\gamma = \frac{p(n-1)}{n-p} > 1$. Then

$$\begin{aligned}
 \left(\int_{\mathbb{R}^n} |u|^{p^*} dx \right)^{\frac{n-1}{n}} &= \left(\int_{\mathbb{R}^n} |u|^{\frac{\gamma n}{n-1}} dx \right)^{\frac{n-1}{n}} \\
 &\leq \int_{\mathbb{R}^n} |D|u|^\gamma| dx \\
 &= \gamma \int_{\mathbb{R}^n} |u|^{\gamma-1} |Du| dx \\
 &\leq \gamma \left(\int_{\mathbb{R}^n} |u|^{(\gamma-1)\frac{p}{p-1}} dx \right)^{\frac{p-1}{p}} \|Du\|_p \\
 &= \gamma \left(\int_{\mathbb{R}^n} |u|^{p^*} dx \right)^{\frac{p-1}{p}} \|Du\|_p
 \end{aligned}$$

in which case $\frac{\gamma n}{n-1} = (\gamma - 1)\frac{p}{p-1} = \frac{np}{n-p} = p^*$. So we have

$$\left(\int_{\mathbb{R}^n} |u|^{p^*} dx \right)^{\frac{1}{p^*}} \leq \gamma \left(\int_{\mathbb{R}^n} |Du|^p dx \right)^{\frac{1}{p}}$$

□

Theorem 1.5 (Estimates for $W_0^{1,p}$). Assume Ω is a bounded, open subset of \mathbb{R}^n . Suppose $u \in W_0^{1,p}(\Omega)$ for some $1 \leq p < n$. Then we have the estimate

$$\|u\|_{L^{p^*}(\Omega)} \leq C \|Du\|_{L^p(\Omega)}$$

Furthermore, $\|u\|_{L^q(\Omega)} \leq C' \|Du\|_{L^p(\Omega)}$ for each $q \in [1, p^*]$, the constant C' depending only on p, q, n and Ω .

Proof: Since $u \in W_0^{1,p}(\Omega)$, there exist functions $u_m \in C_c^\infty(\Omega)$ converging to u in $W^{1,p}(\Omega)$. We extend each function u_m to be 0 on $\mathbb{R}^n - \bar{\Omega}$ (that $C_c^\infty(\Omega) \subset C_c^\infty(\mathbb{R}^n)$) and apply the estimate for $C_c^\infty(\mathbb{R}^n)$ to discover

$$\|u_m\|_{L^{p^*}(\Omega)} = \|u_m\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du_m\|_{L^p(\mathbb{R}^n)} = C \|Du_m\|_{L^p(\Omega)}$$

Thus

$$\|u_m - u_n\|_{L^{p^*}(\Omega)} \leq C \|Du_m - Du_n\|_{L^p(\Omega)} \leq C \|u_m - u_n\|_{W^{1,p}(\Omega)} \rightarrow 0$$

Since $L^{p^*}(\Omega)$ is complete, we conclude that $u_m \xrightarrow{L^{p^*}(\Omega)} u$ and

$$\|u\|_{L^{p^*}(\Omega)} \leq C \|Du\|_{L^p(\Omega)}$$

Remark In view of this estimate, on $W_0^{1,p}(\Omega)$ the norm $\|Du\|_{L^p(\Omega)}$ is equivalent to $\|u\|_{W^{1,p}(\Omega)}$, if Ω is a bounded, open subset of \mathbb{R}^n .

$$\|Du\|_{L^p(\Omega)} \leq \|u\|_{W^{1,p}(\Omega)}$$

$$\|u\|_{W^{1,p}(\Omega)}^p = \|u\|_{L^p(\Omega)}^p + \|Du\|_{L^p(\Omega)}^p \leq C \|Du\|_{L^p(\Omega)}^p$$

since $p \in [1, p^*]$.

Theorem 1.6 (Estimates for $W^{1,p}$, $1 \leq p < n$). *Let Ω be a bounded, open subset of \mathbb{R}^n , and suppose $\partial\Omega$ is C^1 . If $1 \leq p < n$, and $u \in W^{1,p}(\Omega)$. Then $u \in L^{p^*}(\Omega)$, with the estimate*

$$\|u\|_{L^{p^*}(\Omega)} \leq C \|u\|_{W^{1,p}(\Omega)}$$

the constant C depending only on p, n , and Ω .

Proof: Since $\partial\Omega$ is C^1 , there exists an extension $Eu = \bar{u} \in W^{1,p}(\mathbb{R}^n)$, such that

$$\begin{cases} \bar{u} = u \text{ in } \Omega, \bar{u} \text{ has compact support within } V \\ \|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\Omega)} \end{cases}$$

Because \bar{u} has compact support within V ($V \subset \Omega_\varepsilon$), we know that there exist functions $u_m \in C_c^\infty(V) \subset C_c^\infty(\mathbb{R}^n)$ such that

$$u_m \rightarrow \bar{u} \quad \text{in } W^{1,p}(\mathbb{R}^n)$$

Now according to inequality for $C_c^\infty(\mathbb{R}^n)$,

$$\|u_m - u_l\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du_m - Du_l\|_{L^p(\mathbb{R}^n)} \leq C \|u_m - u_l\|_{W^{1,p}(\mathbb{R}^n)}$$

and

$$\|u_m\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|Du_m\|_{L^p(\mathbb{R}^n)}$$

for all $l, m \geq 1$. Thus

$$u_m \xrightarrow{L^{p^*}(\mathbb{R}^n)} \bar{u}$$

and

$$\|\bar{u}\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|D\bar{u}\|_{L^p(\mathbb{R}^n)}$$

Then we conclude that

$$\|u\|_{L^{p^*}(\Omega)} \leq \|\bar{u}\|_{L^{p^*}(\mathbb{R}^n)} \leq C \|D\bar{u}\|_{L^p(\mathbb{R}^n)} \leq C \|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\Omega)}$$

§2 Morrey's inequality

Theorem 2.1. *Let Ω be a bounded Lipschitz domain in \mathbb{R}^n . If $u \in W^{k,p}(\Omega)$ and $kp > n$ then there exists a constant $C = C(k, p, n, \Omega)$ such that*

$$\|u\|_{C^{m,\gamma}(\bar{\Omega})} \leq C \|u\|_{W^{k,p}(\Omega)}$$

where $m = k - 1 - \left\lfloor \frac{n}{p} \right\rfloor$ and $\gamma = \frac{n}{p} - \left\lfloor \frac{n}{p} \right\rfloor$.

§2.1 Proof of Morrey's inequality

Lemma 2.2. *For all $u \in C^1$, we claim there exists a constant $C = \frac{1}{n\alpha(n)}$, depending only on n , such that*

$$\int_{B(x,r)} |u(y) - u(x)| \, dy \leq C \int_{B(x,r)} \frac{|Du(y)|}{|y - x|^{n-1}} \, dy$$

$$\begin{aligned} \int_{B(x,r)} |u(y) - u(x)| \, dy &= \frac{1}{\alpha(n)r^n} \int_0^r \int_{\partial B(0,1)} |u(x + s\omega) - u(x)| s^{n-1} \, dS_\omega \, ds \\ &= \frac{1}{\alpha(n)r^n} \int_0^r s^{n-1} \int_{\partial B(0,1)} \left| \int_0^s \frac{d}{dt} u(x + tw) \, dt \right| \, dS_\omega \, ds \\ &\leq \frac{1}{\alpha(n)r^n} \int_0^r s^{n-1} \int_{\partial B(0,1)} \int_0^s |Du(x + tw)| \, dt \, dS_\omega \, ds \\ &= \frac{1}{\alpha(n)r^n} \int_0^r s^{n-1} \int_{B(x,s)} \frac{|Du(y)|}{|y - x|^{n-1}} \, dy \, ds \\ &\leq \frac{1}{\alpha(n)r^n} \int_0^r s^{n-1} \int_{B(x,r)} \frac{|Du(y)|}{|y - x|^{n-1}} \, dy \, ds \\ &= \frac{1}{n\alpha(n)} \int_{B(x,r)} \frac{|Du(y)|}{|y - x|^{n-1}} \, dy \end{aligned}$$

Theorem 2.3 ($C^1(\mathbb{R}^n) \cap W^{1,p}(\mathbb{R}^n)$). *Assume $n < p \leq \infty$. Then there exists a constant C ,*

depending only on p and n , such that

$$\|u\|_{C^{0,\gamma}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\mathbb{R}^n)}$$

for all $u \in C^1(\mathbb{R}^n)$, where $\gamma = 1 - n/p$

Proof: 1. Now fix $x \in \mathbb{R}^n$. We apply lemma as follows:

$$\begin{aligned} |u(x)| &\leq \int_{B(x,r)} |u(x) - u(y)| \, dy + \int_{B(x,r)} |u(y)| \, dy \\ &\leq C \int_{B(x,r)} \frac{|Du(y)|}{|x - y|^{n-1}} \, dy + C \|u\|_{L^p(B(x,r))} \\ &\leq C \|Du\|_{L^p(\mathbb{R}^n)} \left(\int_{B(x,r)} \frac{1}{|x - y|^{(n-1)\frac{p}{p-1}}} \, dy \right)^{\frac{p-1}{p}} + C \|u\|_{L^p(\mathbb{R}^n)} \\ &\leq C \|u\|_{W^{1,p}(\mathbb{R}^n)} \end{aligned}$$

The last estimate holds since $p > n$ implies $(n-1)\frac{p}{p-1} < n$. As $x \in \mathbb{R}^n$ is arbitrary, it follows that

$$\|u\|_{C(\bar{\Omega})} = \sup_{x \in \mathbb{R}^n} |u| \leq C \|u\|_{W^{1,p}(\mathbb{R}^n)}$$

2. Next, choose any two points $x, y \in \mathbb{R}^n$ and write $r := |x - y|$. Let $W := B(x, r) \cap B(y, r)$.

Then

$$|u(x) - u(y)| \leq \int_W |u(x) - u(z)| \, dz + \int_W |u(y) - u(z)| \, dz$$

But lemma allows us to estimate

$$\begin{aligned} \int_W |u(x) - u(z)| \, dz &\leq \int_{B(x,r)} |u(x) - u(z)| \, dz \\ &\leq C \int_{B(x,r)} \frac{|Du(z)|}{|z - x|^{n-1}} \, dz \\ &\leq C \|Du\|_{L^p(\mathbb{R}^n)} \left(\int_{B(x,r)} \frac{dz}{|x - z|^{(n-1)\frac{p}{p-1}}} \right)^{\frac{p-1}{p}} \\ &= C \frac{n\alpha(n)(p-1)}{pn} r^{1-\frac{n}{p}} \|Du\|_{L^p(\mathbb{R}^n)} \end{aligned}$$

Likewise,

$$\int_W |u(y) - u(z)| \, dz \leq C r^{1-\frac{n}{p}} \|Du\|_{L^p(\mathbb{R}^n)}$$

These estimates yield

$$|u(x) - u(y)| \leq C r^{1-\frac{n}{p}} \|Du\|_{L^p(\mathbb{R}^n)} = C |x - y|^\gamma \|Du\|_{L^p(\mathbb{R}^n)}$$

Thus

$$[u]_{C^{0,1-n/p}(\mathbb{R}^n)} = \sup_{x \neq y} \left\{ \frac{|u(x) - u(y)|}{|x - y|^{1-n/p}} \right\} \leq C \|Du\|_{L^p(\mathbb{R}^n)}$$

This complete the proof.

Theorem 2.4 (Estimates for $W^{1,p}$). *Let Ω be a bounded, open subset of \mathbb{R}^n , and suppose $\partial\Omega$ is C^1 . Assume $n < p \leq \infty$ and $u \in W^{1,p}(\Omega)$. Then u has a version $u^* \in C^{0,\gamma}(\bar{\Omega})$, for $\gamma = 1 - \frac{n}{p}$, with the estimate*

$$\|u^*\|_{C^{0,\gamma}(\bar{\Omega})} \leq C \|u\|_{W^{1,p}(\Omega)}$$

The constant C depends only on p, n and Ω . In view of this Theorem, we will henceforth always identify a function $u \in W^{1,p}(\Omega)$ ($p > n$) with its continuous version.

Proof: Since $\partial\Omega$ is C^1 , there exists an extension $Eu = \bar{u} \in W^{1,p}(\mathbb{R}^n)$ such that

$$\begin{cases} \bar{u} = u \text{ in } \Omega \\ \bar{u} \text{ has compact support, and} \\ \|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\Omega)} \end{cases}$$

Assume first $n < p < \infty$. Since \bar{u} has compact support, we obtain functions $u_m \in C_c^\infty(\mathbb{R}^n)$ such that

$$u_m \xrightarrow{W^{1,p}(\mathbb{R}^n)} \bar{u}$$

Now according to previous theorem ,

$$\|u_m - u_l\|_{C^{0,\gamma}(\mathbb{R}^n)} \leq C \|u_m - u_l\|_{W^{1,p}(\mathbb{R}^n)}$$

and

$$\|u_m\|_{C^{0,\gamma}} \leq C \|u_m\|_{W^{1,p}(\mathbb{R}^n)}$$

for all $l, m \geq 1$, whence there exists a function $u^* \in C^{0,\gamma}(\mathbb{R}^n)$ such that

$$u_m \xrightarrow{C^{0,\gamma}(\mathbb{R}^n)} u^*$$

We see that $u^* = u$ a.e. on Ω , so that u^* is a version of u and

$$\|u^*\|_{C^{0,1-n/p}(\mathbb{R}^n)} \leq C \|\bar{u}\|_{W^{1,p}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\Omega)}$$

2. If $p = \infty$, $\gamma = 1$ we can assume that $\bar{u} \leq \|\bar{u}\|_{L^\infty(\mathbb{R}^n)}$ for all $x \in \mathbb{R}^n$. Then

$$\sup_{x \in \mathbb{R}^n} \bar{u} \leq \|\bar{u}\|_{L^\infty(\mathbb{R}^n)}$$

and

$$\begin{aligned} |\bar{u}(y) - \bar{u}(x)| &= \left| \int_0^{|y-x|} \frac{d}{dt} \bar{u}\left(x + t \frac{y-x}{|y-x|}\right) dt \right| \\ &\leq \left| \frac{y-x}{|y-x|} \cdot D\bar{u}\left(x + t \frac{y-x}{|y-x|}\right) \right| |y-x| \\ &\leq \|D\bar{u}\|_{L^\infty(\mathbb{R}^n)} |y-x| \end{aligned}$$

for every $x, y \in \mathbb{R}^n$. Thus $\bar{u} \in C^{0,1}(\mathbb{R}^n)$ and $\bar{u}|_\Omega$ is a version of u . Also,

$$\|u\|_{C^{0,1}(\bar{\Omega})} \leq \|\bar{u}\|_{C^{0,1}(\mathbb{R}^n)} \leq \|\bar{u}\|_{W^{1,\infty}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,\infty}(\Omega)}$$

§3 Poincaré's inequality

Theorem 3.1 (Poincaré's inequality). *Let Ω be a bounded, connected, open subset of \mathbb{R}^n , with a C^1 boundary $\partial\Omega$. Assume $1 \leq p \leq \infty$. Then there exists a constant C , depending only on n, p and Ω , such that*

$$\|u - (u)_\Omega\|_{L^p(\Omega)} \leq C \|Du\|_{L^p(\Omega)}$$

for each function $u \in W^{1,p}(\Omega)$.

Proof: We argue by contradiction. Were the stated estimate false, there would exist for each integer $k = 1, \dots$ a function $u_k \in W^{1,p}(\Omega)$ satisfying

$$\|u_k - (u_k)_\Omega\|_{L^p(\Omega)} > k \|Du_k\|_{L^p(\Omega)}$$

We renormalize by defining

$$v_k := \frac{u_k - (u_k)_\Omega}{\|u_k - (u_k)_\Omega\|_{L^p(\Omega)}} \quad (k = 1, \dots)$$

Then

$$(v_k)_\Omega = 0, \|v_k\|_{L^p(\Omega)} = 1$$

and (2) implies

$$\|Dv_k\|_{L^p(\Omega)} < \frac{1}{k} \quad (k = 1, 2, \dots)$$

In particular the functions $\{v_k\}_{k=1}^\infty$ are bounded in $W^{1,p}(\Omega)$. Then there exist a subsequence $\{v_{k_j}\}_{j=1}^\infty \subset \{v_k\}_{k=1}^\infty$ and a function $v \in L^p(\Omega)$ such that

$$v_{k_j} \rightarrow v \quad \text{in } L^p(\Omega)$$

It follows that

$$(v)_\Omega = 0, \|v\|_{L^p(\Omega)} = 1$$

On the other hand, since $\|Dv_k\|_{L^p(\Omega)} \rightarrow 0$

$$\int_{\Omega} v \phi_{x_i} dx = \lim_{k_j \rightarrow \infty} \int_{\Omega} v_{k_j} \phi_{x_i} dx = - \lim_{k_j \rightarrow \infty} \int_{\Omega} v_{k_j, x_i} \phi dx = 0$$

for each $\phi \in C_c^\infty(\Omega)$. Consequently $v \in W^{1,p}(\Omega)$, with $Dv = 0$ a.e.

Thus v is constant, since Ω is connected. Since v is constant and $(v)_\Omega = 0$, we must have $v \equiv 0$, in which case $\|v\|_{L^p(\Omega)} = 0$. This contradiction establishes estimate.

§4 Compactness

Definition 4.1. Let X and Y be Banach spaces, $X \subset Y$. We say that X is **compactly embedded** in Y , written

$$X \subset\subset Y$$

provided the imbedding operator is a linear compact operator i.e.

(i) $\|u\|_Y \leq C \|u\|_X$ for all $u \in X$, where C is a constant,

(ii) each bounded sequence in X is precompact in Y .

Theorem 4.2 (Rellich-Kondrachov Compactness Theorem). Assume Ω is a bounded open subset of \mathbb{R}^n and $\partial\Omega$ is C^1 . Suppose $1 \leq p < n$. Then

$$W^{1,p}(\Omega) \subset\subset L^q(\Omega)$$

for each $1 \leq q < p^*$.

Proof: Step 1. Fix $1 \leq q < p^*$ and note that since Ω is bounded and $\partial\Omega$ is C^1 , Gagliardo-Nirenberg-Sobolev inequality for $W^{1,p}(\Omega)$ implies

$$W^{1,p}(\Omega) \subset L^q(\Omega), \quad \|u\|_{L^q(\Omega)} \leq C \|u\|_{W^{1,p}(\Omega)}.$$

It remains therefore to show that if $\{u_m\}_{m=1}^\infty$ is a bounded sequence in $W^{1,p}(\Omega)$, there exists a subsequence $\{u_{m_j}\}_{j=1}^\infty$ which converges in $L^q(\Omega)$.

Step 2. In view of the Extension Theorem, we may with no loss of generality assume that $\Omega = \mathbb{R}^n$ and the functions $\{u_m\}_{m=1}^\infty$ all have compact support in some bounded open set $V \subset \mathbb{R}^n$. We also may assume

$$\sup_m \|u_m\|_{W^{1,p}(V)} < \infty.$$

Step 3. Let us first study the smoothed functions

$$u_m^\varepsilon := \eta_\varepsilon * u_m \quad (\varepsilon > 0, m = 1, 2, \dots),$$

where η_ε denotes the usual mollifier. We may suppose the functions $\{u_m^\varepsilon\}_{m=1}^\infty$ all have support in V as well.

Step 4. We first claim

$$u_m^\varepsilon \xrightarrow{L^q(V)} u_m \text{ as } \varepsilon \rightarrow 0, \text{ uniformly in } m.$$

To prove this, we first note that if u_m is smooth, then

$$\begin{aligned} u_m^\varepsilon(x) - u_m(x) &= \frac{1}{\varepsilon^n} \int_{B(x,\varepsilon)} \eta\left(\frac{x-z}{\varepsilon}\right) (u_m(z) - u_m(x)) dz \\ &= \int_{B(0,1)} \eta(y) (u_m(x - \varepsilon y) - u_m(x)) dy \\ &= \int_{B(0,1)} \eta(y) \int_0^1 \frac{d}{dt} (u_m(x - \varepsilon ty)) dt dy \\ &= -\varepsilon \int_{B(0,1)} \eta(y) \int_0^1 Du_m(x - \varepsilon ty) \cdot y dt dy \end{aligned}$$

Thus

$$\begin{aligned} \int_V |u_m^\varepsilon(x) - u_m(x)| dx &\leq \varepsilon \int_{B(0,1)} \eta(y) \int_0^1 \int_V |Du_m(x - \varepsilon ty)| dx dt dy \\ &\leq \varepsilon \int_V |Du_m(z)| dz. \end{aligned}$$

By approximation, this estimate holds if $u_m \in W^{1,p}(V)$. Hence

$$\|u_m^\varepsilon - u_m\|_{L^1(V)} \leq \varepsilon \|Du_m\|_{L^1(V)} \leq \varepsilon C \|Du_m\|_{L^p(V)},$$

We thereby discover

$$u_m^\varepsilon \rightarrow u_m \text{ in } L^1(V), \text{ uniformly in } m.$$

But then since $1 \leq q < p^$, we see using the interpolation inequality for L^p -norms that*

$$\|u_m^\varepsilon - u_m\|_{L^q(V)} \leq \|u_m^\varepsilon - u_m\|_{L^1(V)}^\theta \|u_m^\varepsilon - u_m\|_{L^{p^*}(V)}^{1-\theta},$$

where $\frac{1}{q} = \frac{\theta}{1} + \frac{1-\theta}{p^}$, $0 < \theta < 1$. Consequently, the Gagliardo-Nirenberg-Sobolev inequality imply*

$$\|u_m^\varepsilon - u_m\|_{L^q(V)} \leq C \|u_m^\varepsilon - u_m\|_{L^1(V)}^\theta,$$

whence assertion (2) follows from (3).

Step 5. Next we claim

$$\left\{ \begin{array}{l} \text{for each fixed } \varepsilon > 0, \text{ the sequence } \{u_m^\varepsilon\}_{m=1}^\infty \\ \text{is uniformly bounded and equicontinuous.} \end{array} \right.$$

Indeed, if $x \in \mathbb{R}^n$, then

$$|u_m^\varepsilon(x)| \leq \int_{B(x,\varepsilon)} \eta_\varepsilon(x-y) |u_m(y)| dy$$

$$\leq \|\eta_\varepsilon\|_{L^\infty(\mathbb{R}^n)} \|u_m\|_{L^1(V)} \leq \frac{C}{\varepsilon^n} < \infty$$

for $m = 1, 2, \dots$. Similarly,

$$\begin{aligned} |Du_m^\varepsilon(x)| &\leq \int_{B(x,\varepsilon)} |D\eta_\varepsilon(x-y)| |u_m(y)| dy \\ &\leq \|D\eta_\varepsilon\|_{L^\infty(\mathbb{R}^n)} \|u_m\|_{L^1(V)} \leq \frac{C}{\varepsilon^{n+1}} < \infty, \end{aligned}$$

for $m = 1, \dots$. Assertion (4) follows from these two estimates.

Step 6. We now observe that since the functions $\{u_m\}_{m=1}^\infty$, and thus the functions $\{u_m^\varepsilon\}_{m=1}^\infty$, have support in some fixed bounded set $V \subset \mathbb{R}^n$, we may utilize (4) and the Arzela-Ascoli compactness criterion, to obtain a subsequence $\{u_{m_j}^\varepsilon\}_{j=1}^\infty \subset \{u_m^\varepsilon\}_{m=1}^\infty$ which converges uniformly on V . In particular therefore

$$\limsup_{j,k \rightarrow \infty} \|u_{m_j}^\varepsilon - u_{m_k}^\varepsilon\|_{L^q(V)} = 0$$

Now fix $\delta > 0$. We will show there exists a subsequence $\{u_{m_j}\}_{j=1}^\infty \subset \{u_m\}_{m=1}^\infty$ such that

$$\limsup_{j,k \rightarrow \infty} \|u_{m_j} - u_{m_k}\|_{L^q(V)} \leq \delta.$$

To see this, let us first employ assertion (2) to select $\varepsilon > 0$ so small that

$$\|u_m^\varepsilon - u_m\|_{L^q(V)} \leq \delta/2$$

for $m = 1, 2, \dots$

But then (6) and (7) imply

$$\limsup_{j,k \rightarrow \infty} \|u_{m_j} - u_{m_k}\|_{L^q(V)} \leq \delta$$

and so (5) is proved.

Step 7. We next employ assertion (5) with $\delta = 1, \frac{1}{2}, \frac{1}{3}, \dots$ and use a standard diagonal argument to extract a subsequence $\{u_{m_l}\}_{l=1}^\infty \subset \{u_m\}_{m=1}^\infty$ satisfying

$$\limsup_{l,k \rightarrow \infty} \|u_{m_l} - u_{m_k}\|_{L^q(V)} = 0.$$

Corollary 4.3. Assume Ω is a bounded open subset of \mathbb{R}^n and $\partial\Omega$ is C^1 . We have in particular

$$W^{1,p}(\Omega) \subset\subset L^p(\Omega)$$

for all $1 \leq p \leq \infty$.

Proof: 1. If $1 \leq p < n$, it is obvious from Rellich-Kondrachov Compactness Theorem.

2. If $n < p \leq \infty$, there is a version $u_n^* \in C^{0,\gamma}(\bar{\Omega})$ of u_n for bounded sequence $\{u_n\}$ in $W^{1,p}(\Omega)$

and also

$$\|u_n^*\|_{C^{0,\gamma}(\bar{\Omega})} \leq C\|u_n\|_{W^{1,p}(\Omega)} \leq M$$

which implies u_n^* is uniformly bounded and equicontinuous. By Arzela-Ascoli theorem, $\{u_{n_k}\}$ convergent uniformly to $u \in C(\Omega)$ and also

$$u_{n_k} \rightarrow u \quad \text{in } L^p(\Omega)$$

since Ω is bounded.

Theorem 4.4. Assume Ω is a bounded open subset of \mathbb{R}^n . Suppose $1 \leq p < n$. Then

$$W^{1,p}(\Omega) \subset\subset L^q(\Omega)$$

for each $1 \leq q < p^*$.

§5 Difference quotients and $W^{1,p}$

Definition 5.1. Assume $u : \Omega \rightarrow \mathbb{R}$ is a locally summable function and $V \subset\subset \Omega$.

(1) The i^{th} -difference quotient of size h is

$$D_i^h u(x) = \frac{u(x + he_i) - u(x)}{h} \quad (i = 1, \dots, n)$$

for $x \in V$ and $h \in \mathbb{R}, 0 < |h| < \text{dist}(V, \partial\Omega)$.

(2) $D^h u := (D_1^h u, \dots, D_n^h u)$.

Proposition 5.2. Choose $i = 1, \dots, n, \phi \in C_c^\infty(V)$, and note for small enough h that

$$\int_V u(x) \left[\frac{\phi(x + he_i) - \phi(x)}{h} \right] dx = - \int_V \left[\frac{u(x) - u(x - he_i)}{h} \right] \phi(x) dx$$

that is,

$$\int_V u (D_i^h \phi) dx = - \int_V (D_i^{-h} u) \phi dx$$

Theorem 5.3 (Difference quotients and weak derivatives). Let Ω be a open sets of \mathbb{R}^n .

(1) Suppose $1 \leq p < \infty$ and $u \in W^{1,p}(\Omega)$. Then for each $V \subset\subset \Omega$

$$\|D^h u\|_{L^p(V)} \leq C \|Du\|_{L^p(\Omega)}$$

for some constant C depending on Ω, V and all $0 < |h| < \frac{1}{2} \text{dist}(V, \partial\Omega)$.

(2) Assume $1 < p < \infty, u \in L^p(V)$, and there exists a constant C such that

$$\|D^h u\|_{L^p(V)} \leq C$$

for all $0 < |h| < \frac{1}{2} \text{dist}(V, \partial\Omega)$. Then

$$u \in W^{1,p}(V), \quad \text{with} \quad \|Du\|_{L^p(V)} \leq C$$

Proof: Step 1. Assume $1 \leq p < \infty$, and temporarily suppose u is smooth. Then for each $x \in V, i = 1, \dots, n$, and $0 < |h| < \frac{1}{2} \text{dist}(V, \partial\Omega)$, we have

$$u(x + he_i) - u(x) = h \int_0^1 u_{x_i}(x + the_i) dt$$

so that

$$|u(x + he_i) - u(x)| \leq |h| \int_0^1 |Du(x + the_i)| dt$$

Consequently

$$\begin{aligned} \int_V |D^h u|^p dx &\leq C \sum_{i=1}^n \int_V \int_0^1 |Du(x + the_i)|^p dt dx \\ &= C \sum_{i=1}^n \int_0^1 \int_V |Du(x + the_i)|^p dx dt \end{aligned}$$

Thus

$$\int_V |D^h u|^p dx \leq C \int_\Omega |Du|^p dx$$

This estimate holds should u be smooth, and thus is valid by approximation for arbitrary $u \in W^{1,p}(\Omega)$.

2. Now suppose estimate holds for all $0 < |h| < \frac{1}{2} \text{dist}(V, \partial\Omega)$ and some constant C . Estimate (9) implies

$$\sup_h \|D_i^{-h} u\|_{L^p(V)} < \infty$$

and therefore, since $1 < p < \infty$, there exists a function $v_i \in L^p(V)$ and a subsequence $h_k \rightarrow 0$ such that

$$D_i^{-h_k} u \rightharpoonup v_i \quad \text{weakly in } L^p(V)$$

But then

$$\begin{aligned} \int_V u \phi_{x_i} dx &= \int_\Omega u \phi_{x_i} dx \\ &= \lim_{h_k \rightarrow 0} \int_\Omega u D_i^{h_k} \phi dx \\ &= - \lim_{h_k \rightarrow 0} \int_V D_i^{-h_k} u \phi dx \\ &= - \int_V v_i \phi dx \\ &= - \int_\Omega v_i \phi dx \end{aligned}$$

Thus $v_i = u_{x_i}$ in the weak sense ($i = 1, \dots, n$), and so $Du \in L^p(V)$. As $u \in L^p(V)$, we deduce therefore that $u \in W^{1,p}(V)$.

§5.1 Lipschitz functions and $W^{1,\infty}$

Theorem 5.4 (Characterization of $W^{1,\infty}$). *Let Ω be open and bounded, with $\partial\Omega$ of class C^1 . Then $u : \Omega \rightarrow \mathbb{R}$ is Lipschitz continuous if and only if $u \in W^{1,\infty}(\Omega)$.*

*Proof. Step 1. First assume $\Omega = \mathbb{R}^n$ and u has compact support. Suppose $u \in W^{1,\infty}(\mathbb{R}^n) \subset C^{0,\gamma}$. Then $u^\varepsilon := \eta_\varepsilon * u$ is smooth and satisfies*

$$\begin{cases} u^\varepsilon \rightarrow u \text{ uniformly as } \varepsilon \rightarrow 0 \\ \|Du^\varepsilon\|_{L^\infty(\mathbb{R}^n)} \leq \|Du\|_{L^\infty(\mathbb{R}^n)} \end{cases}$$

Choose any two points $x, y \in \mathbb{R}^n, x \neq y$. We have

$$\begin{aligned} u^\varepsilon(x) - u^\varepsilon(y) &= \int_0^1 \frac{d}{dt} u^\varepsilon(tx + (1-t)y) dt \\ &= \int_0^1 Du^\varepsilon(tx + (1-t)y) dt \cdot (x - y) \end{aligned}$$

and so

$$|u^\varepsilon(x) - u^\varepsilon(y)| \leq \|Du^\varepsilon\|_{L^\infty(\mathbb{R}^n)} |x - y| \leq \|Du\|_{L^\infty(\mathbb{R}^n)} |x - y|.$$

We let $\varepsilon \rightarrow 0$ to discover

$$|u(x) - u(y)| \leq \|Du\|_{L^\infty(\mathbb{R}^n)} |x - y|.$$

Hence u is Lipschitz continuous.

Step 2. On the other hand assume now u is Lipschitz continuous; we must prove that u has essentially bounded weak first derivatives. Since u is Lipschitz, we see

$$\|D_i^{-h} u\|_{L^\infty(\mathbb{R}^n)} \leq \text{Lip}(u)$$

and thus there exists a function $v_i \in L^\infty(\mathbb{R}^n)$ and a subsequence $h_k \rightarrow 0$ such that

$$D_i^{-h_k} u \rightharpoonup v_i \quad \text{weakly in } L_{loc}^2(\mathbb{R}^n)$$

Consequently

$$\begin{aligned} \int_{\mathbb{R}^n} u \phi_{x_i} dx &= \lim_{h_k \rightarrow 0} \int_{\mathbb{R}^n} u D_i^{h_k} \phi dx \\ &= - \lim_{h_k \rightarrow 0} \int_{\mathbb{R}^n} D_i^{-h_k} u \phi dx \\ &= - \int_{\mathbb{R}^n} v_i \phi dx \end{aligned}$$

by (12). The above equality holds for all $\phi \in C_c^\infty(\mathbb{R}^n)$, and so $v_i = u_{x_i}$ in the weak sense ($i = 1, \dots, n$). Consequently $u \in W^{1,\infty}(\mathbb{R}^n)$.

3. In the general case that Ω is bounded, with $\partial\Omega$ of class C^1 , we as usual extend u to $Eu = \bar{u}$ and apply the above argument.

Corollary 5.5. *Let Ω be a open set $u \in W_{loc}^{1,\infty}(\Omega)$ if and only if u is locally Lipschitz continuous in Ω .*

Theorem 5.6 (Differentiability almost everywhere). *Assume $u \in W_{loc}^{1,p}(\Omega)$ for some $n < p \leq \infty$. Then u is differentiable a.e. in Ω , and its gradient equals its weak gradient a.e.*

Proof: Recall that we always identify u with its continuous version.

1. Assume first $n < p < \infty$. From the remark after the proof of Theorem 4 in §5.6.2, we recall Morrey's estimate

$$|v(y) - v(x)| \leq Cr^{1-\frac{n}{p}} \left(\int_{B(x,2r)} |Dv(z)|^p dz \right)^{1/p} \quad (y \in B(x,r))$$

valid for any C^1 function v and thus, by approximation, for any $v \in W^{1,p}$.

2. Choose $u \in W_{loc}^{1,p}(\Omega)$. Now for a.e. $x \in \Omega$, a version of Lebesgue's Differentiation Theorem (§E.4) implies

$$\int_{B(x,r)} |Du(x) - Du(z)|^p dz \rightarrow 0$$

as $r \rightarrow 0$, Du denoting as usual the weak derivative of u . Fix any such point x and set

$$v(y) := u(y) - u(x) - Du(x) \cdot (y - x)$$

in estimate (14), where

$$r = |x - y|$$

We find

$$\begin{aligned} & |u(y) - u(x) - Du(x) \cdot (y - x)| \\ & \leq Cr^{1-n/p} \left(\int_{B(x,2r)} |Du(x) - Du(z)|^p dz \right)^{1/p} \\ & \leq Cr \left(\int_{B(x,2r)} |Du(x) - Du(z)|^p dz \right)^{1/p} \\ & = o(r) \quad \text{by (15)} \\ & = o(|x - y|) \quad \text{by (16).} \end{aligned}$$

Thus u is differentiable at x , and its gradient equals its weak gradient at x . 3. In case $p = \infty$, we note $W_{loc}^{1,\infty}(\Omega) \subset W_{loc}^{1,p}(\Omega)$ for all $1 \leq p < \infty$ and apply the reasoning above.

Theorem 5.7 (Rademacher's Theorem). *Let u be locally Lipschitz continuous in Ω . Then u is differentiable almost everywhere in Ω .*

Part IV

Second Order Elliptic Equatons