

A Primer of Algebraic D-modules

HHH

December 18, 2025

Contents

I	Weyl Algebra	2
§1	Definition	2
§2	Degree	4
§3	Ideal Structure	4
II	Rings of Differential Operators	5
§1	Definition	5
§2	The Weyl Algebra	6
III	Jacobian Conjecture	8
§1	Polynomial Maps	8
§2	Jacobian Conjecture	9
§3	Derivation	9
§4	Automorphisms	11
IV	Modules Over The Weyl Algebra	13
§1	The Polynomial Ring	13
§2	Twisting	14
§3	Holomorphic Functions	15
V	Differential Equations	16
§1	The D-Modules of Equations	16
§2	Direct Limit of Modules	17
§3	Microfunctions	18
VI		20
§1	Graded and filtered	20
§1.1	Graded	20
§1.2	Filtered	21
§1.3	Induced filtrations	21
VII	Dimension	23
§1	Hilbert Functions	23

Throughout this book, K denotes a field of characteristic zero and $K[X]$ the ring of polynomials $K[x_1, \dots, x_n]$ in n commuting indeterminates over K .

Chapter I

Weyl Algebra

§1 Definition

Definition 1.1. The ring $K[X]$ is a vector space of infinite dimension over K . Its algebra of K -linear operators is denoted by $\text{End}_K(K[X])$. Let $\hat{x}_1, \dots, \hat{x}_n$ be the operators of $K[X]$ which are defined on a polynomial $f \in K[X]$ by the formulae $\hat{x}_i(f) = x_i \cdot f$. Similarly, $\partial_1, \dots, \partial_n$ are the operators defined by $\partial_i(f) = \partial f / \partial x_i$. These are linear operators of $K[X]$.

The n -th **Weyl algebra** A_n is the K -subalgebra of $\text{End}_K(K[X])$ generated by the operators $\hat{x}_1, \dots, \hat{x}_n$ and $\partial_1, \dots, \partial_n$. For the sake of consistency, we write $A_0 = K$.

Proposition 1.2.

$$[\partial_i, \hat{x}_j] = \delta_{ij} \cdot 1$$

$$[\partial_i, \partial_j] = [\hat{x}_i, \hat{x}_j] = 0$$

$$[\partial_i, f] = \frac{\partial f}{\partial x_i}$$

Proposition 1.3. We have

1.

$$\partial^\beta x^\alpha = \sum_{\substack{\beta_1 + \beta_2 = \beta \\ \beta_i \in \mathbb{N}^n}} \binom{\beta}{\beta_1} \frac{\partial x^\alpha}{\partial x^{\beta_1}} \cdot \partial^{\beta_2}$$

thus $\partial^\beta x^\alpha = x^\alpha \partial^\beta + D'$ where $\deg D' \leq |\alpha| + |\beta| - 2$

2. The set $\mathbf{B} = \{x^\alpha \partial^\beta : \alpha, \beta \in \mathbb{N}^n\}$ is a basis of A_n as a vector space over K .

Theorem 1.4. Let the free K -algebra $K\{z_1, \dots, z_{2n}\}$ and J be the two-sided ideal generated by $[z_{i+n}, z_i] - 1$ for $i = 1, 2, \dots, n$ and $[z_i, z_j]$ for $j \neq i + n$ and $1 \leq i, j \leq 2n$. We may define a surjective homomorphism $\phi : K\{z_1, \dots, z_{2n}\} \rightarrow A_n$ by $\phi(z_i) = x_i$ and $\phi(z_{i+n}) = \partial_i$, for $i = 1, 2, \dots, n$.

$$\begin{array}{ccc} \{z_1, \dots, z_n\} & \xrightarrow{\quad} & A_n \\ \downarrow \iota & \nearrow \phi & \\ K\{z_1, \dots, z_n\} & & \end{array}$$

It follows that $J \subseteq \ker \phi$. Thus ϕ induces a homomorphism $\bar{\phi} : K \{z_1, \dots, z_{2n}\} / J \longrightarrow A_n$.

Proof. We may use the relations to show that every element of $K \{z_1, \dots, z_{2n}\} / J$ may be written as a linear combination of monomials of the form

$$z_1^{m_1} \dots z_{2n}^{m_{2n}} + J$$

by 1.3 the images of these monomials under $\bar{\phi}$ form a basis of A_n as a vector space over K . In particular, the monomials must be linearly independent in $K \{z_1, \dots, z_{2n}\} / J$. Hence $\bar{\phi}$ is an isomorphism of vector spaces and, a fortiori, an isomorphism of rings. \square

Corollary 1.5. *Let $m < n$ be positive integers. Choose polynomials $f_i \in K[X]$, for $1 \leq i \leq n$, as follows: if $i \leq m$, then f_i is a polynomial in the variables x_{m+1}, \dots, x_n ; otherwise $f_i = 0$. The map $\sigma : A_n \longrightarrow A_n$ defined by the formulae*

$$\begin{aligned} \sigma(x_i) &= x_i + f_i \\ \sigma(\partial_i) &= \partial_i - \sum_1^n \frac{\partial f_k}{\partial x_i} \partial_k \end{aligned}$$

induce an automorphism of A_n .

Proof. Define a homomorphism ϕ of $K \{z_1, \dots, z_{2n}\}$ to A_n by $\phi(z_i) = x_i + f_i$ and

$$\phi(z_{i+n}) = \partial_i - \sum_1^n \frac{\partial f_k}{\partial x_i} \partial_k$$

Choose i, j such that $1 \leq i, j \leq n$. It is clear that $\phi([z_i, z_j]) = 0$.

Let us calculate $\phi([z_{i+n}, z_j])$. Since ϕ is a ring homomorphism, this is the same as $[\phi(z_{i+n}), \phi(z_j)]$, which equals

$$\begin{aligned} \phi([z_{i+n}, z_j]) &= [\phi(z_{i+n}), \phi(z_j)] \\ &= \left[\partial_i - \sum_1^n \frac{\partial f_k}{\partial x_i} \partial_k, x_j + f_j \right] \\ &= \delta_{ij} + \frac{\partial f_j}{\partial x_i} - \frac{\partial f_j}{\partial x_i} - \sum_1^n \frac{\partial f_k}{\partial x_i} \cdot \frac{\partial f_j}{\partial x_k} \\ &= \delta_{ij} \end{aligned}$$

A similar calculation shows that $\phi([z_{i+n}, z_{j+n}]) = 0$. Thus ϕ induces an endomorphism $\sigma = \bar{\phi}$ of A_n .

A similar argument shows that the map τ defined by the formulae

$$\begin{aligned} \tau(x_i) &= x_i - f_i \\ \tau(\partial_i) &= \partial_i + \sum_1^n \frac{\partial f_k}{\partial x_i} \partial_k \end{aligned}$$

is an endomorphism of A_n . It is now easy to check that τ is the inverse of σ . Thus σ is an automorphism of A_n . \square

§2 Degree

Definition 2.1. Let $D \in A_n$. The **degree** of D is the largest length of the multi-indices $(\alpha, \beta) \in \mathbb{N}^n \times \mathbb{N}^n$ for which $x^\alpha \partial^\beta$ appears with non-zero coefficient in the canonical form of D . It is denoted by $\deg(D)$. As with the degree of a polynomial, we use the convention that the zero polynomial has degree $-\infty$.

Lemma 2.2. We have

$$\partial^\beta x^\alpha = \sum_{\substack{\beta_1 + \beta_2 = \beta \\ \beta_i \in \mathbb{N}^n}} \partial^{\beta_1}(x^\alpha) \partial^{\beta_2}$$

thus $\partial^\beta x^\alpha = x^\alpha \partial^\beta + D'$ where $\deg D' \leq |\alpha| + |\beta| - 2$

Theorem 2.3. The degree satisfies the following properties; for $D, D' \in A_n$:

1. $\deg(D + D') \leq \max\{\deg(D), \deg(D')\}$. if $\deg(D) \neq \deg(D')$ then we have equality in the above formula.
2. $\deg(DD') = \deg(D) + \deg(D')$.
3. $\deg[D, D'] \leq \deg(D) + \deg(D') - 2$.

Corollary 2.4. The algebra A_n is a domain.

§3 Ideal Structure

Theorem 3.1. The algebra A_n is simple.

Proof. Let I be a non-zero two-sided ideal of A_n . Choose an element $D \neq 0$ of smallest degree in I . If D has degree 0, it is a constant, and $I = A_n$. Assume that D has degree $k > 0$ and let us aim at a contradiction.

Suppose that (α, β) is a multi-index of length k . If $x^\alpha \partial^\beta$ is a summand of D with non-zero coefficient and $\beta_i \neq 0$, then $[x_i, D]$ is non-zero and has degree $k - 1$. Since I is a two-sided ideal of A_n , we have that $[x_i, D] \in I$. But this contradicts the minimality of D . Thus $\beta = (0, \dots, 0)$. Since $k > 0$, we must have that $\alpha_i \neq 0$, for some $i = 1, 2, \dots, n$. Hence $[\partial_i, D]$ is a non-zero element of I of degree $k - 1$, and once again we have a contradiction. \square

Corollary 3.2. Every (ring, algebra) endomorphism of A_n is injective.

Theorem 3.3. Every left ideal of A_n is generated by two elements.

Chapter II

Rings of Differential Operators

Let R be a commutative K -algebra.

§1 Definition

Definition 1.1. Let R be a commutative K -algebra.

- (i) We will identify an element $a \in R$ with the operator of $\text{End}_K(R)$ defined by the rule $r \mapsto ar$, for every $r \in R$. An operator $P \in \text{End}_K(R)$ has order zero if $[a, P] = 0$, for every $a \in R$.
- (ii) Suppose we have defined operators of order $< n$. An operator $P \in \text{End}_K(R)$ has order n if it does not have order less than n and $[a, P]$ has order less than n for every $a \in R$.

Let $D_n(R)$ denote the K -vector space of all operators of $\text{End}_K(R)$ of order $< n$. The **ring of differential operators** of R is defined as the subring $D(R) = \bigcup_{n \geq 0} D_n(R)$.

Remark. It follows that $D_n(R) = \{P \in \text{End}_K(R) : [\cdots [P, a_1], a_2], \cdots, a_{n+1}] = 0 \text{ for any } a_i \in R\}$ and the set of order n is $D_n - D_{n-1}$.

Definition 1.2. A **derivation** of the K -algebra R is a K -linear operator D of R which satisfies Leibniz's rule:

$$D(ab) = aD(b) + bD(a) \quad \text{for every } a, b \in R$$

Let $\text{Der}_K(R)$ denote the K -vector space of all derivations of R . If $D \in \text{Der}_K(R)$ and $a \in R$, we define a new derivation aD by $(aD)(b) = aD(b)$ for every $b \in R$. The K -vector space $\text{Der}_K(R)$ is a left R -module for this action.

Proposition 1.3. Let R be a commutative K -algebra.

1. The elements of order zero are the elements of R i.e. $D^0(R) = R$
2. The operators of order ≤ 1 correspond to the elements of $\text{Der}_K(R) + R$ i.e.

$$D^1(R) \subset \text{Der}_K(R) + R$$

Proof. 2. Let $Q \in D^1(R)$ and put $P = Q - Q(1)$. Note that $P(1) = 0$ and that P has order ≤ 1 . Hence $[P, a]$ has order zero for every $a \in R$. Thus for every $b \in R$, we have that $[[P, a], b] = 0$. Writing the commutators explicitly, one obtains the equality

$$(Pa)b - (aP)b - b(Pa) + b(aP) = 0$$

Applying this operator to $1 \in R$, we end up with $P(ab) = aP(b) + bP(a)$, it follows that P is a derivation of R . But $Q = P + Q(1) \in \text{Der}_K(R) + R$, as required. \square

Proposition 1.4. *Let $P \in D^n(R)$ and $Q \in D^m(R)$, then $P \cdot Q \in D^{n+m}(R)$.*

Proof. The proof is by induction on $m + n$. If $m + n = 0$ the result is obvious. Suppose the result true whenever $m + n < k$. If $m + n = k$ and $a \in R$, we have that

$$[PQ, a] = P[Q, a] + [P, a]Q$$

The definition of order implies that $[Q, a] \in D^{m-1}(R)$ and $[P, a] \in D^{n-1}(R)$. Thus, by the induction hypothesis $P[Q, a], [P, a]Q \in D^{n+m-1}$. Hence $[PQ, a]$ belongs to D^{n+m-1} , as required. \square

§2 The Weyl Algebra

Proposition 2.1. *Every derivation of $K[X]$ is of the form $\sum_1^n f_i \partial_i$, for some $f_1, \dots, f_n \in K[X]$.*

Lemma 2.2. *Let $P \in D(K[X])$. If $[P, x_i] = 0$ for every $i = 1, \dots, n$, then $P \in K[X]$.*

Definition 2.3. *Define C_r to be the set of operators in A_n which can be written in the form $\sum_\alpha f_\alpha \partial^\alpha$ with $|\alpha| \leq r$. A simple calculation shows that*

$$C_r = C_{r+1} \cap D^r(K[X])$$

By Proposition 1.3, we have that $C_1 = \text{Der}_K(K[X]) + K[X]$ and that $C_0 = K[X]$. We will use the convention that if $k < n$ then \mathbb{N}^k is embedded in \mathbb{N}^n as the set of n -tuples whose last $n - k$ components are zero.

Lemma 2.4. *It follows from the identity $[\partial^\beta, x_n] = \beta_n \partial^{\beta - e_n}$ that $[\partial^\beta, x_n] = 0 \Leftrightarrow \beta_n = 0$. Thus $[G, x_n] = 0$ implies that G can be written as a linear combination of monomials of the form $x^\alpha \partial^\beta$ with $\beta \in \mathbb{N}^{n-1}$.*

Lemma 2.5. *Let $P_1, \dots, P_n \in C_{r-1}$ and assume that $[P_i, x_j] = [P_j, x_i]$ whenever $1 \leq i, j \leq n$. Then there exists $Q \in C_r$ such that $P_i = [Q, x_i]$, for $i = 1, \dots, n$.*

Proof. Suppose, by induction, that we have determined $Q' \in C_r$ such that $[Q', x_i] = P_i$ for $k+1 \leq i \leq n$. Write $G = [Q', x_k] - P_k$, then

$$G = \sum_{\alpha \in \mathbb{N}^k} f_\alpha \partial^\alpha.$$

since $[G, x_i] = [[Q', x_k], x_i] - [P_k, x_i] = [[Q', x_i], x_k] - [P_k, x_i] = 0$ for $k+1 \leq i \leq n$.

Now write

$$Q'' = \sum_{\alpha \in \mathbb{N}^k} (\alpha_k + 1)^{-1} f_\alpha \partial^{\alpha + e_k} \in C^r$$

We have $[Q'', x_k] = G$ and $[Q'', x_i] = 0$. Thus $[Q' - Q'', x_i] = P_i$, for $k \leq i \leq n$; and the induction is complete. \square

Theorem 2.6. *The ring of differential operators of $K[X]$ is $A_n(K)$ i.e. . Besides this, .*

$$1. D^k(K[X]) = C_k$$

$$2. D(K[X]) = A_n(K)$$

Proof. It is enough to prove that $D^k(K[X]) \subseteq C_k$. Let $P \in D(K[X])$. If $P \in D^1(K[X])$ then by Lemma 1.1, $P \in \text{Der}_K(K[X]) + K[X]$. Thus $P \in C_1$ by Proposition 1.3.

Suppose, by induction, that $D^k(K[X]) = C_k$ for $k \leq m-1$. Let $P \in D^m(K[X])$. Write $P_i = [P, x_i]$. Since P_i has order $k \leq m-1$, it follows that $P_i \in C_{m-1}$. But, for all $1 \leq i, j \leq n$,

$$[P_i, x_j] = [[P, x_i], x_j] = [[P, x_j], x_i] = [P_j, x_i].$$

Thus by Lemma 2.2 there exists $Q \in C_m$ such that $[Q, x_i] = P_i$, $1 \leq i \leq n$. Hence $[Q - P, x_i] = 0$ in $D(K[X])$. Since this holds whenever $1 \leq i \leq n$, we conclude by Lemma 2.1 that $Q - P \in K[X] = C_0$. Therefore $P \in C_m$. Hence $D^m(K[X]) \subseteq C^m$, as we wanted to prove. \square

Theorem 2.7. $\text{ord}(PQ) = \text{ord}(P) + \text{ord}(Q)$

Chapter III

Jacobian Conjecture

§1 Polynomial Maps

Definition 1.1. Let $F : K^n \rightarrow K^m$. We say that F is **polynomial** if there exist $F_1, \dots, F_m \in K[x_1, \dots, x_n]$ such that $F(p) = (F_1(p), \dots, F_m(p))$ for all $p \in K^n$.

A polynomial map is called an isomorphism or a **polynomial isomorphism** if it has an inverse which is also a polynomial map.

For the rest of the section we shall write X, Y for the spaces K^n and K^m ; and $K[X], K[Y]$ for the polynomial rings $K[x_1, \dots, x_n]$ and $K[y_1, \dots, y_m]$.

Definition 1.2. Suppose that $F : X \rightarrow Y$ is a polynomial map. We may define a map,

$$F^\sharp : K[Y] \rightarrow K[X], \quad \text{given that } g \mapsto g \circ F$$

The map F^\sharp is called the **comorphism** of F .

Suppose that a ring homomorphism $\phi : K[Y] \rightarrow K[X]$ is given. Then we may use it to construct a polynomial map from X to Y . Now let

$$\phi_\sharp : X \rightarrow Y, \quad \mathbf{x} \mapsto (\phi(y_1)(\mathbf{x}), \dots, \phi(y_m)(\mathbf{x}))$$

Theorem 1.3. Let $F : X \rightarrow Y$ and $G : Y \rightarrow Z$ be polynomial maps, then

1. $(F^\sharp)_\sharp = F$
2. $G \cdot F : X \rightarrow Z$ is a polynomial map and $(G \cdot F)^\sharp = F^\sharp \cdot G^\sharp$

Theorem 1.4. If $\phi : K[Y] \rightarrow K[X]$ and $\psi : K[Z] \rightarrow K[Y]$ are homomorphism of polynomial rings, then

1. $(\phi_\sharp)^\sharp = \phi$.
2. $(\phi \cdot \psi)_\sharp = \psi_\sharp \cdot \phi_\sharp$.

Corollary 1.5. A polynomial map $F : X \rightarrow Y$ is an isomorphism if and only if F^\sharp is an isomorphism.

§2 Jacobian Conjecture

Jacobian conjecture. Let $F : K^n \rightarrow K^n$ be a polynomial map. If $\Delta F = 1$ on K^n then F has a polynomial inverse on the whole of K^n .

Lemma 2.1. *Let $F : X \rightarrow X$ be a polynomial map and suppose that $\Delta F \neq 0$ everywhere in X . Then F^\sharp is injective.*

Proof. Suppose that F^\sharp is not injective, and choose the non-constant polynomial $g \in K[X]$ of smallest degree such that $F^\sharp(g) = 0$. Then $g \circ F = 0$. Let $g_i = \partial g / \partial x_i$ and

$$\mathbf{v} = (g_1(F_1, \dots, F_n), \dots, g_n(F_1, \dots, F_n)).$$

Hence, by the chain rule,

$$D(g \circ F) = \mathbf{v}(p) \cdot JF(p) = 0$$

for every $p \in X$. Since

$$\Delta F(p) = \det JF(p) \neq 0,$$

we conclude that $\mathbf{v}(p) = 0$ for every $p \in X$. Thus $g_i(F_1, \dots, F_n) = 0$ for $1 \leq i \leq n$. Since g is not constant, at least one of the g_i must be non-zero. But g_1 has degree smaller than g , a contradiction. \square

Proposition 2.2. *Denote by $K[F_1, \dots, F_n]$ the subalgebra of $K[X]$ generated by the coordinate functions of F . This is the image of the homomorphism F^\sharp . Thus the Jacobian conjecture may be rephrased as follows.*

Let $F : K^n \rightarrow K^n$ be a polynomial map and assume that $\Delta F = 1$ in K^n . Then $K[F_1, \dots, F_n] = K[x_1, \dots, x_n]$.

§3 Derivation

Definition 3.1. *Let D be a derivation of a K -algebra S .*

1. *It follows from Leibniz's rule that the kernel of D is a subring (subalgebra) of S , it is called the **ring of constants** of D .*
2. *The derivation D is **locally nilpotent** if given $a \in S$, there exists $k \in \mathbb{N}$ such that $D^k(a) = 0$.*
3. *Let S be a ring and D a locally nilpotent derivation. Define a map $\phi : S \rightarrow S[x]$ by the rule*

$$\phi(a) = \sum_0^\infty \frac{D^n(a)}{n!} x^n$$

for every $a \in S$. It is easy to check that ϕ is a ring homomorphism which satisfies

$$\phi \cdot D = \partial \cdot \phi$$

Proposition 3.2. *Let S be a K -algebra and D_1, \dots, D_n be commuting locally nilpotent derivations of S . Suppose that there exist $t_1, \dots, t_n \in S$ such that $D_i(t_j) = \delta_{ij}$. Then*

1. $S = R[t_1, \dots, t_n]$, where R is the ring of constants with respect to D_1, \dots, D_n ,
2. t_1, \dots, t_n are algebraically independent over R ,
3. $D_i = \partial/\partial t_i$ for $i = 1, \dots, n$.

Proof. Proof: We firstly prove it when $n = 1$. Put $\bar{S} = S/St$. Let $\rho : S \rightarrow \bar{S}[x]$ be the composition of ϕ defined above and the projection $S[x] \rightarrow \bar{S}[x]$. We want to show that ρ is an isomorphism. Note that $\rho(t) = x$.

To prove that ρ is surjective it is enough to prove that its image contains \bar{S} . Let $a \in S$. Denote by \bar{a} its image in \bar{S} . Since D is locally nilpotent, there exists $n \in \mathbb{N}$ such that $D^k(a) = 0$ for $k > n$. Thus,

$$\rho(a) = \sum_0^n \frac{\overline{D^i(a)}}{i!} x^i.$$

If $n = 0$, then $\rho(a) = \bar{a}$. If $n > 0$ put $a_0 = a$ and define $a_{j+1} = a_j - D^{n-j}(a_j)t^{n-j}/(n-j)!$, for $j = 1, \dots, n$. It is easy to show, by induction on j , that $D^k(a_j) = 0$ for $k > n - j$ and that

$$\rho(a_j) = \sum_0^{n-j} \frac{\overline{D^i(a_j)}}{i!} x^i.$$

Thus $\rho(a_n) = \bar{a}_n$. However, since $\bar{t} = 0$, we have that $\rho(a_n) = \bar{a}$. Thus ρ is surjective.

Let us prove that ρ is injective. If not, then there exists a non-zero $a \in S$ such that $\rho(a) = 0$. Thus $D^k(a) \in tS$, for every $k \in \mathbb{N}$. Hence $a = a_1 \cdot t$, for some $a_1 \in S$. Since $\rho(t) = x$, we have that $\rho(a_1) = 0$. Thus $a_1 \in tS$ and $a = a_2 \cdot t^2$, for some $a_2 \in S$. Continuing this way we conclude that t^n divides a for all $n \geq 0$. But this is impossible, unless $a = 0$. Indeed, ϕ maps t to $t + x$. Thus if t^n divides a , we also have that $\phi(t^n) = (t + x)^n$ divides $\phi(a)$ in the polynomial ring $S[x]$. Hence, if $a \neq 0$ we have that $\deg(\phi(a)) \geq n$ for every $n > 0$, which is clearly impossible. Thus $a = 0$, as required.

We conclude that the homomorphism $\rho : S \rightarrow \bar{S}[x]$ is an isomorphism. Since $\rho \cdot D = d/dx \cdot \rho$, we have that $R = \rho^{-1}(\bar{S})$. The result now follows if we recall that $\rho(t) = x$.

We proceed by induction on the number n of derivations. By Lemma 3.2, $S = R_1[t_1]$, where R_1 is the ring of constants of D_1 . But t_1 is algebraically independent over R_1 and $D_1 = d/dt_1$. Since D_1 commutes with D_i for $i > 1$, we have that $D_i(R_1) \subseteq R_1$. Thus, by the induction hypothesis, $R_1 = R[t_2, \dots, t_n]$, and the proposition follows. \square

§4 Automorphisms

Definition 4.1. Let $X = K^n$. The rational function field of $K[X]$ will be denoted by $K(X)$. Let $F : X \rightarrow X$ be a polynomial map with coordinate functions F_1, \dots, F_n . Assume that

$$\Delta = \Delta F \neq 0$$

everywhere on X .

1. Define a map $D_i : K(X) \rightarrow K(X)$ by

$$D_i(g) = \Delta^{-1} \det J(F_1, \dots, F_{i-1}, g, F_{i+1}, \dots, F_n).$$

It is easy to check that D_i is a derivation of $K(X)$.

2. Now let $K[X, \Delta^{-1}]$ be the K -subalgebra of $K(X)$ of all rational functions whose denominator is a power of Δ . Then D_i restricts to a derivation of $K[X, \Delta^{-1}]$, since $D_i(\Delta^{-1}) = -\Delta^{-2} D_i(\Delta) \in K[X, \Delta^{-1}]$

Proposition 4.2. Let R be a commutative ring and D, D' be derivations of R , then $[D, D']$ is a derivation of R .

Proposition 4.3. Let D be a K -derivation of $K[x_1, \dots, x_n]$.

1. D can be extended to the power series ring $K[[x_1, \dots, x_n]]$.
2. If Δ is a power series such that $\Delta(0) \neq 0$ then $\Delta^{-1} \cdot D$ is a derivation of the power series ring $K[[x_1, \dots, x_n]]$.

Lemma 4.4. As derivations of $K[X, \Delta^{-1}]$ the D_i satisfy:

1. $D_i(F_j) = \delta_{ij}$.
2. The D_i commute pairwise.

Proof. Note first that $\Delta(0) \neq 0$. Thus Δ is invertible as a power series and $K[X, \Delta^{-1}] \subseteq K[[X]]$. On the other hand, $\Delta \cdot D_i$ is a derivation of $K[x_1, \dots, x_n]$ which can be extended to a derivation on the power series ring $K[[X]] = K[[x_1, \dots, x_n]]$. Since Δ is invertible as a power series, then D_i can also be extended to a derivation of $K[[X]]$.

Put derivation $B = [D_i, D_j]$. We want to show that $B = 0$ on $K[X, \Delta^{-1}]$. It is enough to show that $B = 0$ on the power series ring $K[[X]]$.

Moreover $B(F_k) = 0$, for $1 \leq k \leq n$; and so B is zero in the subalgebra $K[F_1, \dots, F_n]$. But F_1, \dots, F_n are algebraically independent, by Lemma 2.2. Hence we may consider B as a derivation on the power series ring $K[[F_1, \dots, F_n]]$. By (1), B is zero on $K[[F_1, \dots, F_n]]$.

For $1 \leq i \leq n$ let $a_i = F_i(0)$. The jacobian matrices of $(F_1 - a_1, \dots, F_n - a_n)$ and F coincide. Since the latter is invertible in $K[[x_1, \dots, x_n]]$, we conclude from the local inversion theorem (see Appendix 2) that

$$K[[x_1, \dots, x_n]] = K[[F_1 - a_1, \dots, F_n - a_n]] = K[[F_1, \dots, F_n]].$$

Thus B is zero on $K[[x_1, \dots, x_n]]$, as required. \square

Definition 4.5. Let $a \in A_n$. The map $\text{ad}_a : A_n \rightarrow A_n$ is defined by

$$\text{ad}_a(b) = [a, b].$$

This is a K -linear map, but it is not a K -algebra homomorphism.

Theorem 4.6. Let $F : K^n \rightarrow K^n$ be a polynomial map and assume that $\Delta F = 1$ everywhere on K^n . If every endomorphism of A_n is an automorphism, then the Jacobian conjecture holds.

Proof. Since $\Delta F = 1$, it follows from Lemma 4.1 that D_1, \dots, D_n are derivations of $K[X]$ which satisfy

$$[D_i, F_j] = D_i(F_j) = \delta_{ij} \text{ and } [D_i, D_j] = 0$$

for $1 \leq i, j \leq n$. By, there exists an endomorphism $\phi : A_n \rightarrow A_n$ such that $\phi(x_i) = F_i$ and $\phi(\partial_i) = D_i$, for $1 \leq i \leq n$. Note that for $b \in A_n$,

$$\deg(\text{ad}_{\partial_i}(b)) = \deg[\partial_i, b] \leq \deg b - 1$$

Thus given $b \in A_n$, there exists $k \in \mathbb{N}$ such that $(\text{ad}_{\partial_1})^k(b) = 0$. Since

$$\phi(\text{ad}_{\partial_i}(b)) = \text{ad}_{D_i} \phi(b)$$

we have that $(\text{ad}_{D_i})^k(\phi(b)) = 0$. Assuming that ϕ is an automorphism, we conclude that D_i is locally nilpotent. It then follows by Proposition 3.1 that $K[F_1, \dots, F_n] = K[x_1, \dots, x_n]$, which is the Jacobian conjecture as stated in 2.3. \square

Chapter IV

Modules Over The Weyl Algebra

§1 The Polynomial Ring

Proposition 1.1. *Let R be a ring and M an irreducible left R -module.*

1. $M = Rm$ for every $0 \neq m \in M$.
2. If $0 \neq u \in M$, then $M \cong R/\text{ann}_R(u)$.
3. If R is not a division ring, then M is a torsion module.

Proposition 1.2. *The A_n -module $K[X]$ is an irreducible, torsion A_n -module. Besides this, we have isomorphism of A_n -module*

$$K[X] \cong A_n / \sum_1^n A_n \partial_i.$$

Proof. First of all 1 is clearly a generator of A_n -module $K[X]$ and the annihilator of 1 is the left ideal generated by $\partial_1, \dots, \partial_n$.

$$\begin{array}{ccc} A_n & \xrightarrow{\quad} & K[X] \\ \downarrow & \nearrow & \\ A_n / \sum_1^n A_n \partial_i & & \end{array}$$

□

Remark. Choose $g_1, \dots, g_n \in K[X]$ and consider the left ideal J of A_n generated by $\partial_1 - g_1, \dots, \partial_n - g_n$. It is easy to check that the map

$$\psi : A_n/J \longrightarrow K[X] \text{ defined by } \psi(f + J) = f$$

is an isomorphism of A_n -modules.

Proposition 1.3. *Another A_n -module is $A_n / \sum_1^n A_n \cdot x_i$.*

$$\overline{x^\alpha \partial^\beta} = (-1)^{|\alpha|} \overline{\partial^{\beta-\alpha}}$$

As a K -vector space it is isomorphic to $K[\partial] = K[\partial_1, \dots, \partial_n]$, the set of polynomials in $\partial_1, \dots, \partial_n$. Using this isomorphism, we may identify the action of A_n directly on $K[\partial]$: the ∂ 's act by multiplication, whilst x_i acting on ∂_j gives $-\delta_{ij} \cdot 1$.

§2 Twisting

Definition 2.1. Let R be a ring and M a left R module. Suppose that σ is an automorphism of R . We shall define a new left module M_σ , as follows. As an abelian group, $M_\sigma = M$. Let $a \in R$ and $u \in M$, define

$$r \cdot m = \sigma(r)m$$

It is called the **twisted module of M by σ** .

Proposition 2.2. Let R be a ring, M a left R -module and σ an automorphism of R . Then:

1. M_σ is irreducible if and only if M is irreducible.
2. M_σ is a torsion module if and only if M is a torsion module.
3. $M_\sigma \oplus M'_\sigma \cong (M \oplus M')_\sigma$.
4. If N is a submodule of M then $(M/N)_\sigma \cong M_\sigma/N_\sigma$.
5. Let J be a left ideal of R . Set $\sigma(J) = \{\sigma(r) : r \in J\}$. Then $\sigma(J)$ is a left ideal of R and $(R/J)_\sigma \cong R/\sigma^{-1}(J)$.

Proposition 2.3. The Fourier transform of $K[X]$ is $K[\partial]$.

Proof. Let \mathcal{F} be the Fourier transform automorphism of A_n . By Proposition 5.2, the twisted module $(K[X])_{\mathcal{F}}$ is isomorphic to

$$A_n / \sum_1^n A_n \cdot \mathcal{F}^{-1}(\partial_i) = A_n / \sum_1^n A_n \cdot x_i \cong K[\partial]$$

□

Theorem 2.4. For every positive integer r let σ_r be the automorphism of A_n which satisfies $\sigma_r(x_i) = x_i$ and $\sigma_r(\partial_i) = \partial_i - x_i^r$. The modules $K[X]_{\sigma_r}$ form an infinite family of pairwise nonisomorphic irreducible modules over A_n .

Proof. Let $r < t$, and suppose that there exists an isomorphism, $\phi : K[X]_{\sigma_r} \rightarrow K[X]_{\sigma_t}$. Since $K[X]_{\sigma_r}$ is irreducible, it is generated by 1. Thus ϕ is completely determined by the image of 1; say $\phi(1) = f \neq 0$. Now the equation $\phi(\partial_i \bullet 1) = \partial_i \bullet \phi(1)$ translates as the differential equation

$$\phi(\partial_i \bullet 1) = \phi((\partial_i - x_i^r)(1)) = \phi(-x_i^r) = -\phi(x_i^r) \text{ and } \partial_i \bullet \phi(1) = (\partial_i - x_i^r)(f)$$

thus

$$\frac{\partial f}{\partial x_i} = (x_i^t - x_i^r) f.$$

The left hand side of the equation has degree $\leq \deg f - 1$. Since $f \neq 0$ and $r < t$, the right hand side has degree $\deg f + t$. This is a contradiction, so the theorem is proved. \square

§3 Holomorphic Functions

Lemma 3.1. *Let $h(z)$ be the holomorphic function $\exp(\exp(z))$. For every positive integer m there exists a polynomial $F_m(x) \in \mathbb{C}[x]$ of degree m such that*

$$d^m h / dz^m = F_m(e^z) h(z).$$

Proof. The proof is by induction on m . If $m = 1$ then $dh/dz = e^z h(z)$, so we may take $F_1(x) = x$. Suppose that the result is true for $m = k$. Then

$$d^{k+1} h / dz^{k+1} = d/dz (F_k(e^z) h(z)) = (e^z F'_k(e^z) + e^z F_k(e^z)) h(z).$$

Thus we may take

$$F_{k+1}(x) = x F'_k(x) + x F_k(x)$$

which is a polynomial of degree $k + 1$. \square

Proposition 3.2. *The function $h(z) = \exp(\exp(z))$ is not a torsion element of the $A_1(\mathbb{C})$ -module $\mathcal{H}(U)$.*

Proof. Suppose that there exists a non-zero operator $P \in A_1(\mathbb{C})$ such that $P \cdot h = 0$. Write

$$P = \sum_{i=0}^m f_i(z) \partial^i$$

with $f_m(z) \neq 0$. By Lemma 6.1, we have

$$0 = P \cdot h = \sum_{i=0}^m f_i(z) F_i(e^z) h(z).$$

Since $h(z) \neq 0$, we conclude that

$$\sum_{i=0}^m f_i(z) F_i(e^z) = 0.$$

But this is impossible since F_m has degree m and $f_m(z) \neq 0$. \square

Chapter V

Differential Equations

§1 The D-Modules of Equations

Definition 1.1. Let P be an operator in A_n . This differential operator gives rise to the equation

$$P(f) = \sum_{\alpha} g_{\alpha} \partial_{\alpha}(f) = 0 \text{ in } K[X],$$

where $f \in K[X]$. More generally, if P_1, \dots, P_m are differential operators in A_n , then we have a system of differential equations

$$P_i(f) = 0, \quad i = 1, \dots, m. \quad (1)$$

The A_n -**module associated to the system (1)** is $A_n / \sum_1^m A_n P_i$. A polynomial solution of (1) is a polynomial $f \in K[X]$ which satisfies $P_i(f) = 0$, for $i = 1, \dots, m$. The set of all polynomial solutions of (1) forms a K -vector space.

Theorem 1.2. Let M be the A_n -module associated with the system (1). The K -vector space of polynomial solutions of the system (1) is isomorphic to $\text{Hom}_{A_n}(M, K[X])$.

Proof. Step1. Let f be a polynomial solution of (1). Define a map $\phi_f : A_n \rightarrow K[X]$ by the rule

$$\phi_f(D) = D(f)$$

It is easy to check that ϕ_f is an A_n -module homomorphism. Moreover, if $D \in \sum_1^m A_n P_i$, then

$$\phi_f(D) = D(f) = \sum_1^m D_i P_i(f) = 0.$$

that is $\sum_{i=1}^m A_n P_i \subseteq \ker \phi_f$. Thus ϕ_f induces an A_n -module homomorphism $\bar{\phi}_f : M \rightarrow K[X]$.

$$\begin{array}{ccc} A_n & \xrightarrow{\phi_f} & K[X] \\ \downarrow \pi & \nearrow \bar{\phi}_f & \\ M & & \end{array}$$

Step 2. Conversely, let $\psi : M \rightarrow K[X]$ be an A_n -module homomorphism. Define $f = \psi(\bar{1})$, where $\bar{1}$ is the image of 1 in M . Then for $i = 1, \dots, m$,

$$P_i(f) = P_i(\psi(\bar{1})) = \psi(P_i(\bar{1})) = 0.$$

Thus f is a polynomial solution of (1).

It is easy to check that the maps $f \mapsto \bar{\phi}_f$ and $\psi \mapsto \psi(\bar{1})$ are inverse to each other. Hence we have established the required isomorphism. \square

Definition 1.3. Let \mathcal{S} be a left A_n -module; and let $M = (P_i)$ be a finitely generated left A_n -module. We will call $\text{Hom}_{A_n}(M, \mathcal{S})$ the solution space of M in \mathcal{S} which isomorphic to K -vector space of solutions of

$$P_i(f) = 0 \text{ in } \mathcal{S}, \quad i = 1, \dots, m.$$

§2 Direct Limit of Modules

Definition 2.1. Let I be a set with a relation \leq . We say that I is **partially ordered** if \leq is reflexive, antisymmetric, and transitive.

A partially ordered set I is **directed** if, given $i, j \in I$, there exists $k \in I$ such that $i \leq k$ and $j \leq k$.

Definition 2.2. Let R be a ring and I be a directed set. Suppose that to every $i \in I$ we associate a left R -module M_i . A **directed family** of left R -modules consist of the data of

(i) For every $i \in I$, there exist a left R -module M_i

(ii) given any $i, j \in I$, satisfying $i \leq j$, there exists a homomorphism of R -modules

$$f_{ij} : M_i \longrightarrow M_j,$$

such that

(iii) For every $i \in I$, $f_{ii} = \text{id}_{M_i}$

(iv) if $i \leq j \leq k$, then

$$f_{ij} \circ f_{jk} = f_{ik}.$$

Definition 2.3. Let (M_i, f_{ij}) be a directed family of left R -modules. Denote by M the disjoint union of the modules M_i

$$M = \bigsqcup_{i \in I} M_i.$$

We define an equivalence relation in M as follows: $(u, i), (v, j) \in U$ are equivalent if and only if there exists $k \in I$ such that $i \leq k, j \leq k$ and $\pi_{ik}(u) = \pi_{jk}(v)$. The **direct limit** of the family $\{M_i : i \in I\}$, denoted by $\varinjlim M_i$, is the quotient set of M by this equivalence relation.

Remark. The operation on M is defined

$$\overline{(a_i, i)} + \overline{(a_j, j)} = \overline{(f_{ik}(a_i) + f_{jk}(a_j), k)}$$

for some $k \in I$ such that $i \leq k$ and $j \leq k$,

$$r(\overline{(a_i, i)}) = \overline{(ra_i, i)}$$

We sometimes simply write $[a_i]$ to replace $\overline{(a_i, i)}$ in M

§3 Microfunctions

Let $D(\epsilon)$ be the open disk of \mathbb{C} of centre 0 and radius ϵ , $D'(\epsilon) = D(\epsilon) \setminus \{0\}$ and $\mathcal{H}(\Omega)$ be the set of holomorphic functions in the open set $\Omega \subseteq \mathbb{C}$ viewed as an $A_1(\mathbb{C})$ -module.

Proposition 3.1. Let directed set $I = \{D(\epsilon) : \epsilon \in \mathbb{R}\}$ such that $D(\epsilon) \leq D(\epsilon')$ iff $D(\epsilon) \supseteq D(\epsilon')$, and a directed family $\{\mathcal{H}(D(\epsilon)) : D(\epsilon) \in I\}$. The homomorphisms $\tau_{\epsilon\epsilon'} : \mathcal{H}(D(\epsilon)) \rightarrow \mathcal{H}(D(\epsilon'))$ are defined by restriction of holomorphic functions.

The elements of $\mathcal{H}_0 = \varinjlim \mathcal{H}(D(\epsilon))$ are called **germs of holomorphic functions** at 0.

Proposition 3.2. The **universal cover** of $D'(\epsilon)$ is the set $\tilde{D}(\epsilon) = \{z \in \mathbb{C} : \operatorname{Re}(z) < \log(\epsilon)\}$. The projection π of $\tilde{D}(\epsilon)$ on $D'(\epsilon)$ is defined by $\pi(z) = e^z$. We have the commutative diagram

$$\begin{array}{ccc} \tilde{D}(\epsilon) & & \\ \downarrow \pi & & \\ D'(\epsilon) & \hookrightarrow & D(\epsilon) \end{array}$$

Proposition 3.3. Let $h \in \mathcal{H}(\tilde{D}(\epsilon))$. The action of a polynomial $f \in \mathbb{C}[x]$ on h is given by $f \bullet h = f(e^z)h(z)$. The operator $\partial = d/dx$ acts on h by the formula $\partial \bullet h = h'(z)e^{-z}$. Then the map

$$\pi^* : \mathcal{H}(D'(\epsilon)) \rightarrow \mathcal{H}(\tilde{D}(\epsilon))$$

defined by $\pi^*(h) = h \circ \pi$ is an injective homomorphism of $A_1(\mathbb{C})$ modules.

Definition 3.4. Let \mathcal{M}_ϵ denote the quotient module $\mathcal{H}(\tilde{D}(\epsilon))/\pi^*(\mathcal{H}(D(\epsilon)))$. If $D(\epsilon) \leq D(\epsilon')$, then $\tilde{D}(\epsilon') \subseteq \tilde{D}(\epsilon)$ and $\mathcal{H}(\tilde{D}(\epsilon)) \subseteq \mathcal{H}(\tilde{D}(\epsilon'))$. This induces a homomorphism of $A_1(\mathbb{C})$ -modules

$$\tau_{\epsilon\epsilon'} : \mathcal{M}_\epsilon \longrightarrow \mathcal{M}_{\epsilon'}.$$

Hence $\{\mathcal{M}_\epsilon : \epsilon \in \mathbb{R}\}$ is a directed family of $A_1(\mathbb{C})$ -modules, and we may take its direct limit called the **module of microfunctions**, denoted by \mathcal{M} .

Chapter VI

§1 Graded and filtered

§1.1 Graded

Definition 1.1. Let R be a ring.

1. If we have abelian group direct sum

$$R = \bigoplus_{i \geq 0} R_i$$

with $R_i R_j \subseteq R_{i+j}$ for all i, j , then R is called a **graded ring**. The R_i are called the **homogeneous components** of R .

2. A ideal (left, right, two-sided) I of R is called **graded ideal** if $I = \bigoplus_{i \geq 0} (R_i \cap I)$.
3. If I is a two-sided graded ideal of R , then quotient graded ring $R/I = \bigoplus_{i \geq 0} R_i / (R_i \cap I)$ is well-defined.

Remark. It clear that a ideal I is graded if and only if it is generated by homogeneous elements.

Definition 1.2. Let $R = \bigoplus_{i \geq 0} R_i$ and $S = \bigoplus_{i \geq 0} S_i$ be graded rings, the **graded ring homomorphism** is a ring homomorphism $f : R \rightarrow S$ such that $f(R_i) \subseteq S_i$ for all i .

Proposition 1.3. Let $R = \bigoplus_{i \geq 0} R_i$ and $S = \bigoplus_{i \geq 0} S_i$ be graded rings and $f : R \rightarrow S$ a graded ring homomorphism

1. $\ker f$ (the kernel of ring homomorphism) is a graded ideal.

Definition 1.4. Let $R = \bigoplus_{i \geq 0} R_i$ be a graded ring. A left R -module M is called a **graded left R -module** if

- (i) it is a direct sum $M = \bigoplus_{i \geq 0} M_i$ of abelian groups
- (ii) such that $R_i M_j \subseteq M_{i+j}$ for all i, j .

The M_i are called the **homogeneous components** of degree i of M .

§1.2 Filtered

Definition 1.5. Let R be a ring. A **increasing filtration** of R is a sequence of subgroups $\{F_i R\}$ of R such that

- (i) $F_0 R \subset F_1 R \subset F_2 R \subset \cdots \subset F_n R \subset \cdots \subset R$
- (ii) $\bigcup_{i \geq 0} F_i R = R$.
- (iii) $F_i R \cdot F_j R \subseteq F_{i+j} R$ for all i, j .

R is called a **filtered ring** if it has an increasing filtration.

Proposition 1.6. Let R be a ring.

1. If $R = \bigoplus_{i \geq 0} R_i$ is a graded ring, then the sequence of subgroups $\{F_k R := \bigoplus_{i=0}^k R_i\}$ is an increasing filtration of R .
2. If $\mathcal{F} = \{F_i R\}$ is a filtration of R , then

$$\mathrm{gr}_{\mathcal{F}} R := \bigoplus_{i \geq 0} F_{i+1} R / F_i R$$

is a graded ring (multiplication follows from R), called the **associated graded ring of R associated with the filtration \mathcal{F}** .

Definition 1.7. Let R be a filtered ring with filtration $\mathcal{F} = \{F_i R\}$. A left R -module M is called a **filtered left R -module** if it has a sequence of subgroups $\Gamma = \{\Gamma_i M\}$ such that

- (i) $\Gamma_0 M \subset \Gamma_1 M \subset \Gamma_2 M \subset \cdots \subset \Gamma_n M \subset \cdots \subset M$
- (ii) $\bigcup_{i \geq 0} \Gamma_i M = M$
- (iii) $F_i R \cdot \Gamma_j M \subseteq \Gamma_{i+j} M$ for all i, j .

Proposition 1.8. The **graded module of M associated with the filtration Γ** is defined by

$$\mathrm{gr}_{\Gamma} M = \bigoplus_{i \geq 0} \Gamma_{i+1} M / \Gamma_i M$$

§1.3 Induced filtrations

Definition 1.9. Let R be a filtered ring with filtration $\mathcal{F} = \{F_i R\}$ and M be a left filtered R -module.

1. If N is a submodule of M , then the **induced filtration** on N is a filtration $\Gamma' = \{\Gamma_i N\}$ defined by

$$\Gamma_i N := N \cap \Gamma_i M.$$

Remark. associated graded module $\text{gr}_{\Gamma'} N$ of N and graded module homomorphism $i_k : N \cap \Gamma_k M / N \cap \Gamma_{k-1} M \rightarrow \Gamma_k M / \Gamma_{k-1} M$

2. the **quotient filtration** $\Gamma'' = \{\Gamma_i(M/N)\}$ on M/N is defined by

$$\Gamma_i(M/N) := \Gamma_i M / (N \cap \Gamma_i M).$$

Remark. associated graded module $\text{gr}_{\Gamma''}(M/N)$ of M/N and graded module homomorphism $\pi_k : \Gamma_k M / \Gamma_{k-1} M \rightarrow (\Gamma_k M / (N \cap \Gamma_k M)) / (\Gamma_{k-1} M / (N \cap \Gamma_{k-1} M))$ (note that the right hand side is isomorphic to $\Gamma_k M / (\Gamma_{k-1} M + \Gamma_k M \cap N)$, $(A_1/B_1) / (A_2/B_2) \cong A_1 / (B_1 + A_2)$)

Proposition 1.10. Let R be a filtered ring with filtration $\mathcal{F} = \{F_i R\}$ and M be a left filtered R -module with filtration $\Gamma = \{\Gamma_i M\}$. If N is a submodule of M , then there exists a short exact sequence of graded $\text{gr}_{\mathcal{F}} R$ -modules

$$0 \longrightarrow \text{gr}_{\Gamma'} N \xrightarrow{i} \text{gr}_{\Gamma} M \xrightarrow{\pi} \text{gr}_{\Gamma''}(M/N) \longrightarrow 0.$$

is exact.

Proof. It is clear that the each

$$0 \longrightarrow \Gamma'_k \cap \Gamma'_{k-1} \xrightarrow{i_k} \Gamma_k / \Gamma_{k-1} \xrightarrow{\pi_k} \Gamma''_k / \Gamma''_{k-1} \longrightarrow 0$$

is exact in **Ab**. Thus the proposition follows. □