A Primer of Algebraic D-modules

HHH

February 2, 2026

Contents

I **2**

§1 2

§2 Algebra 2

§3 Rings of Differential Operators 2

§4 Twisting 3

II Weyl Algebra **5**

§1 Definition 5

§2 Degree and order 6

§3 Ideal Structure of A_n 7

§4 The Rings of Differential Operators of $k[X]$ 7

§5 Modules Over A_n 8

III Jacobian Conjecture **11**

§1 Polynomial Maps 11

§2 Jacobian Conjecture 12

§3 Derivation 12

§4 Automorphisms 14

IV Differential Equations **16**

§1 The D-Modules of Equations 16

§2 Microfunctions 17

V **18**

§1 Filtration and Associated Graded Rings and Modules 18

§1.1 Increasing filtration 18

§2 Good filtration 19

§3 Induced filtrations 19

§4 Dimension 20

§4.1 Upper boundness 21

§4.2 Lower boundness 22

VI Holonomic Modules **23**
§1 Definition 23
§2 Basic properties 23

VII **25**

VIII **27**
§1 Tensor Product 27
§2 External tensor product 27
§3 Filtration 28

IX **29**
§1 Inverse Image 29
§1.1 Projections 30
§1.2 Embeddings 30
§2 Direct Image 31
§2.1 31

Throughout this book, K denotes a field of characteristic zero and $k[X]$ the ring of polynomials $K[x_1, \ldots, x_n]$ in n commuting indeterminates over K.

Chapter I

§1

Proposition 1.1. *Let A be a ring and M an irreducible left A-module.*

1. $M = Am$ *for every* $0 \neq m \in M$.
2. *If* $0 \neq u \in M$, *then* $M \cong A/ann_A(u)$.
3. *If A is not a division ring, then M is a torsion module.*

§2 Algebra

§3 Rings of Differential Operators

Let A be a commutative k-algebra.

Definition 3.1. *Let A be a commutative k-algebra.*

(i) We will identify an element $a \in A$ with the operator of $\mathrm{End}_k(A)$ *defined by the rule $r \mapsto ar$, for every $r \in A$. An operator $P \in \mathrm{End}_k(A)$ has order zero if $[a, P] = 0$, for every $a \in A$.*

(ii) Suppose we have defined operators of order $< n$. An operator $P \in \mathrm{End}_k(A)$ has order n if it does not have order less than n and $[a, P]$ has order less than n for every $a \in A$.

Let $\mathrm{Diff}^n(A)$ *denote the k-vector space of all operators of* $\mathrm{End}_k(A)$ *of order $< n$. The* ***ring of differential operators*** *of A is defined as the subring* $\mathrm{Diff}(A) = \bigcup_{n \geq 0} \mathrm{Diff}^n(A)$.

Remark. *It follows that* $\mathrm{Diff}^n(A) = \{P \in \mathrm{End}_k A : [\cdots[[P, a_1], a_2], \cdots, a_{n+1}] = 0$ *for any* $a_i \in A\}$ *and the set of order n is* $\mathrm{Diff}^n - \mathrm{Diff}^{n-1}$.

Definition 3.2. *A* ***derivation*** *of the k-algebra A is a k-linear operator D of A which satisfies Leibniz's rule:*

$$D(ab) = aD(b) + bD(a) \quad \textit{for every } a, b \in A$$

Let $\mathrm{Der}_k(A)$ *denote the k-vector space of all derivations of A. If $D \in \mathrm{Der}_k(A)$ and $a \in A$, we define a new derivation aD by $(aD)(b) = aD(b)$ for every $b \in A$. The k-vector space* $\mathrm{Der}_k(A)$ *is a left A-module for this action.*

Proposition 3.3. *Let A be a commutative k-algebra.*

1. *The elements of order zero are the elements of A i.e.* $\mathrm{Diff}^0(A) = A$
2. *The operators of order ≤ 1 correspond to the elements of* $\mathrm{Der}_k(A) + A$ *.i.e.*

$$\mathrm{Diff}^1(A) \subset \mathrm{Der}_k(A) + A$$

Proof. 2. Let $Q \in \mathrm{Diff}^1(A)$ and put $P = Q - Q(1)$. Note that $P(1) = 0$ and that P has order ≤ 1. Hence $[P, a]$ has order zero for every $a \in A$. Thus for every $b \in A$, we have that $[[P, a], b] = 0$. Writing the commutators explicitly, one obtains the equality

$$(Pa)b - (aP)b - b(Pa) + b(aP) = 0$$

Applying this operator to $1 \in R$, we end up with $P(ab) = aP(b) + bP(a)$, it follows that P is a derivation of R. But $Q = P + Q(1) \in \mathrm{Der}_k(R) + R$, as required. □

Proposition 3.4. *Let $P \in \mathrm{Diff}^n(A)$ and $Q \in \mathrm{Diff}^m(A)$, then $P \cdot Q \in \mathrm{Diff}^{n+m}(A)$.*

Proof. The proof is by induction on $m + n$. If $m + n = 0$ the result is obvious. Suppose the result true whenever $m + n < k$. If $m + n = k$ and $a \in A$, we have that

$$[PQ, a] = P[Q, a] + [P, a]Q$$

The definition of order implies that $[Q, a] \in \mathrm{Diff}^{m-1}(A)$ and $[P, a] \in \mathrm{Diff}^{n-1}(A)$. Thus, by the induction hypothesis $P[Q, a], [P, a]Q \in \mathrm{Diff}^{n+m-1}(A)$. Hence $[PQ, a]$ belongs to $\mathrm{Diff}^{n+m-1}(A)$, as required. □

§4 Twisting

Definition 4.1. *Let A be a ring and M a left A module. Suppose that σ is an automorphism of A. We shall define a new left module M_σ, as follows. As an abelian group, $M_\sigma = M$. Let $a \in A$ and $u \in M$, define*

$$r \cdot m = \sigma(r)m$$

It is called the ***twisted module of*** *M* ***by*** *σ.*

Proposition 4.2. *Let A be a ring, M a left A-module and σ an automorphism of A. Then:*

1. *M_σ is irreducible if and only if M is irreducible.*
2. *M_σ is a torsion module if and only if M is a torsion module.*
3. *$M_\sigma \oplus M'_\sigma \cong (M \oplus M')_\sigma$.*
4. *If N is a submodule of M then $(M/N)_\sigma \cong M_\sigma/N_\sigma$.*

5. *Let J be a left ideal of R. Set $\sigma(J) = \{\sigma(r) : r \in J\}$. Then $\sigma(J)$ is a left ideal of J and $(R/J)_\sigma \cong R/\sigma^{-1}(J)$.*

Chapter II

Weyl Algebra

§1 Definition

Let $k[X]$ be the polynomial ring $k[x_1, \ldots, x_n]$ in n commuting indeterminates over a field k of characteristic zero. Its algebra of k-linear operators is denoted by $\mathrm{End}_k(k[X])$. Let $\hat{x}_1, \ldots, \hat{x}_n$ be the operators of $k[X]$ which are defined on a polynomial $f \in k[X]$ by the formulae $\hat{x}_i(f) = x_i \cdot f$. Similarly, $\partial_1, \ldots, \partial_n$ are the operators defined by $\partial_i(f) = \partial f / \partial x_i$.

Definition 1.1. *The n-**th Weyl algebra** $A_n(k)$ is the k-subalgebra of $\mathrm{End}_k(k[X])$ generated by the operators $\hat{x}_1, \ldots, \hat{x}_n$ and $\partial_1, \ldots, \partial_n$. For the sake of consistency, we write $A_0 = K$.*

Proposition 1.2.

$$[\partial_i, \hat{x}_j] = \delta_{ij} \cdot 1$$
$$[\partial_i, \partial_j] = [\hat{x}_i, \hat{x}_j] = 0$$
$$[\partial_i, f] = \frac{\partial f}{\partial x_i}$$

Proposition 1.3. *We have*

1. $$\partial^\beta x^\alpha = \sum_{\substack{\beta_1+\beta_2=\beta \\ \beta_i \in \mathbb{N}^n}} \binom{\beta}{\beta_1} \frac{\partial x^\alpha}{\partial x^{\beta_1}} \cdot \partial^{\beta_2}$$

   thus $\partial^\beta x^\alpha = x^\alpha \partial^\beta + D'$ where $\deg D' \leq |\alpha| + |\beta| - 2$

2. *The set $\mathbf{B} = \left\{x^\alpha \partial^\beta : \alpha, \beta \in \mathbb{N}^n\right\}$ is a basis of A_n as a vector space over K.*

Theorem 1.4. *Let the free k-algebra $k\langle z_1, \ldots, z_{2n}\rangle$ and J be the two-sided ideal generated by $[z_{i+n}, z_i] - 1$ for $i = 1, 2, \ldots, n$ and $[z_i, z_j]$ for $j \neq i + n$ and $1 \leq i, j \leq 2n$.*

$$A_n(k) \cong k\langle z_1, \ldots, z_{2n}\rangle / J$$

as k-algebras.

Proof. We may define a surjective homomorphism $\phi : k\langle z_1, \ldots, z_{2n}\rangle \to A_n$ by $\phi(z_i) = x_i$ and $\phi(z_{i+n}) = \partial_i$, for $i = 1, 2, \ldots, n$.

$$\begin{tikzcd} \{z_1, \ldots, z_n\} \arrow[r] \arrow[d, "\iota"] & A_n \\ k\langle z_1, \ldots, z_n\rangle \arrow[ur, "\phi"'] & \end{tikzcd}$$

It follows that $J \subseteq \ker\phi$. Thus ϕ induces a homomorphism $\bar{\phi} : k\langle z_1, \ldots, z_{2n}\rangle / J \longrightarrow A_n$. We may use the relations to show that every element of $k\langle z_1, \ldots, z_{2n}\rangle / J$ may be written as a linear combination of monomials of the form

$$z_1^{m_1} \ldots z_{2n}^{m_{2n}} + J$$

by 1.3 the images of these monomials under $\bar{\phi}$ form a basis of A_n as a vector space over k. In particular, the monomials must be linearly independent in $k\langle z_1, \ldots, z_{2n}\rangle / J$. Hence $\bar{\phi}$ is an isomorphism of vector spaces and, a fortiori, an isomorphism of rings. □

Corollary 1.5. *Let $m < n$ be positive integers. Choose polynomials $f_i \in k[X]$, for $1 \leq i \leq n$, as follows: if $i \leq m$, then f_i is a polynomial in the variables $x_{m+1}, \ldots, x_n$; otherwise $f_i = 0$. The map $\sigma : A_n \longrightarrow A_n$ defined by the formulae*

$$\sigma(x_i) = x_i + f_i$$
$$\sigma(\partial_i) = \partial_i - \sum_1^n \frac{\partial f_k}{\partial x_i}\partial_k$$

induce an automorphism of A_n.

§2 Degree and order

Definition 2.1. *Let $D \in A_n$. The **degree** of D is the largest length of the multi-indices $(\alpha, \beta) \in \mathbb{N}^n \times \mathbb{N}^n$ for which $x^\alpha \partial^\beta$ appears with non-zero coefficient in the canonical form of D. It is denoted by* $\deg(D)$. *As with the degree of a polynomial, we use the convention that the zero polynomial has degree $-\infty$.*

Theorem 2.2. *The degree satisfies the following properties; for $D, D' \in A_n$:*

1. $\deg(D + D') \leq \max\{\deg(D), \deg(D')\}$. *if* $\deg(D) \neq \deg(D')$ *then we have equality in the above formula.*
2. $\deg(DD') = \deg(D) + \deg(D')$.
3. $\deg[D, D'] \leq \deg(D) + \deg(D') - 2$.

Corollary 2.3. *The algebra A_n is a domain.*

Definition 2.4. *Let $D \in A_n$. If we write D in its canonical form*

$$D = \sum_{\alpha,\beta} a_{\alpha,\beta} x^\alpha \partial^\beta$$

*then the **order** of D is defined to be* $\mathrm{ord}(D) = \max\{|\beta| : a_{\alpha,\beta} \neq 0\}$.

Proposition 2.5. *The order satisfies the following property; for $P, Q \in A_n$:*

1. $\mathrm{ord}(P+Q) \leq \max\{\mathrm{ord}(P), \mathrm{ord}(Q)\}$, *with equality if* $\mathrm{ord}(P) \neq \mathrm{ord}(Q)$.
2. $\mathrm{ord}(PQ) = \mathrm{ord}(P) + \mathrm{ord}(Q)$
3. $\mathrm{ord}([P,Q]) \leq \mathrm{ord}(P) + \mathrm{ord}(Q) - 1$

§3 Ideal Structure of A_n

Theorem 3.1. *The algebra A_n is simple.*

Proof. Let I be a non-zero two-sided ideal of A_n. Choose an element $D \neq 0$ of smallest degree in I. If D has degree 0 , it is a constant, and $I = A_n$. Assume that D has degree $k > 0$ and let us aim at a contradiction.

Suppose that (α, β) is a multi-index of length k. If $x^\alpha \partial^\beta$ is a summand of D with non-zero coefficient and $\beta_i \neq 0$, then $[x_i, D]$ is non-zero and has degree $k-1$. Since I is a two-sided ideal of A_n, we have that $[x_i, D] \in I$. But this contradicts the minimality of D. Thus $\beta = (0, \ldots, 0)$. Since $k > 0$, we must have that $\alpha_i \neq 0$, for some $i = 1, 2, \ldots, n$. Hence $[\partial_i, D]$ is a non-zero element of I of degree $k-1$, and once again we have a contradiction. □

Corollary 3.2. *Every (ring, algebra) endomorphism of A_n is injective.*

Theorem 3.3. *Every left ideal of A_n is generated by two elements.*

§4 The Rings of Differential Operators of $k[X]$

We denote the set of element of A_n with order less than or equal to r by $C^r(k[X])$, simplified as C^r.

Proposition 4.1. *Every derivation of $k[X]$ is of the form $\sum_1^n f_i \partial_i$, for some $f_1, \ldots, f_n \in k[X]$.*

Lemma 4.2. *Let $P \in \mathrm{Diff}(k[X])$. If $[P, x_i] = 0$ for every $i = 1, \ldots, n$, then $P \in k[X]$.*

Lemma 4.3. $C_r = C_{r+1} \cap \mathrm{Diff}^r(k[X])$

Lemma 4.4. *It follows from the identity $\left[\partial^\beta, x_n\right] = \beta_n \partial^{\beta - e_n}$ that $\left[\partial^\beta, x_n\right] = 0 \Leftrightarrow \beta_n = 0$. Thus $[G, x_n] = 0$ implies that G can be written as a linear combination of monomials of the form $x^\alpha \partial^\beta$ with $\beta \in \mathbb{N}^{n-1}$.*

Lemma 4.5. *Let $P_1, \ldots, P_n \in C_{r-1}$ and assume that $[P_i, x_j] = [P_j, x_i]$ whenever $1 \leq i, j \leq n$. Then there exists $Q \in C_r$ such that $P_i = [Q, x_i]$, for $i = 1, \ldots, n$.*

Proof. Suppose, by induction, that we have determined $Q' \in C_r$ such that $[Q', x_i] = P_i$ for $k+1 \leq i \leq n$. Write $G = [Q', x_k] - P_k$, then

$$G = \sum_{\alpha \in \mathbb{N}^k} f_\alpha \partial^\alpha.$$

since $[G, x_i] = [[Q', x_k], x_i] - [P_k, x_i] = [[Q', x_i], x_k] - [P_k, x_i] = 0$ for $k+1 \leq i \leq n$.

Now write

$$Q'' = \sum_{\alpha \in \mathbb{N}^k} (\alpha_k + 1)^{-1} f_\alpha \partial^{\alpha + e_k} \in C^r$$

We have $[Q'', x_k] = G$ and $[Q'', x_i] = 0$. Thus $[Q' - Q'', x_i] = P_i$, for $k \leq i \leq n$; and the induction is complete. □

Theorem 4.6. *The ring of differential operators of $k[X]$ is $A_n(K)$ i.e. . Besides this, .*

1. $\mathrm{Diff}^k(k[X]) = C_k$
2. $\mathrm{Diff}(k[X]) = A_n(K)$

Proof. It is enough to prove that $\mathrm{Diff}^k(k[X]) \subseteq C_k$. Let $P \in \mathrm{Diff}(k[X])$. If $P \in \mathrm{Diff}^1(k[X])$ then by Lemma 1.1, $P \in \mathrm{Der}_k(k[X]) + k[X]$. Thus $P \in C_1$ by Proposition 1.3.

Suppose, by induction, that $\mathrm{Diff}^k(k[X]) = C_k$ for $k \leq m-1$. Let $P \in \mathrm{Diff}^m(k[X])$. Write $P_i = [P, x_i]$. Since P_i has order $k \leq m-1$, it follows that $P_i \in C_{n-1}$. But, for all $1 \leq i, j \leq n$,

$$[P_i, x_j] = [[P, x_i], x_j] = [[P, x_j], x_i] = [P_j, x_i].$$

Thus by Lemma 2.2 there exists $Q \in C_m$ such that $[Q, x_i] = P_i, 1 \leq i \leq n$. Hence $[Q - P, x_i] = 0$ in $\mathrm{Diff}(k[X])$. Since this holds whenever $1 \leq i \leq n$, we conclude by Lemma 2.1 that $Q - P \in k[X] = C_0$. Therefore $P \in C_m$. Hence $\mathrm{Diff}^m(k[X]) \subseteq C^m$, as we wanted to prove. □

§5 Modules Over A_n

Noted that a A_n-module M satisfies the following relation:

$$[\partial_i, x_j]\, m = \delta_{ij} m$$
$$[x_i, x_j]\, m = \left[\partial_{x_i}, \partial_{x_j}\right] m = 0$$

Proposition 5.1. *The left A_n-module $k[X]$ is an irreducible, torsion A_n-module. Besides this, we have isomorphism of A_n-module*

$${}_{A_n}k[X] \cong A_n / \sum_1^n A_n \partial_i.$$

Proof. First of all 1 is clearly a generator of ${}_{A_n}k[X]$ and the annihilator of 1 is the left ideal generated by $\partial_1, \ldots, \partial_n$.

$$\begin{tikzcd} A_n \arrow[r] \arrow[d] & k[X] \\ A_n/\sum_1^n A_n\partial_i \arrow[ur] & \end{tikzcd}$$

□

Remark. *Choose $g_1, \ldots, g_n \in k[X]$ and consider the left ideal J of A_n generated by $\partial_1 - g_1, \ldots, \partial_n - g_n$. It is easy to check that the map*

$$\psi : A_n/J \longrightarrow k[X] \text{ defined by } \psi(f+J) = f$$

is an isomorphism of A_n-modules.

Proposition 5.2. *Another A_n-module is $A_n/\sum_1^n A_n \cdot x_i$.*

$$\overline{x^\alpha \partial^\beta} = (-1)^{|\alpha|}\, \overline{\partial^{\beta-\alpha}}$$

As a k-vector space it is isomorphic to $K[\partial] = K[\partial_1, \ldots, \partial_n]$, the set of polynomials in $\partial_1, \ldots, \partial_n$. Using this isomorphism, we may identify the action of A_n directly on $K[\partial]$: the ∂ 's act by multiplication, whilst x_i acting on ∂_j gives $-\delta_{ij} \cdot 1$.

Proposition 5.3. *The Fourier transform of $k[X]$ is $K[\partial]$.*

Proof. Let $\mathcal{F}$ be the Fourier transform automorphism of A_n. By proposition 4.2 , the twisted module $(k[X])_{\mathcal{F}}$ is isomorphic to

$$A_n/\sum_1^n A_n \cdot \mathcal{F}^{-1}(\partial_i) = A_n/\sum_1^n A_n \cdot x_i \cong K[\partial]$$

□

Theorem 5.4. *For every positive integer r let σ_r be the automorphism of A_n which satisfies $\sigma_r(x_i) = x_i$ and $\sigma_r(\partial_i) = \partial_i - x_i^r$. The modules $k[X]_{\sigma_r}$ form an infinite family of pairwise nonisomorphic irreducible modules over A_n.*

Proof. Let $r < t$, and suppose that there exists an isomorphism, $\phi : k[X]_{\sigma_r} \longrightarrow k[X]_{\sigma_t}$. Since $k[X]_{\sigma_r}$ is irreducible, it is generated by 1 . Thus ϕ is completely determined by the image of 1 ; say $\phi(1) = f \neq 0$. Now the equation $\phi(\partial_i \bullet 1) = \partial_i \bullet \phi(1)$ translates as the differential equation

$$\phi(\partial_i \bullet 1) = \phi((\partial_i - x_i^r)(1)) = \phi(-x_i^r) = -\phi(x_i^r) \text{ and } \partial_i \bullet \phi(1) = (\partial_i - x^r)(f)$$

thus

$$\frac{\partial f}{\partial x_i} = \left(x_i^t - x_i^r\right) f.$$

The left hand side of the equation has degree $\leq \deg f - 1$. Since $f \neq 0$ and $r < t$, the right hand side has degree $\deg f + t$. This is a contradiction, so the theorem is proved. $\square$

Proposition 5.5. *The function $h(z) = \exp(\exp(z))$ is not a torsion element of the $A_1(\mathbb{C})$-module $\mathcal{H}(U)$.*

Proof. Suppose that there exists a non-zero operator $P \in A_1(\mathbb{C})$ such that $P \cdot h = 0$. Write

$$P = \sum_{i=0}^{m} f_i(z)\partial^i$$

with $f_m(z) \neq 0$. By Lemma 6.1, we have

$$0 = P \cdot h = \sum_{i=0}^{m} f_i(z) F_i\left(e^z\right) h(z).$$

Since $h(z) \neq 0$, we conclude that

$$\sum_{i=0}^{m} f_i(z) F_i\left(e^z\right) = 0.$$

But this is impossible since F_m has degree m and $f_m(z) \neq 0$. $\square$

Chapter III

Jacobian Conjecture

For the rest of the section we shall write X, Y for the vector spaces k^n and k^m; and $k[X], k[Y]$ for the polynomial rings $k[x_1, \ldots, x_n]$ and $k[y_1, \ldots, y_m]$.

§1 Polynomial Maps

Definition 1.1. *Let $F : k^n \to k^m$. We say that F is **polynomial** if there exist $F_1, \ldots, F_m \in k[x_1, \ldots, x_n]$ such that $F(p) = (F_1(p), \ldots, F_m(p))$ for all $p \in k^n$.*

*A polynomial map is called an isomorphism or a **polynomial isomorphism** if it has an inverse which is also a polynomial map.*

Definition 1.2. *Suppose that $F : X \to Y$ is a polynomial map. We may define a map,*

$$F^\sharp : k[Y] \to k[X], \quad \text{given that } g \mapsto g \circ F$$

*The map $F^\sharp$ is called the **comorphism** of F.*

Suppose that a ring homomorphism $\phi : K[Y] \to k[X]$ is given. Then we may use it to construct a polynomial map from X to Y. Now let

$$\phi_\sharp : X \to Y, \quad \mathbf{x} \mapsto (\phi(y_1)(\mathbf{x}), \ldots, \phi(y_m)(\mathbf{x}))$$

Theorem 1.3. *Let $F : X \to Y$ and $G : Y \to Z$ be polynomial maps, then*

1. $\left(F^\sharp\right)_\sharp = F$
2. *$G \cdot F : X \to Z$ is a polynomial map and $(G \cdot F)^\sharp = F^\sharp \cdot G^\sharp$*

Theorem 1.4. *If $\phi : K[Y] \to k[X]$ and $\psi : K[Z] \to K[Y]$ are homomorphism of polynomial rings, then*

1. $(\phi_\sharp)^\sharp = \phi$.
2. $(\phi \cdot \psi)_\sharp = \psi_\sharp \cdot \phi_\sharp$.

Corollary 1.5. *A polynomial map $F : X \to Y$ is an isomorphism if and only if $F^\sharp$ is an isomorphism.*

§2 Jacobian Conjecture

Jacobian conJecture. Let $F : K^n \to K^n$ be a polynomial map. If $\Delta F = 1$ on K^n then F has a polynomial inverse on the whole of K^n.

Lemma 2.1. *Let $F : X \to X$ be a polynomial map and suppose that $\Delta F \neq 0$ everywhere in X. Then $F^\sharp$ is injective.*

Proof. Suppose that $F^\sharp$ is not injective, and choose the non-constant polynomial $g \in k[X]$ of smallest degree such that $F^\sharp(g) = 0$. Then $g \circ F = 0$. Let $g_i = \partial g/\partial x_i$ and

$$\mathbf{v} = (g_1(F_1, \ldots, F_n), \ldots, g_n(F_1, \ldots, F_n)).$$

Hence, by the chain rule,

$$D(g \circ F) = \mathbf{v}(p) \cdot JF(p) = 0$$

for every $p \in X$. Since

$$\Delta F(p) = \det JF(p) \neq 0,$$

we conclude that $\mathbf{v}(p) = 0$ for every $p \in X$. Thus $g_i(F_1, \ldots, F_n) = 0$ for $1 \leq i \leq n$. Since g is not constant, at least one of the g_i must be non-zero. But g_1 has degree smaller than g, a contradiction. □

Proposition 2.2. *Denote by $K[F_1, \ldots, F_n]$ the subalgebra of $k[X]$ generated by the coordinate functions of F. This is the image of the homomorphism $F^\sharp$. Thus the Jacobian conjecture may be rephrased as follows.*

Let $F : K^n \to K^n$ be a polynomial map and assume that $\Delta F = 1$ in K^n. Then $K[F_1, \ldots, F_n] = K[x_1, \ldots, x_n]$.

§3 Derivation

Definition 3.1. *Let D be a derivation of a k-algebra S.*

1. *It follows from Leibniz's rule that the kernel of D is a subring (subalgebra) of S, it is called the **ring of constants of** D.*

2. *The derivation D is **locally nilpotent** if given $a \in S$, there exists $k \in \mathbb{N}$ such that $D^k(a) = 0$.*

3. *Let S be a ring and D a locally nilpotent derivation. Define a map $\phi : S \longrightarrow S[x]$ by the rule*

$$\phi(a) = \sum \frac{D^n(a)}{n!} x^n$$

for every $a \in S$. It is easy to check that ϕ is a ring homomorphism which satisfies

$$\phi \cdot D = \partial \cdot \phi$$

Proposition 3.2. *Let S be a k-algebra and $D_1, \ldots, D_n$ be commuting locally nilpotent derivations of S. Suppose that there exist $t_1, \ldots, t_n \in S$ such that $D_i(t_j) = \delta_{ij}$. Then*

1. *$S = R[t_1, \ldots, t_n]$, where R is the ring of constants with respect to $D_1, \ldots, D_n$,*
2. *$t_1, \ldots, t_n$ are algebraically independent over R,*
3. *$D_i = \partial/\partial t_i$ for $i = 1, \ldots, n$.*

Proof. Proof: We firstly prove it whene $n = 1$. Put $\bar{S} = S/St$. Let $\rho : S \longrightarrow \bar{S}[x]$ be the composition of ϕ defined above and the projection $S[x] \longrightarrow \bar{S}[x]$. We want to show that ρ is an isomorphism. Note that $\rho(t) = x$.

To prove that ρ is surjective it is enough to prove that its image contains $\bar{S}$. Let $a \in S$. Denote by $\bar{a}$ its image in $\bar{S}$. Since D is locally nilpotent, there exists $n \in \mathbb{N}$ such that $D^k(a) = 0$ for $k > n$. Thus,

$$\rho(a) = \sum_0^n \frac{\overline{D^i(a)}}{i!} x^i.$$

If $n = 0$, then $\rho(a) = \bar{a}$. If $n > 0$ put $a_0 = a$ and define $a_{j+1} = a_j - D^{n-j}(a_j)\, t^{n-j}/(n-j)!$, for $j = 1, \ldots, n$. It is easy to show, by induction on j, that $D^k(a_j) = 0$ for $k > n - j$ and that

$$\rho(a_j) = \sum_0^{n-j} \frac{\overline{D^i(a_j)}}{i!} x^i.$$

Thus $\rho(a_n) = \bar{a}_n$. However, since $\bar{t} = 0$, we have that $\rho(a_n) = \bar{a}$. Thus ρ is surjective.

Let us prove that ρ is injective. If not, then there exists a non-zero $a \in S$ such that $\rho(a) = 0$. Thus $D^k(a) \in tS$, for every $k \in \mathbb{N}$. Hence $a = a_1 \cdot t$, for some $a_1 \in S$. Since $\rho(t) = x$, we have that $\rho(a_1) = 0$. Thus $a_1 \in tS$ and $a = a_2 \cdot t^2$, for some $a_2 \in S$. Continuing this way we conclude that t^n divides a for all $n \geq 0$. But this is impossible, unless $a = 0$. Indeed, ϕ maps t to $t + x$. Thus if t^n divides a, we also have that $\phi(t^n) = (t + x)^n$ divides $\phi(a)$ in the polynomial ring $S[x]$. Hence, if $a \neq 0$ we have that $\deg(\phi(a)) \geq n$ for every $n > 0$, which is clearly impossible. Thus $a = 0$, as required.

We conclude that the homomorphism $\rho : S \longrightarrow \bar{S}[x]$ is an isomorphism. Since $\rho \cdot D = d/dx \cdot \rho$, we have that $R = \rho^{-1}(\bar{S})$. The result now follows if we recall that $\rho(t) = x$.

We proceed by induction on the number n of derivations. By Lemma 3.2, $S = R_1[t_1]$, where R_1 is the ring of constants of D_1. But t_1 is algebraically independent over R_1 and $D_1 = d/dt_1$. Since D_1 commutes with D_i for $i > 1$, we have that $D_i(R_1) \subseteq R_1$. Thus, by the induction hypothesis, $R_1 = R[t_2, \ldots t_n]$, and the proposition follows. □

§4 Automorphisms

Definition 4.1. *Let $X = K^n$. The rational function field of $k[X]$ will be denoted by $K(X)$. Let $F : X \to X$ be a polynomial map with coordinate functions $F_1, \ldots, F_n$. Assume that*

$$\Delta = \Delta F \neq 0$$

everywhere on X.

1. *Define a map $D_i : K(X) \to K(X)$ by*

$$D_i(g) = \Delta^{-1} \det J\left(F_1, \ldots, F_{i-1}, g, F_{i+1}, \ldots, F_n\right).$$

   It is easy to check that D_i is a derivation of $K(X)$.

2. *Now let $K\left[X, \Delta^{-1}\right]$ be the K-subalgebra of $K(X)$ of all rational functions whose denominator is a power of Δ. Then D_i restricts to a derivation of $K\left[X, \Delta^{-1}\right]$, since $D_i\left(\Delta^{-1}\right) = -\Delta^2 D_i(\Delta) \in K\left[X, \Delta^{-1}\right]$*

Proposition 4.2. *Let R be a commutative ring and D, D' be derivations of R, then $[D, D']$ is a derivation of R.*

Proposition 4.3. *Let D be a K-derivation of $K\left[x_1, \ldots, x_n\right]$.*

1. *D can be extended to the power series ring $K\left[\left[x_1, \ldots, x_n\right]\right]$.*

2. *If Δ is a power series such that $\Delta(0) \neq 0$ then $\Delta^{-1} \cdot D$ is a derivation of the power series ring $K\left[\left[x_1, \ldots, x_n\right]\right]$.*

Lemma 4.4. *As derivations of $K\left[X, \Delta^{-1}\right]$ the D_i satisfy:*

1. *$D_i\left(F_j\right) = \delta_{ij}$.*

2. *The D_i commute pairwise.*

Proof. Note first that $\Delta(0) \neq 0$. Thus Δ is invertible as a power series and $K\left[X, \Delta^{-1}\right] \subseteq K[[X]]$. On the other hand, $\Delta \cdot D_i$ is a derivation of $K\left[x_1, \ldots, x_n\right]$ which can be extended to a derivation on the power series ring $K[[X]] = K\left[\left[x_1, \ldots, x_n\right]\right]$. Since Δ is invertible as a power series, then D_i can also be extended to a derivation of $K[[X]]$.

Put derivation $B = [D_i, D_j]$. We want to show that $B = 0$ on $K\left[X, \Delta^{-1}\right]$. It is enough to show that $B = 0$ on the power series ring $K[[X]]$.

Moreover $B\left(F_k\right) = 0$, for $1 \leq k \leq n$; and so B is zero in the subalgebra $K\left[F_1, \ldots, F_n\right]$. But $F_1, \ldots, F_n$ are algebraically independent, by Lemma 2.2. Hence we may consider B as a derivation on the power series ring $K\left[\left[F_1, \ldots, F_n\right]\right]$. By (1), B is zero on $K\left[\left[F_1, \ldots, F_n\right]\right]$.

For $1 \leq i \leq n$ let $a_i = F_i(0)$. The jacobian matrices of $(F_1 - a_1, \ldots, F_n - a_n)$ and F coincide. Since the latter is invertible in $K[[x_1, \ldots, x_n]]$, we conclude from the local inversion theorem (see Appendix 2) that

$$K[[x_1, \ldots, x_n]] = K[[F_1 - a_1, \ldots, F_n - a_n]] = K[[F_1, \ldots, F_n]].$$

Thus B is zero on $K[[x_1, \ldots, x_n]]$, as required. □

Definition 4.5. *Let $a \in A_n$. The map $\mathrm{ad}_a : A_n \to A_n$ is defined by*

$$\mathrm{ad}_a(b) = [a, b].$$

This is a K-linear map, but it is not a K-algebra homomorphism.

Theorem 4.6. *Let $F : K^n \to K^n$ be a polynomial map and assume that $\Delta F = 1$ everywhere on K^n. If every endomorphism of A_n is an automorphism, then the Jacobian conjecture holds.*

Proof. Since $\Delta F = 1$, it follows from Lemma 4.1 that $D_1, \ldots, D_n$ are derivations of $k[X]$ which satisfy

$$[D_i, F_j] = D_i(F_j) = \delta_{ij} \text{ and } [D_i, D_j] = 0$$

for $1 \leq i, j \leq n$. By, there exists an endomorphism $\phi : A_n \to A_n$ such that $\phi(x_i) = F_i$ and $\phi(\partial_i) = D_i$, for $1 \leq i \leq n$. Note that for $b \in A_n$,

$$\deg(\mathrm{ad}_{\partial_i}(b)) = \deg[\partial_i, b] \leq \deg b - 1$$

Thus given $b \in A_n$, there exists $k \in \mathbb{N}$ such that $(\mathrm{ad}_{\partial_1})^k(b) = 0$. Since

$$\phi(\mathrm{ad}_{\partial_i}(b)) = \mathrm{ad}_{D_i}\phi(b)$$

we have that $(\mathrm{ad}_{D_i})^k(\phi(b)) = 0$. Assuming that ϕ is an automorphism, we conclude that D_i is locally nilpotent. It then follows by Proposition 3.1 that $K[F_1, \ldots, F_n] = K[x_1, \ldots, x_n]$, which is the Jacobian conjecture as stated in 2.3. □

Chapter IV

Differential Equations

§1 The D-Modules of Equations

Definition 1.1. *Let $\mathcal{S}$ be a left A_n-module and $P_1, P_2, \ldots, P_m \in A_n$; consider the system of differential equations*

$$P_i(f) = 0 \text{ in } \mathcal{S}, \quad i = 1, \ldots, m.$$

*And let $M := (P_i)$ be a finitely generated left A_n-module called A_n**-module associated to the system**. We will call* $\mathrm{Hom}_{A_n}(M, \mathcal{S})$ *the **solution space** of M in $\mathcal{S}$*

Remark. *The set of all solutions of the system is a k-vector space.*

Theorem 1.2. *Let M be the A_n-module associated with the system (1). The k-vector space of solutions of the system (1) is isomorphic to* $\mathrm{Hom}_{A_n}(M, \mathcal{S})$.

Proof. Step1. Let f be a solution of (1). Define a map $\phi_f : A_n \to \mathcal{S}$ by the rule

$$\phi_f(D) = D(f)$$

It is easy to check that ϕ_f is an A_n-module homomorphism. Moreover, if $D \in \sum_1^m A_n P_i$, then

$$\phi_f(D) = D(f) = \sum_1^m D_i P_i(f) = 0.$$

that is $\sum_1^m A_n P_i \subseteq \ker \phi_f$. Thus ϕ_f induces an A_n-module homomorphism $\bar{\phi}_f : M \to \mathcal{S}$.

$$\begin{tikzcd} A_n \arrow[r, "\phi_f"] \arrow[d, "\pi"] & \mathcal{S} \\ M \arrow[ur, "\bar{\phi}_f"] & \end{tikzcd}$$

Step 2. Conversely, let $\psi : M \to \mathcal{S}$ be an A_n-module homomorphism. Define $f = \psi(\overline{1})$,

where $\overline{1}$ is the image of 1 in M. Then for $i = 1, \ldots, m$,

$$P_i(f) = P_i(\psi(\overline{1})) = \psi(P_i(\overline{1})) = 0.$$

Thus f is a solution of (1).

It is easy to check that the maps $f \mapsto \bar{\phi}_f$ and $\psi \mapsto \psi(\overline{1})$ are inverse to each other. Hence we have established the required isomorphism. $\square$

§2 Microfunctions

Let $D(\epsilon)$ be the open disk of $\mathbb{C}$ of centre 0 and radius ϵ, $D'(\epsilon) = D(\epsilon)\backslash 0$ and $\mathcal{H}(\Omega)$ be the set of holomorphic functions in the open set $\Omega \subseteq \mathbb{C}$ viewed as an $A_1(\mathbb{C})$-module.

Proposition 2.1. *Let directed set $I = \{D(\epsilon) : \epsilon \in \mathbb{R}\}$ such that $D(\epsilon) \leq D(\epsilon')$ iif $D(\epsilon) \supseteq D(\epsilon')$, and a directed family $\{\mathcal{H}(D(\epsilon)) : D(\epsilon) \in I\}$. The homomorphisms $\tau_{\epsilon\epsilon'} : \mathcal{H}(D(\epsilon)) \longrightarrow \mathcal{H}(D(\epsilon'))$ are defined by restriction of holomorphic functions.*

*The elements of $\mathcal{H}_0 = \varinjlim \mathcal{H}(D(\epsilon))$ are called **germs of holomorphic functions** at 0.*

Proposition 2.2. *The **universal cover** of $D'(\epsilon)$ is the set $\tilde{D}(\epsilon) = \{z \in \mathbb{C} : \mathrm{Re}(z) < \log(\epsilon)\}$. The projection π of $\tilde{D}(\epsilon)$ on $D'(\epsilon)$ is defined by $\pi(z) = e^z$. We have the commutative diagram*

$$\begin{tikzcd} \tilde{D}(\epsilon) \arrow[d, "\pi"] & \\ D'(\epsilon) \arrow[r, hook] & D(\epsilon) \end{tikzcd}$$

Proposition 2.3. *Let $h \in \mathcal{H}(\tilde{D}(\epsilon))$. The action of a polynomial $f \in \mathbb{C}[x]$ on h is given by $f \bullet h = f(e^z)\, h(z)$. The operator $\partial = d/dx$ acts on h by the formula $\partial \bullet h = h'(z)e^{-z}$. Then the map*

$$\pi^* : \mathcal{H}(D'(\epsilon)) \longrightarrow \mathcal{H}(\tilde{D}(\epsilon))$$

defined by $\pi^*(h) = h \circ \pi$ is an injective homomorphism of $A_1(\mathbb{C})$ modules.

Definition 2.4. *Let $\mathcal{M}_\epsilon$ denote the quotient module $\mathcal{H}(\tilde{D}(\epsilon))/\pi^*(\mathcal{H}(D(\epsilon)))$. If $D(\epsilon) \leq D(\epsilon')$, then $\tilde{D}(\epsilon') \subseteq \tilde{D}(\epsilon)$ and $\mathcal{H}(\tilde{D}(\epsilon)) \subseteq \mathcal{H}\left(\tilde{D}(\epsilon')\right)$. This induces a homomorphism of $A_1(\mathbb{C})$-modules*

$$\tau_{\epsilon\epsilon'} : \mathcal{M}_\epsilon \longrightarrow \mathcal{M}_{\epsilon'}.$$

*Hence $\{\mathcal{M}_\epsilon : \epsilon \in \mathbb{R}\}$ is a directed family of $A_1(\mathbb{C})$-modules, and we may take its direct limit called the **module of microfunctions**, denoted by $\mathcal{M}$.*

Chapter V

§1 Filtration and Associated Graded Rings and Modules

§1.1 Increasing filtration

Definition 1.1. *Let A be a ring. A **increasing filtration** of A is a sequence of subgroups $\{F_iA\}$ of A such that*

(i) $F_0A \subset F_1A \subset F_2A \subset \cdots \subset F_nA \subset \cdots \subset A$

(ii) $\bigcup_{i\geq 0} F_iA = A$.

(iii) $F_iA \cdot F_jA \subseteq F_{i+j}A$ *for all* i, j.

*A is called a **filtered ring** if it has an increasing filtration.*

Proposition 1.2. *Let A be a ring.*

1. *If $A = \bigoplus_{i\geq 0} A_i$ is a graded ring, then the sequence of subgroups $\left\{F_kA := \bigoplus_{i=0}^{k} A_i\right\}$ is an increasing filtration of A.*

2. *If $\mathcal{F} = \{F_iA\}$ is a filtration of A, then*

$$\mathrm{gr}_{\mathcal{F}} A := \bigoplus_{i\geq 0} F_{i+1}A/F_iA$$

*is a graded ring (multiplication follows from A), called the **associated graded ring of** A **associated with the filtration** $\mathcal{F}$.*

Definition 1.3. *Let A be a filtered ring with increasing filtration $\mathcal{F} = \{F_iA\}$. A left A-module M is called a **filtered left** A**-module** if it has a sequence of subgroups $\Gamma = \{\Gamma_iM\}$ such that*

(i) $\Gamma_0M \subset \Gamma_1M \subset \Gamma_2M \subset \cdots \subset \Gamma_nM \subset \cdots \subset M$

(ii) $\bigcup_{i\geq 0} \Gamma_iM = M$

(iii) $F_iA \cdot \Gamma_jM \subseteq \Gamma_{i+j}M$ *for all* i, j.

Proposition 1.4. *Let M be a left A-module where A is a filtered ring with increasing filtration $\mathcal{F} = \{F_i A\}$.*

1. *If $M = \bigoplus_{i \geq 0} M_i$ is a graded A-module, then the sequence of subgroups $\Gamma_k M := \bigoplus_{i=0}^{k} M_i$ is an increasing filtration of M.*

2. *If $\Gamma = \{\Gamma_i M\}$ is a filtration of M, then The **graded module of** M **associated with the filtration** Γ is defined by*
$$\mathrm{gr}_\Gamma M = \bigoplus_{i \geq 0} \Gamma_{i+1} M / \Gamma_i M$$
which is a graded $\mathrm{gr}_{\mathcal{F}} A$-module.

Remark. *In (1), the $\mathrm{gr}_{\mathcal{F}} M \cong M$ (as graded modules).*

But in (2), let $A = \mathbb{Z}$ with trivial filtration and $M = \mathbb{Z}_{p^2}$ with filtration $F_0 M = 0, F_1 M = p\mathbb{Z}_{p^2}, F_2 M = M, \ldots$, then $\mathrm{gr}_{\mathcal{F}} M \cong \mathbb{Z}_p^2 \ncong M$ in ${}_{\mathbb{Z}}\mathbf{Mod}$.

The functor $\mathrm{gr}(-) : {}_R\mathbf{FiltMod} \to {}_R\mathbf{GrMod}$ is not faithful

§2 Good filtration

Definition 2.1. *Let A be a filtered ring with an increasing filtration $\{F_n A\}$, and M a filtered A-module with an increasing filtration $\{F_n M\}$. The filtration of M is called a **good filtration** if $\mathrm{gr}(M)$ is finitely generated over $\mathrm{gr}(A)$.*

§3 Induced filtrations

Definition 3.1. *Let A be a filtered ring with filtration $\mathcal{F} = \{F_i A\}$ and M be a left filtered A-module with filtration $\Gamma = \{\Gamma_i\}$.*

1. *If N is a submodule of M, then the **induced filtration** on N is a filtration $\Gamma' = \{\Gamma_i N\}$ defined by*
$$\Gamma'_i := N \cap \Gamma_i.$$

   Remark. *associated graded module $\mathrm{gr}_{\Gamma'} N$ of N and graded module homomorphism $i_k : N \cap \Gamma_k / N \cap \Gamma_{k-1} \to \Gamma_k / \Gamma_{k-1}$*

2. *the **quotient filtration** $\Gamma'' = \{\Gamma''_i\}$ on M/N is defined by*
$$\Gamma''_i := \Gamma_i / (N \cap \Gamma_i).$$

   Remark. *associated graded module $\mathrm{gr}_{\Gamma''}(M/N)$ of M/N and graded module homomorphism $\pi_k : \Gamma_k M / \Gamma_{k-1} M \to (\Gamma_k M / (N \cap \Gamma_k M)) \,/\, (\Gamma_{k-1} M / (N \cap \Gamma_{k-1} M))$ (note that the right hand side is isomorphic to $\Gamma_k M / (\Gamma_{k-1} M + \Gamma_k M \cap N)$, $(A_1/B_1) \,/\, (A_2/B_2) \cong A_1 / (B_1 + A_2)$)*

Proposition 3.2. *Then there exists a short exact sequence of graded* $\mathrm{gr}_{\mathcal{F}} R$*-modules*

$$0 \longrightarrow \mathrm{gr}_{\Gamma'} N \xrightarrow{i} \mathrm{gr}_{\Gamma} M \xrightarrow{\pi} \mathrm{gr}_{\Gamma''}(M/N) \longrightarrow 0.$$

is exact.

Proof. It is clear that the each

$$0 \longrightarrow \Gamma'_k \cap \Gamma'_{k-1} \xrightarrow{i_k} \Gamma_k/\Gamma_{k-1} \xrightarrow{\pi_k} \Gamma''_k/\Gamma''_{k-1} \longrightarrow 0$$

is exact in **Ab**. Thus the proposition follows. □

§4 Dimension

Definition 4.1. *Let* M *be a finitely generated left* A_n*-module. Suppose that* Γ *is a good filtration of* M *with respect to the Bernstein filtration* $\mathcal{B}$*.*

1. *Denote by* $\chi(t, \Gamma, M)$ *the* ***Hilbert polynomial*** *of the graded module* $gr^{\Gamma} M$ *over* S_n*. We have*

$$\chi(t, \Gamma, M) = \sum_{0}^{t} \dim_k (\Gamma_i/\Gamma_{i-1}) = \dim_k (\Gamma_t).$$

   The dimension $d(M)$ *of* M *is defined the* $\deg \chi(t, \Gamma, M)$*.*

2. *Let* $a_{d(M)}$ *be the leading coefficient of* $\chi(t, \Gamma, M)$*. The* ***multiplicity*** *of* M *is* $m(M) = d! a_{d(M)}$*.*

Remark. *The dimension and multiplicity of* M *do not depend on the choice of the good filtration* Γ*.*

Example 4.2. *First let* M *be the left* A_n*-module* A_n*. The Bernstein filtration* $\mathcal{B}$ *is a good filtration of* M *and it is possible to calculate* $\chi(t, \mathcal{B}, M)$ *explicitly in this case.*

$$\chi(t, \mathcal{B}, M) = \#\left\{x^{\alpha}\partial^{\beta} : \alpha, \beta \in \mathbb{N}^n, |\alpha| + |\beta| \leq t\right\} = \binom{t+2n}{2n}$$

that is

$$\frac{(t+2n)(t+2n-1)\cdots(t+1)}{(2n)!}$$

Thus $d(A_n) = 2n$ *and* $m(A_n) = 1$*.*

Example 4.3. *Another* A_n*-module that we know very well is* $k[X] = K[x_1, \ldots, x_n]$ *with good filtration* $\Gamma_i := \{f(x_1, \ldots, x_n) : \deg f \leq i\}$*. It is easy to show that*

$$\chi(t, \mathcal{B}, k[X]) = \dim_k \Gamma_t = \binom{n+t}{n}$$

Hence $d(k[X]) = n$ *and* $m(k[X]) = 1$*.*

§4.1 Upper boundness

Theorem 4.4. *Let M be a finitely generated left A_n-module and N a submodule of M.*

1. $d(M) = \max\{d(N), d(M/N)\}$.
2. *If* $d(N) = d(M/N)$ *then* $m(M) = m(N) + m(M/N)$.

Proof. Let M be a finitely generated left A_n-module and Γ a good filtration of M with respect to $\mathcal{B}$. Let N be a submodule of M. Denote by Γ' and Γ'' the filtrations induced by Γ in N and M/N, we have an exact sequence of S_n-modules, namely

$$0 \to gr_{\Gamma'} N \to gr_{\Gamma} M \to gr_{\Gamma''} M/N \to 0.$$

Since Γ is good, $gr_{\Gamma} M$ is finitely generated. But S_n is a noetherian ring. Hence $gr_{\Gamma'} N$ and $gr_{\Gamma''}(M/N)$ are also finitely generated. Therefore both Γ' and Γ'' are good filtrations. On the other hand, since the sequence of vector spaces

$$0 \to \Gamma'_m/\Gamma'_{m-1} \to \Gamma_m/\Gamma_{m-1} \to \Gamma''_m/\Gamma''_{m-1} \to 0$$

is exact, we have that

$$\dim_k \Gamma'_m/\Gamma'_{m-1} + \dim_k \Gamma''_m/\Gamma''_{m-1} = \dim_k \Gamma_m/\Gamma_{m-1}.$$

Summing these terms for $m = 0, 1, \ldots, s$ and assuming that $s \gg 0$ one obtains

$$\chi(s, \Gamma', N) + \chi(s, \Gamma'', M/N) = \chi(s, \Gamma, M)$$

The result now follows from the properties of polynomials. □

Corollary 4.5. *Let $M_1, \ldots, M_m$ be finitely generated left A_n-modules, and $M = M_1 \oplus \cdots \oplus M_m$.*

1. $d(M) = \max\{d(M_1), \ldots, d(M_m)\}$.
2. *If* $d(M) = d(M_i)$ *for* $1 \le i \le k$, *then* $m(M) = \sum_1^k m(M_i)$.

Corollary 4.6. *Let M be a finitely generated A_n-module. Then $d(M) \le 2n$.*

Proof. Suppose that M is generated by r elements. Then there exists a surjective homomorphism $\phi : {A_n}^{\oplus r} \to M$. It follows from the theorem that $d\left({A_n}^{\oplus r}\right) = \max\{d(M), d(\ker\phi)\}$, thus $d(M) \le d\left({A_n}^{\oplus r}\right) = 2n$. □

Corollary 4.7. *Let I be a non-zero left ideal of A_n. Then $d(A_n/I) \le 2n - 1$.*

Proof. First consider the case of a cyclic left ideal. Let $d \in A_n$, and put $I = A_n d$. Then we have an exact sequence

$$0 \to A_n \xrightarrow{\theta} A_n \to A_n/A_n d \to 0$$

where the map θ is defined by $\theta(a) = ad$, for every $a \in A_n$. Suppose, by contradiction, that $d\left(A_n/A_n d\right) = 2n$. Then

$$m\left(A_n\right) = m\left(A_n\right) + m\left(A_n/A_n d\right).$$

Since $m\left(A_n\right) = 1$ and the multiplicity is a positive number, this equation is impossible. Hence $d\left(A_n/A_n d\right) \leq 2n - 1$.

Now for the general case. Let I be a non-zero left ideal of A_n and choose $0 \neq d \in I$. Since $A_n d \subseteq I$, we have that A_n/I is a quotient of $A_n/A_n d$. Since the latter has dimension $\leq 2n-1$, so does A_n/I. $\square$

§4.2 Lower boundness

Theorem 4.8 (Bernstein's Inequality)**.** *If M is a finitely generated non-zero left A_n-module, then $d(M) \geq n$.*

Chapter VI

Holonomic Modules

§1 Definition

Definition 1.1. *A finitely generated left A_n-module is* ***holonomic*** *if it is zero, or if it has dimension n.*

Proposition 1.2. *Let n be a positive integer.*

1. *Submodules and quotients of holonomic A_n-modules are holonomic.*
2. *Finite direct sums of holonomic A_n-modules are holonomic.*

Corollary 1.3. *Finitely generated torsion A_1-modules are holonomic.*

Proposition 1.4. *Holonomic A_n-modules are torsion modules.*

§2 Basic properties

Theorem 2.1. *Holonomic modules are artinian.*

Proof. Let M be a holonomic left A_n-module. Suppose that M has a descending chain of submodules

$$M = N_0 \supseteq N_1 \supseteq N_2 \supseteq \cdots \supseteq N_r$$

it follows that $m\left(N_i\right) = m\left(N_{i+1}\right) + m\left(N_i/N_{i+1}\right)$. Putting these together, we get that

$$m(M) = \sum_{0}^{r-1} m\left(N_i/N_{i+1}\right) + m\left(N_r\right) \geq r$$

Hence M cannot have a descending chain of more than r submodules. In particular M cannot have an infinite descending chain. $\square$

Remark. *Noted that the regular left module A_n is not artinian (ring A_n is not left Artinian). It is easy to construct an infinite descending chain; take for instance*

$$A_n x_n \supseteq A_n x_n^2 \supseteq A_n x_n^3 \supseteq \dots .$$

Corollary 2.2. *Every holonomic A_n-module has finite length that cannot exceed its multiplicity.*

Corollary 2.3. *Every irreducible holonomic A_n-module has multiplicity* 1.

Chapter VII

Lemma 0.1. *Let Ω and Γ be both good filtration of finitely generated left A_n-module M. Then*

$$\mathrm{Rad}\left(\mathrm{Ann}\left(\mathrm{gr}_\Gamma M\right)\right) = \mathrm{Rad}\left(\mathrm{Ann}\left(\mathrm{gr}_\Omega M\right)\right)$$

in S_n.

Definition 0.2. *The ideal*

$$\mathcal{I}(M) = \mathrm{rad}(\mathrm{ann}(M, F))$$

*is called the **characteristic ideal** of M. Thus the affine variety*

$$\mathrm{Ch}(M) = \mathcal{Z}(\mathcal{I}(M)) \subseteq \mathbb{C}^{2n}$$

*is called the **characteristic variety** of M.*

Remark. *Since $\mathcal{I}(M)$ is a homogeneous ideal, the variety $\mathrm{Ch}(M)$ is homogeneous. This means that if $p \in \mathrm{Ch}(M)$ and $\lambda \in \mathbb{C}$, then $\lambda p \in \mathrm{Ch}(M)$. Note that $\mathrm{Ch}(M)$ is a subvariety of $\mathbb{C}^{2n}$, since S_n is a polynomial ring in $2n$ variables!*

Proposition 0.3. *Let M be a finitely generated left A_n-module and N a submodule of M. Then*

$$\mathrm{Ch}(M) = \mathrm{Ch}(N) \cup \mathrm{Ch}(M/N).$$

Definition 0.4. *Let J be an ideal of $S_n = \mathbb{C}\left[y_1, \ldots, y_{2n}\right]$ and put algebric set $V = \mathcal{Z}(J)$.*

1. *If p is a point of V then the **Zariski tangent space** of V at p is the linear subspace of $\mathbb{C}^{2n}$ defined by the equations*

$$\sum_1^{2n} \frac{\partial F}{\partial y_i}(p) y_i = 0$$

   for all $F \in J$. This space is denoted by $T_p(V)$; it is a complex vector subspace of $\mathbb{C}^{2n}$.

2. *The **dimension** of V equals $\inf\{\dim_{\mathbb{C}} T_p(V) : p \in V\}$.*

Remark. *Actually one need not look at every point of* V*. If* p *is a non-singular point of* V*, then* $\dim(V) = \dim_{\mathbb{C}} T_p(V)$*. This is equivalent to the definition in terms of heights of prime ideals, see [Hartshorne 77, Ch. 1, Exercise 5.10].*

Theorem 0.5. *Let* M *be a finitely generated left module over* A_n*. Then* $\dim \mathrm{Ch}(M) = d(M)$.

Chapter VIII

Recall that we have the notations $X = k^n, Y = k^m, k[X] = k[x_1, \ldots, x_n], Y = k[y_1, \ldots, y_m]$. Let $F : X \to Y$ be a polynomial map, its comorphism $F^\sharp : k[Y] \to k[X]$ is an algebra homomorphism defined by

$$F^\sharp(\varphi) := \varphi \circ F = \varphi\left(F_1, \ldots, F_m\right)$$

§1 Tensor Product

Proposition 1.1. *Let A be a k-algebra and let A_1 and A_2 be subalgebras of A. Suppose that*

(i) $A = A_1 A_2$,

(ii) $[A_1, A_2] = 0$,

(iii) there exist k-bases $\{a_i\}$ and $\{b_j\}$ of A_1 and A_2, respectively, such that $\{a_i b_j\}$ is a k-basis of A.

Then $A \cong A_1 \otimes_k A_2 := A_1 \widehat{\otimes} A_2$ as k-algebras.

Corollary 1.2. *The following isomorphisms are induced by the multiplication map:*

1. $k[X] \otimes_k k[Y] \cong k[X, Y]$.
2. $A_m \otimes_k A_n \cong A_{m+n}$.

§2 External tensor product

Definition 2.1. *Let A, B be k-algebras. Suppose that M is a left A-module and that N is a left B-module. Then $M \otimes_k N$ is a left $A \otimes_k B$-module with action defined by*

$$(a \otimes b)(u \otimes v) = au \otimes bv.$$

*We use the notation $M \boxtimes N$ to denote this module, called the **external tensor product** of M and N.*

Proposition 2.2. *If M and N are finitely generated modules, then so is $M \boxtimes N$.*

Theorem 2.3. *Let M be a finitely generated left A_m-module and N a finitely generated left A_n-module. Then*

1. $d(M \boxtimes N) = d(M) + d(N)$,
2. $m(M \boxtimes N) \leq m(M)m(N)$.

§3 Filtration

Lemma 3.1. *The Bernstein filtration of A_{m+n} satisfies*

$$B_t\left(A_{m+n}\right) = \sum_{p+q=t} B_p\left(A_m\right) B_q\left(A_n\right)$$

Lemma 3.2. *The k-vector spaces*

$$\Gamma_t(M \boxtimes N) := \sum_{p+q=t} \Gamma_p(M) \otimes_k \Gamma_q(N)$$

form a good filtration of $M \boxtimes N$ as an A_{m+n}-module.

Lemma 3.3.

$$\Gamma_t(M \boxtimes N)/\Gamma_{t-1}(M \boxtimes N) \cong \bigoplus_{p+q=t} \left(\Gamma_p(M)/\Gamma_{p-1}(M)\right) \otimes_k \left(\Gamma_q(N)/\Gamma_{q-1}(N)\right)$$

Remark. *it can be simplified as*

$$\mathrm{gr}_t(M \boxtimes N) \cong \mathrm{gr}_t(M) \boxtimes \mathrm{gr}_t(N)$$

where $\mathrm{gr}_t(M) := \Gamma_t(M)/\Gamma_{t-1}(M)$

Chapter IX

§1 Inverse Image

Definition 1.1. *Let polynomial map*

$$F : X \to Y$$

be given. Its comorphism is the algebra homomorphism

$$F^\sharp : k[Y] \to k[X]$$

defined by $\varphi \mapsto \varphi \circ F$. *It may be used to make* $k[Y]$*-modules* M *into* $k[X]$*-modules:*

$$F^\star : {}_{k[Y]}\mathbf{Mod} \to {}_{k[X]}\mathbf{Mod}$$

defined by $F^\star(M) := k[X] \otimes_{k[Y]} M$ *called the* ***inverse image*** *of* M *by the polynomial map* F.

Definition 1.2. *If* M *is a left* A_m*-module, it follows that* M *is also a left* $k[Y]$*-module. We now turn* $k[X]$*-module* $F^\star(M)$ *into a* A_n*-module*

$$F^\star = {}_{A_m}\mathbf{Mod} \to {}_{A_n}\mathbf{Mod}$$

by giving the recipe for the action

$$x_i\,(q \otimes u) = (x_i q) \otimes u$$

$$\partial_{x_i}(q \otimes u) = \partial_{x_i}(q) \otimes u + \sum_{k=1}^{m} q\partial_{x_i}\,(F_k) \otimes \partial_{y_k} u$$

for $i = 1, \ldots, n$.

Remark. *The inverse image functor* $F^* \cong \mathcal{D}_{X\to Y} \otimes_{A_m}$, *where* $\mathcal{D}_{X\to Y} := F^* A_m = k[X] \otimes_{k[Y]} A_m$ *is a* A_n*-*A_m *bimodule.*

Theorem 1.3. *Let* $F : X \to Y$ *and* $G : Y \to Z$ *be polynomial maps and* M *be an* A_r*-module. Then*

$$F^\star G^\star M \cong (GF)^\star M$$

Corollary 1.4. *Let $F : X \to X$ be a polynomial isomorphism and G its inverse. If M is a left A_n-module, then $F^\star G^\star M \cong M$.*

Theorem 1.5 (Preservation of holonomy)**.** *Let $F : X \to Y$ be a polynomial map. If M is a holonomic A_m-module, then $F^\star M$ is a holonomic A_n-module.*

§1.1 Projections

Example 1.6. *The second example is the projection $\pi : X \times Y \to Y$ onto the second coordinate. The comorphism $\pi^\sharp$ maps a polynomial $p \in k[Y]$ to itself, but considered as an element of $k[X, Y]$.*

Let M be a module over $k[Y]$, then

$$\pi^\star(M) = k[X, Y] \otimes_{k[Y]} M \cong k[X] \widehat{\otimes} M$$

as $k[X, Y]$-modules by

$$q_\alpha x^\alpha \otimes u \longmapsto q_\alpha \otimes q_\alpha u$$

Theorem 1.7. *Let M be a finitely generated left A_m-module and π the projection defined above. Then:*

1. *$\pi^\star M$ is a finitely generated A_{m+n}-module.*
2. *$d(\pi^\star M) = n + d(M)$.*
3. *$m(\pi^\star M) \leq m(M)$.*

§1.2 Embeddings

Definition 1.8. *We begin by considering the standard embedding*

$$\iota : X \to X \times Y$$

defined by $\iota(X) = (X, 0)$, where 0 denotes the origin of Y. The comorphism

$$\iota^\sharp : K[X, Y] \to k[X]$$

is defined by $\iota^\sharp(g(X, Y)) = g(X, 0)$ (gives a isomorpjism $k[X, Y]/(Y) \cong k[X]$). It may be used to make $k[X, Y]$-modules M into a $k[X]$-module:

$$\iota^\star : {}_{k[X,Y]}\mathbf{Mod} \to {}_{k[X]}\mathbf{Mod}$$

defined by $\iota^\star(M) := k[X] \otimes_{k[X,Y]} M$.

Now consider left A_{n+m}-module M, the A_n-modules $\iota^\star(M)$ is defined by the action

$$\partial_{x_i}(q \otimes u) = \frac{\partial q}{\partial x_i} \otimes u + q \otimes \partial_{x_i} u$$

Proposition 1.9. *Let M be a left A_{n+m}-module. Then*

$$k[X] \otimes_{k[X,Y]} M \cong {}_{A_n}M/\left(Y\right){}_{A_n}M$$

as A_n-modules.

Proof. If $u \in M$, let $\bar{u}$ denote its image in $M/(Y)M$. Define the map

$$\phi : \iota^\star M \to M/(Y)M$$

by $\phi(q \otimes u) = q\bar{u}$. It is a homomorphism of $k[X]$-modules. It suffices to show that the action of the derivations $\partial_{x_1}, \ldots, \partial_{x_n}$ is compatible with ϕ and ϕ is injective. □

§2 Direct Image

Definition 2.1. *Let $F : X \to Y$ be a polynomial map. Let M be a left A_m-module. The **direct image** of N by F is the right A_m-module defined by*

$$F_\star M := M \otimes_{A_n} D_{X \to Y}$$

Theorem 2.2. *Let $F : X \to Y$ and $G : Y \to Z$ be two polynomial maps. Let M be a left A_n-module. Then*

$$(GF)_\star M \cong G_\star F_\star M.$$

Theorem 2.3. *Let $F : X \to Y$ be a polynomial map. If M is a holonomic A_n-module, then $F_\star M$ is a holonomic A_m-module.*

§2.1

Theorem 2.4. *Let $\iota : X \to X \times Y$ be the standard embedding. If M is a finitely generated left A_n-module, then:*

1. *$\iota_\star M$ is a finitely generated left A_{m+n}-module.*
2. *$d\left(\iota_\star M\right) = m + d(M)$.*
3. *$m\left(\iota_\star M\right) \leq m(M)$.*