

Commutative Algebra

HHH

July 7, 2026

Contents

I	1
§1 Ideals quotient	1
§2 Contraction and extension of ideal	1
§3 Nil and nilpotent ideals	2
§4	2
II Radical Ideals	3
III Fractions and Localization	5
§1 Rings of Quotient	5
§2 Extensions and Contractions in ring of fractions	6
§3 Fractions of modules	7
§4 Localization and Local rings	8
§5 Local properties	8
IV Chain Condition	9
§1	9
§2 Normal series and Composition Series of Modules	10
V Noetherian Ringa and Modules	12
§1 Basic Criterion	12
§1.1 General case	12
§1.2 Commutative case	12
§2 Primary Decomposition	14
VI Artinian	17
§1 Artinian Rings	17
§2 Commutative Artinian Rings	18
VIIThe Structure of Rings and Modules	20
§1 Simplicity	20
§1.1 Primitivity	21
§1.2 Jacobson Density Theorem	21

§1.3	Artinian simple ring	23
§2	Semisimplicity	24
§2.1	Structure of semisimple rings	24
§2.2	Structure of modules over semisimple rings	26
§3	Jacobson Radical	26
§3.1	Jacobson radical	26
§3.2	Nakayama's Lemma	28
§4	Characterizations of semisimple rings	28
§5	Algebra	28
VIII	Integral	30
§1	Rings Extensions	30
§1.1	Integral extension	31
§2	Discrete Valuation Ring	32
§3	Dedekind Domain	33
§3.1	Unique factorization of ideals	33
§3.2	The ideal class group	34
§4	Discrete valuations	35
§5	Integral closures of Dedekind domains	36
IX	Completions	37
§1		37
§2	Graded rings and graded modules	37
§2.1	Graded ring	37
§2.2	Graded modules	38
§2.3	Graded algebra	39
§3	Filtrations	39
§4	Associated graded ring	40
X	Dimension	42
§1	Hilbert Functions	42
§2	Krull Dimension	43
XI		45
§1		45
§2	Spectrum of a Ring	45

Chapter I

§1 Ideals quotient

Definition 1.1. If $\mathfrak{a}, \mathfrak{b}$ are ideals in a commutative ring R , their **ideal quotient** is

$$(\mathfrak{a} : \mathfrak{b}) = \{x \in R : x\mathfrak{b} \subseteq \mathfrak{a}\}$$

which is an ideal. If $\mathfrak{b}$ is a principal ideal (x) , we shall write $(\mathfrak{a} : x)$ in place of $(\mathfrak{a} : (x))$.

Proposition 1.2. Let R be a commutative ring. Then

1. $\mathfrak{a} \subseteq (\mathfrak{a} : \mathfrak{b})$
2. $(\mathfrak{a} : \mathfrak{b})\mathfrak{b} \subseteq \mathfrak{a}$
3. $((\mathfrak{a} : \mathfrak{b}) : \mathfrak{c}) = (\mathfrak{a} : \mathfrak{b}\mathfrak{c}) = ((\mathfrak{a} : \mathfrak{c}) : \mathfrak{b})$
4. $(\bigcap_i \mathfrak{a}_i : \mathfrak{b}) = \bigcap_i (\mathfrak{a}_i : \mathfrak{b})$
5. $(\mathfrak{a} : \sum_i \mathfrak{b}_i) = \bigcap_i (\mathfrak{a} : \mathfrak{b}_i)$.

§2 Contraction and extension of ideal

Definition 2.1. Let R be a ring and $f : A \rightarrow B$ be a ring homomorphism,

1. the **extension** of ideal $\mathfrak{a}$ of A is the ideal generated by $f(\mathfrak{a})$ in B , denoted by $\mathfrak{a}^e$.
2. the **contraction** of $\mathfrak{b}$ is $f^{-1}(\mathfrak{b})$, denoted by $\mathfrak{b}^c$.

Epecially if A be a subring of B and $i : A \rightarrow B$, the contraction of ideal of B is $A \cap \mathfrak{b}$.

Proposition 2.2. .

1. $\mathfrak{a} \subseteq \mathfrak{a}^{ec}, \mathfrak{b} \supseteq \mathfrak{b}^{ce};$
2. $\mathfrak{b}^c = \mathfrak{b}^{cec}, \mathfrak{a}^e = \mathfrak{a}^{ece};$

3. If $\mathcal{C}$ is the set of all contracted ideals in A and if $\mathcal{E}$ is the set of all extended ideals in B , then $\mathcal{C} = \{\mathfrak{a} \mid \mathfrak{a}^{ec} = \mathfrak{a}\}$, $\mathcal{E} = \{\mathfrak{b} \mid \mathfrak{b}^{ce} = \mathfrak{b}\}$, and $\mathfrak{a} \mapsto \mathfrak{a}^e$ is a bijective map, whose inverse is $\mathfrak{b} \mapsto \mathfrak{b}^c$.

Proposition 2.3. .

$$\begin{aligned} (\mathfrak{a}_1 + \mathfrak{a}_2)^e &= \mathfrak{a}_1^e + \mathfrak{a}_2^e & (\mathfrak{b}_1 + \mathfrak{b}_2)^c &\supseteq \mathfrak{b}_1^c + \mathfrak{b}_2^c, \\ (\mathfrak{a}_1 \cap \mathfrak{a}_2)^e &\subseteq \mathfrak{a}_1^e \cap \mathfrak{a}_2^e & (\mathfrak{b}_1 \cap \mathfrak{b}_2)^c &= \mathfrak{b}_1^c \cap \mathfrak{b}_2^c, \\ (\mathfrak{a}_1 \mathfrak{a}_2)^e &= \mathfrak{a}_1^e \mathfrak{a}_2^e & (\mathfrak{b}_1 \mathfrak{b}_2)^c &\supseteq \mathfrak{b}_1^c \mathfrak{b}_2^c, \\ (\mathfrak{a}_1 : \mathfrak{a}_2)^e &\subseteq (\mathfrak{a}_1^e : \mathfrak{a}_2^e) & (\mathfrak{b}_1 : \mathfrak{b}_2)^c &\subseteq (\mathfrak{b}_1^c : \mathfrak{b}_2^c) \\ \text{Rad}(\mathfrak{a})^e &\subseteq \text{Rad}(\mathfrak{a}^e) & \text{Rad}(\mathfrak{b})^c &= \text{Rad}(\mathfrak{b}^c) \end{aligned}$$

if $\mathfrak{b}$ is a prime ideal in B , then so $\mathfrak{b}^c$.

The set $\mathcal{C}$ is closed under the other three operations, and $\mathcal{E}$ is closed under sum and product.

§3 Nil and nilpotent ideals

Definition 3.1. Let A be a ring and $\mathfrak{a}$ be an (left, right, two-sided) ideal of A .

1. $\mathfrak{a}$ of A is **nil** if every element of $\mathfrak{a}$ is a nilpotent element;
2. $\mathfrak{a}$ is **nilpotent** if $\mathfrak{a}^n = 0$ for some integer n .

Theorem 3.2. Let A be a ring.

1. If $a \in A$ is nilpotent, a is both left and right quasiregular with quasi inverse $r = -a + a^2 - a^3 + \cdots + (-1)^{n-1}a^{n-1}$
2. Every nil left ideal is contained in $J(A)$.
3. Thus every nil ring is a radical ring.

Proposition 3.3. If R is a left [resp. right] Noetherian ring, then every nil ideal is nilpotent (Exercise 16).

§4

Definition 4.1. Let A be a ring and M be a left A -module.

1. The element $a \in A$ induces a homomorphism

$$a_M : M \rightarrow M, \quad x \mapsto a \cdot x$$

called the **principal homomorphism associated with** a on M .

2. We say that a_M is **locally nilpotent** if for every $x \in M$, there exists a positive integer $n(x)$ such that $a^{n(x)} \cdot x = 0$.

Remark. It is equivalent that

Chapter II

Radical Ideals

Throughout this chapter, let A be a commutative ring.

Definition 0.1. Let A be a commutative ring. If $\mathfrak{a}$ is any ideal of A , the ideal

$$\text{Rad}(\mathfrak{a}) := \{x \in A : x^n \in \mathfrak{a} \text{ for some } n \in \mathbb{Z}_{\geq 1}\}$$

is called **radical** of $\mathfrak{a}$, sometimes denoted by $\sqrt{\mathfrak{a}}$. The radical of 0 (the set of all nilpotent elements in A) is called **nilradical** of A , denoted by $\text{Nil}(A)$.

Proposition 0.2. Then

1. $\mathfrak{a} \subset r(\mathfrak{a})$
2. $r(r(\mathfrak{a})) = r(\mathfrak{a})$
3. $r(\mathfrak{a}\mathfrak{b}) = r(\mathfrak{a} \cap \mathfrak{b}) = r(\mathfrak{a}) \cap r(\mathfrak{b})$
4. thus $r(\mathfrak{a}_1\mathfrak{a}_2 \cdots \mathfrak{a}_n) = r(\bigcap \mathfrak{a}_i) = \bigcap r(\mathfrak{a}_i)$ and $r(\mathfrak{a}^n) = r(\mathfrak{a})$
5. $r(\mathfrak{a} + \mathfrak{b}) = r(r(\mathfrak{a}) + r(\mathfrak{b}))$
6. if $\mathfrak{p}$ is prime in R , $r(\mathfrak{p}^n) = r(\mathfrak{p}) = \mathfrak{p}$ for all $n > 0$.
7. $r(\mathfrak{a}) = (1) \Leftrightarrow \mathfrak{a} = (1)$

Proposition 0.3. If S is a multiplicative subset which is disjoint from an ideal $\mathfrak{a}$, then there exists a ideal $\mathfrak{p}$ which is maximal in $\mathcal{S} = \{\mathfrak{b} : \mathfrak{a} \subset \mathfrak{b} \text{ and } \mathfrak{b} \cap S = \emptyset\}$. Furthermore, any such ideal $\mathfrak{p}$ is prime.

Proof. Since $S \neq \emptyset$ and every ideal in $\mathcal{S}$ is properly contained in A , set $\mathcal{S}$ is partially ordered by inclusion. By Zorn's Lemma there is an ideal $\mathfrak{p}$ which is maximal in $\mathcal{S}$.

Let $\mathfrak{a}_1, \mathfrak{a}_2$ be ideals of A such that $\mathfrak{a}_1\mathfrak{a}_2 \subset \mathfrak{p}$. If $\mathfrak{a}_1 \not\subset \mathfrak{p}$ and $\mathfrak{a}_2 \not\subset \mathfrak{p}$, then each of the ideals $\mathfrak{p} + \mathfrak{a}_1$ and $\mathfrak{p} + \mathfrak{a}_2$ properly contains $\mathfrak{p}$ and hence must meet S . Consequently, for some $p_i \in \mathfrak{p}, a_i \in \mathfrak{a}_i$.

$$p_1 + a_1 = s_1 \in S \quad \text{and} \quad p_2 + a_2 = s_2 \in S$$

Thus $s_1 s_2 = p_1 p_2 + p_1 a_2 + a_1 p_2 + a_1 a_2 \in \mathfrak{p} + \mathfrak{a}_1 \mathfrak{a}_2 \subset \mathfrak{p}$. This is a contradiction since $s_1 s_2 \in S$ and $S \cap \mathfrak{p} = \emptyset$. Therefore $\mathfrak{a}_1 \subset \mathfrak{p}$ or $\mathfrak{a}_2 \subset \mathfrak{p}$, whence $\mathfrak{p}$ is prime. $\square$

Theorem 0.4. *Let A be a commutative rng and an ideal $\mathfrak{a}$.*

1. *If $\pi : A \rightarrow A/\mathfrak{a}$ is the canonical projection, then $\text{Rad}(\mathfrak{a}) = \pi^{-1}(\text{Nil}(A/\mathfrak{a}))$*
2. *The radical of an ideal $\mathfrak{a}$ is the intersection of the prime ideals which contain $\mathfrak{a}$, that is,*

$$\text{Rad}(\mathfrak{a}) = \bigcap_{\substack{\mathfrak{a} \subset \mathfrak{p}, \\ \mathfrak{p} \in \text{Spec } A}} \mathfrak{p}$$

Proof. It is clear that

$$\text{Rad}(\mathfrak{a}) \subset \bigcap_{\substack{\mathfrak{a} \subset \mathfrak{p} \\ \mathfrak{p} \text{ is prime}}} \mathfrak{p} := \tilde{\mathfrak{p}}$$

If $S = \tilde{\mathfrak{p}} - \text{Rad}(\mathfrak{a})$ is nonempty, whence is a multiplicative subset of R (verify that $x, y \in S \Rightarrow xy \in S$) and disjoint from $\text{Rad}(\mathfrak{a})$, there exist a prime ideal $\mathfrak{p}'$ that contains $\text{Rad}(\mathfrak{a})$ and disjoint from S by 0.3. But $\tilde{\mathfrak{p}} \subset \mathfrak{p}'$ by the definition of $\tilde{\mathfrak{p}}$, this is a contradiction. $\square$

Proposition 0.5. *If A is a commutative ring with identity $\neq 0$, then $A^\times + \text{Nil}(A) \subset A^\times$.*

Chapter III

Fractions and Localization

§1 Rings of Quotient

A is a commutative ring with identity throughout this section unless otherwise stated.

Definition 1.1. Let A be a commutative ring. A subset S called a **multiplicative subset** of A if S is a submonoid of $(A, \times)$.

Remark. In general, we always assume that $0 \notin S$.

Definition 1.2. Let S be a multiplicative subset of A and

1. The relation defined on the set $A \times S$ by

$$(a, s) \sim (a', s') \Leftrightarrow s_1 (as' - a's) = 0 \text{ for some } s_1 \in S$$

is an equivalence relation and the equivalence class containing the element (a, s) is denoted by a/s .

2. $S^{-1}R$ is a commutative ring with identity $1/1$, where addition and multiplication are defined by

$$r/s + r'/s' = (rs' + r's)/ss' \quad \text{and} \quad (r/s)(r'/s') = rr'/ss'$$

is called the **ring of fractions** of R by S .

Remark. The map $\varphi_S : A \rightarrow S^{-1}A$ given by $a \mapsto a/1$ is a well-defined homomorphism of rings and $\varphi(S) \subset (S^{-1}A)^\times$.

Theorem 1.3 (Universal property). Let $\mathcal{C}$ be the category

- whose objects are ring-homomorphisms (commutative rings with identity)

$$f : A \rightarrow B$$

such that for every $s \in S$, the element $f(s)$ is invertible in B .

- If $f : A \rightarrow B$ and $f' : A \rightarrow B'$ are two objects of $\mathcal{C}$, a morphism g of f into f' is a homomorphism

$$g : B \rightarrow B'$$

making the diagram commutative:

We have that $\varphi_S : A \rightarrow S^{-1}A$ is a universal object in this category $\mathcal{C}$.

Theorem 1.4. Let S be a multiplicative subset of A .

1. If S contains no zero divisors, then φ_S is a monomorphism.
2. If A has no zero divisors and $0 \notin S$, then $S^{-1}A$ is an integral domain.
3. If $S \subset A^\times$, then φ_S is an isomorphism.

Definition 1.5. Let A be a commutative ring and S be the set of all nonzero elements of A that are not zero divisors, then $S^{-1}A$ is called the **complete ring of quotients** of the ring A .

The complete ring of quotients of an integral domain A is its **quotient field**, denoted by $\text{Frac}(A)$.

§2 Extensions and Contractions in ring of fractions

Let A be a commutative with identity 1_A and S be a multiplicative subset with $\varphi_S : A \rightarrow S^{-1}A$

Proposition 2.1. 1. If $\mathfrak{a}$ is an ideal in A , then $S^{-1}\mathfrak{a} = \{a/s \mid a \in \mathfrak{a}; s \in S\} = \mathfrak{a}S^{-1}A = \mathfrak{a}^e$.

2. If $\mathfrak{b}$ is an ideal in $S^{-1}A$, then $\varphi_S^{-1}(\mathfrak{b})$ coincides with $\mathfrak{b}^c$

3. let $\mathfrak{a}$ be an ideal of A , then $S^{-1}\mathfrak{a} = S^{-1}A$ if and only if $S \cap \mathfrak{a} \neq \emptyset$.

Corollary 2.2.

$$S^{-1}(\mathfrak{a} + \mathfrak{b}) = S^{-1}\mathfrak{a} + S^{-1}\mathfrak{b}$$

$$S^{-1}(\mathfrak{a}\mathfrak{b}) = (S^{-1}\mathfrak{a})(S^{-1}\mathfrak{b})$$

$$S^{-1}(\mathfrak{a} \cap \mathfrak{b}) = S^{-1}\mathfrak{a} \cap S^{-1}\mathfrak{b}$$

$$S^{-1}\text{Rad}(\mathfrak{a}) = \text{Rad}(S^{-1}\mathfrak{a})$$

Theorem 2.3. .

1. $\mathfrak{b}^{ce} = \mathfrak{b}$ for all ideals $\mathfrak{b}$ of $S^{-1}A$. In other words every ideal in $S^{-1}A$ is of the $S^{-1}\mathfrak{a} = \mathfrak{a}^e$ for some ideal $\mathfrak{a}$ in A by proposition 2.2 .

$$2. \mathfrak{a}^{ec} = \bigcup_{s \in S} (\mathfrak{a} : s)$$

3. If $\mathfrak{p}$ is a prime ideal in A and $S \cap \mathfrak{p} = \emptyset$, then $S^{-1}\mathfrak{p}$ is a prime ideal in $S^{-1}A$

4. there is a one-to-one correspondence between the set $\mathcal{U} = \{\mathfrak{p} : \mathfrak{p} \text{ is prime and disjoint from } S\}$ and the set $\mathcal{V} = \{S^{-1}\mathfrak{p} : S^{-1}\mathfrak{p} \text{ is prime in } S^{-1}R\}$, given by $\mathfrak{p} \mapsto S^{-1}\mathfrak{p}$.

Proof. Let $I = \varphi_S^{-1}(J)$, then $I^e = J^{ce} \subset J$, whence $S^{-1}I \subset J$. Conversely, if $r/s \in J$, then $\varphi_S(r) = rs/s = (r/s)(s^2/s) \in J$, whence $r \in \varphi_S^{-1}(J) = I$. Thus $r/s \in S^{-1}I$ and hence $J \subset S^{-1}I$. $\square$

§3 Fractions of modules

Let A be a commutative ring with identity, M be a A -module and S be a multiplicative subset of A .

Definition 3.1. The *module of fractions* of M with respect to S is the set

$$S^{-1}M = \{m/s \mid m \in M, s \in S\}$$

of equivalence classes of the relation on $M \times S$ defined by

$$(m, s) \sim (m', s') \Leftrightarrow s_1 (s'm - sm') = 0 \text{ for some } s_1 \in S$$

with addition and scalar multiplication defined by

$$m/s + m'/s' = (s'm + sm')/ss' \quad \text{and} \quad (a/t)(m/s) = (am)/ts$$

for all $m, m' \in M, s, s', t \in S$ and $a \in A$.

Lemma 3.2. 1. $S^{-1}(N + P) = S^{-1}(N) + S^{-1}(P)$

2. $S^{-1}(N \cap P) = S^{-1}(N) \cap S^{-1}(P)$

Proposition 3.3. .

1. The functor $S^{-1} : {}_A\mathbf{Mod} \rightarrow {}_{S^{-1}A}\mathbf{Mod}$ is exact

2. $S^{-1}\square$ is additive

3. $S^{-1}\square \cong S^{-1}A \otimes_A \square$ in $[{}_A\mathbf{Mod}, {}_{S^{-1}A}\mathbf{Mod}]$

Corollary 3.4. .

1. $S^{-1}(M/N) \cong (S^{-1}M) / (S^{-1}N)$ in ${}_{S^{-1}A}\mathbf{Mod}$

2. $S^{-1}M \otimes_{S^{-1}A} S^{-1}N \cong S^{-1}(M \otimes_A N)$ in ${}_{S^{-1}A}\mathbf{Mod}$

3. $S^{-1}A$ is a flat A -module

§4 Localization and Local rings

Definition 4.1. Let A be a commutative ring with identity, $\mathfrak{p}$ a prime ideal of A and multiplicative subset $S = A - \mathfrak{p}$. The ring of quotients $S^{-1}A$ is called the **localization of A at $\mathfrak{p}$** and is denoted $A_{\mathfrak{p}}$. If $\mathfrak{a}$ is an ideal in A , then the ideal $\mathfrak{a}^e = S^{-1}\mathfrak{a}$ in $A_{\mathfrak{p}}$.

Remark. We always identify A with its image $\varphi_S(A)$ in $A_{\mathfrak{p}}$ thus A can be considered as a subring of $A_{\mathfrak{p}}$. In this case, the extension ideal $\mathfrak{a}^e = S^{-1}\mathfrak{a} = \mathfrak{a}A_{\mathfrak{p}}$.

Theorem 4.2. Let $\mathfrak{p}$ be a prime ideal in a commutative ring A with identity and localization $A_{\mathfrak{p}}$.

1. There is a one-to-one correspondence between the set $\{\mathfrak{q} : \mathfrak{q} \text{ is prime and contained in } \mathfrak{p}\}$ and the set $\{S^{-1}\mathfrak{q} : S^{-1}\mathfrak{q} \text{ is prime in } A_{\mathfrak{p}}\}$, given by $\mathfrak{q} \mapsto S^{-1}\mathfrak{q}$;
2. The ideal $S^{-1}\mathfrak{p}$ is the unique maximal ideal of $A_{\mathfrak{p}}$.

Definition 4.3. A **local ring** is a commutative ring with identity which has a unique maximal ideal.

Theorem 4.4. The following conditions are equivalent.

1. R is a local ring;
2. all nonunits of R are contained in some ideal $M \neq R$;
3. the nonunits of R form an ideal.
4. for all $r, s \in R$, $r + s = 1_R$ implies r or s is a unit.

Proposition 4.5. Every nonzero homomorphic image of a local ring is local.

§5 Local properties

Definition 5.1. A property P of a ring A (or of an A -module M) is said to be a **local property** if the following is true: A (or M) has $P \Leftrightarrow A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has P , for each prime ideal $\mathfrak{p}$ of A .

Proposition 5.2. The following are local properties:

1. A -module M is zero
2. A -module homomorphism $\phi : M \rightarrow N$ is injective [resp. surjective, bijective]
3. A -module M is flat

Chapter IV

Chain Condition

§1

Definition 1.1. Let A be a ring.

1. A left A -module M is said to satisfy the **ascending chain condition (ACC) on submodules** (or to be **Noetherian**) if for every chain $M_1 \subset M_2 \subset M_3 \subset \cdots$ of submodules of M , there is an integer n such that $M_i = M_n$ for all $i \geq n$.
2. The ring A is **left Noetherian** if left regular module ${}_A A$ is Noetherian. A is said to be **Noetherian** if A is both left and right Noetherian.
3. A left A -module N is said to satisfy, the **descending chain condition (DCC) on submodules** (or to be **Artinian**) if for every chain $N_1 \supset N_2 \supset N_3 \supset \cdots$ of submodules of N , there is an integer m such that $N_i = N_m$ for all $i \geq m$.
4. A is **left Artinian** if ${}_A A$ is Artinian.

A is said to be **Artinian** if A is both left and right Artinian.

Definition 1.2. Let A be a ring, a left A -module M is said to satisfy the **maximum condition** [resp. **minimum condition**] on submodules if every nonempty set of submodules of M contains a maximal [resp. minimal] element (with respect to set theoretic inclusion).

Theorem 1.3. A module satisfies the ascending [resp. descending] chain condition on submodules if and only if A satisfies the maximal [resp. minimal] condition on submodules.

Theorem 1.4. Let A be a ring and $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence of left A -modules. Then ${}_A M$ is Noetherian [resp. Artinian] if and only if ${}_A M'$ and ${}_A M''$ are both Noetherian [resp. Artinian].

Corollary 1.5 (Stability of chain conditions). Let A be a ring, we have

1. Let M be an A -module and $M_1 \subseteq M$ a submodule. Then M is Noetherian [resp. Artinian] if and only if both M_1 and M/M_1 are Noetherian [resp. Artinian].

2. For A -modules $M_1, \dots, M_n$, the direct sum

$$M_1 \oplus \cdots \oplus M_n$$

is Noetherian [resp. Artinian] if and only if each M_i is.

3. Any homomorphic image of a Noetherian [resp. Artinian] module is Noetherian [resp. Artinian].
4. If A is a left Noetherian [resp. Artinian] ring, then every finitely generated left A -module is Noetherian [resp. Artinian].

§2 Normal series and Composition Series of Modules

Definition 2.1. Let A be a ring and a left A -module M .

1. A **normal series** for M is a chain of submodules: $M = M_0 \supset M_1 \supset M_2 \supset \cdots \supset M_n$. The factors of the series are the quotient modules

$$M_i/M_{i+1} \quad (i = 0, 1, \dots, n-1).$$

The **length** of the normal series is the number of proper inclusions (= number of nontrivial factors).

2. A **refinement** of the normal series $M = M_0 \supset M_1 \supset \cdots \supset M_n$ is a normal series obtained by inserting a finite number of additional submodules between the given ones. A **proper refinement** is one which has length larger than the original series.
3. Two normal series are **equivalent** if there is a one-to-one correspondence between the nontrivial factors such that corresponding factors are isomorphic modules. Thus equivalent series necessarily have the same length.
4. A **composition series** for A is a normal series $A = A_0 \supset A_1 \supset A_2 \supset \cdots \supset A_n = 0$ such that each factor A_k/A_{k+1} ($k = 0, 1, \dots, n-1$) is a module with no proper nonempty submodules.

Theorem 2.2. Any two normal series of a module A have refinements that are equivalent. Any two composition series of A are equivalent.

Theorem 2.3. Let A be a ring. A left A -module M has a composition series if and only if M satisfies both the ACC and DCC on submodules.

Proof. ($\Rightarrow$) Suppose A has a composition series S of length n . If either chain condition fails to hold, one, can find submodules

$$A = A_0 \supsetneq A_1 \supsetneq A_2 \supsetneq \cdots \supsetneq A_n \supsetneq A_{n+1}$$

which form a normal series T of length $n + 1$. By 2.2, S and T have refinements that are equivalent. This is a contradiction since equivalent series have equal length. For every refinement of the composition series S has the same length n as S , but every refinement of T necessarily has length at least $n + 1$. Therefore, A satisfies both chain conditions.

($\Leftarrow$) If B is a nonzero submodule of A , let $S(B)$ be the set of all proper submodules C of B . Also define $S(0) = \{0\}$. For each B there is a maximal element B' of $S(B)$ by 1.3. Let S be the set of all submodules of A and define a map $f : S \rightarrow S$ by $f(B) = B'$

Let $A_i = f^{(i)}(A)$, then $A \supset A_1 \supset A_2 \supset \cdots$ is a descending chain by construction, whence for some n , $A_i = A_n$ for all $i \geq n$. Since $A_{n+1} = f(A_n)$, the definition of f shows that $A_{n+1} = A_n$ only if $A_n = 0 = A_{n+1}$. Let m be the smallest integer such that $A_m = 0$. Then $m \leq n$ and $A_k \neq 0$ for all $k < m$. Furthermore for each $k < m$, A_{k+1} is a maximal submodule of A_k such that $A_k \supsetneq A_{k+1}$. Consequently, each A_k/A_{k+1} is nonzero and has no proper submodules by ??. Therefore, $A \supset A_1 \supset \cdots \supset A_m = 0$ is a composition series for A . $\square$

Chapter V

Noetherian Rings and Modules

Contents

§1 Basic Criterion	12
§1.1 General case	12
§1.2 Commutative case	12
§2 Primary Decomposition	14

§1 Basic Criterion

§1.1 General case

theorem 1.3 corollary 1.5

Proposition 1.1. *A left A -module M is Noetherian if and only if every submodule of M is finitely generated.*

§1.2 Commutative case

Proposition 1.2. *Let A be a commutative Noetherian and let S be a multiplicative subset of A . Then the ring of fractions $S^{-1}A$ is Noetherian.*

Theorem 1.3. *Let A be a left Noetherian ring, then $A[x_1, \dots, x_n]$ is also left Noetherian.*

Proof. We prove the case $n = 1$; the general case follows by induction on n .

Let $\mathfrak{a}$ be a left ideal of $A[x]$. For each nonnegative integer d , let $\mathfrak{a}_d$ be the set of all leading coefficients of polynomials in $\mathfrak{a}$ of degree d together with 0. Then $\mathfrak{a}_d$ is a left ideal of A for each d and $\mathfrak{a}_0 \subset \mathfrak{a}_1 \subset \mathfrak{a}_2 \subset \dots$ is an ascending chain of left ideals of A . Since A is left Noetherian, there exists an integer n such that $\mathfrak{a}_d = \mathfrak{a}_n$ for all $d \geq n$.

For each $d = 0, 1, \dots, n$, let $a_{d1}, a_{d2}, \dots, a_{dm_d}$ be generators of the left ideal $\mathfrak{a}_d$. For each a_{di} , choose a polynomial $f_{di}(x) \in \mathfrak{a}$ of degree d with leading coefficient a_{di} .

We claim that the finite set of polynomials

$$\{f_{di}(x) \mid d = 0, 1, \dots, n; i = 1, 2, \dots, m_d\}$$

generates the ideal $\mathfrak{a}$.

Let $f(x)$ be any polynomial in $\mathfrak{a}$. If $\deg f(x) = d$, then the leading coefficient of $f(x)$ is in $\mathfrak{a}_d = \mathfrak{a}_n$, so there exist elements $b_i \in A$ such that

$$\text{leading coefficient of } f(x) = \sum_{i=1}^{m_n} b_i a_{ni}$$

Let $g(x) = \sum_{i=1}^{m_n} b_i x^{d-n} f_{ni}(x)$. Then $g(x) \in \mathfrak{a}$ and $\deg g(x) = d$ with leading coefficient equal to that of $f(x)$. Thus $\deg(f(x) - g(x)) < d$ and by repeating this process, we can express $f(x)$ as an $A[x]$ -linear combination of the polynomials $f_{di}(x)$. Therefore, $\mathfrak{a}$ is finitely generated and $A[x]$ is left Noetherian. $\square$

Corollary 1.4. *The finitely generated algebra over a left Noetherian ring is left Noetherian.*

Proposition 1.5. *Let $A \subseteq B \subseteq C$ be rings. Suppose that A is Noetherian, that C is finitely generated as an A -algebra and that C is either (i) finitely generated as a B -module or (ii) integral over B . Then B is finitely generated as an A -algebra.*

Proposition 1.6. *Let A be a subring of B ; suppose that A is Noetherian and that B is finitely generated as an A -module. Then B is a Noetherian ring.*

Proposition 1.7. *If A is Noetherian and S is any multiplicatively closed subset of A , then $S^{-1}A$ is Noetherian.*

Theorem 1.8 (Hilbert's Basis Theorem). *If A is a commutative Noetherian ring with identity, then so is $R[x_1, \dots, x_n]$ and $R[[x]]$*

Proposition 1.9. *If R is a commutative ring with identity and $\mathfrak{p}$ is an ideal which is maximal in the set of all ideals of R which are not finitely generated, then $\mathfrak{p}$ is prime.*

Proof. Suppose $ab \in \mathfrak{p}$ but $a \notin \mathfrak{p}$ and $b \notin \mathfrak{p}$. Then $\mathfrak{p} + (a)$ and $\mathfrak{p} + (b)$ are ideals properly containing $\mathfrak{p}$ and therefore finitely generated by maximality of $\mathfrak{p}$. Consequently $\mathfrak{p} + (a) = (p_1 + r_1 a, \dots, p_n + r_n a)$ and $\mathfrak{p} + (b) = (p'_1 + r'_1 b, \dots, p'_m + r'_m b)$ for some $p_i, p'_i \in \mathfrak{p}$ and $r_i, r'_i \in R$.

If $J = (\mathfrak{p} : a) = \{r \in R \mid ra \in \mathfrak{p}\}$, then J is an ideal. Since $ab \in \mathfrak{p}$, $(p'_i + r'_i b)a = p'_i a + r'_i ab \in \mathfrak{p}$ for all i , whence $\mathfrak{p} \subset \mathfrak{p} + (b) \subset J$. By maximality, J is finitely generated, say $J = (j_1, \dots, j_k)$.

If $x \in \mathfrak{p}$, then $x \in \mathfrak{p} + (a)$ and hence for some $s_i \in R$, $x = \sum_{i=1}^n s_i (p_i + r_i a) = \sum_{i=1}^n s_i p_i + \sum_{i=1}^n s_i r_i a$. Consequently, $(\sum_i s_i r_i) a = x - \sum_i s_i p_i \in \mathfrak{p}$, whence $\sum_i s_i r_i \in J$. Thus for some $t_i \in R$, $\sum_{i=1}^n s_i r_i = \sum_{i=1}^k t_i j_i$ and $x = \sum_{i=1}^n s_i p_i + \sum_{i=1}^k t_i j_i a$. Therefore, $\mathfrak{p}$ is generated by $p_1, \dots, p_n, j_1 a, \dots, j_k a$, which is a contradiction. Thus $a \in \mathfrak{p}$ or $b \in \mathfrak{p}$ and $\mathfrak{p}$ is prime by ?? $\square$

Theorem 1.10 (I.S.Cohen). *A commutative ring R with identity is Noetherian if and only if every prime ideal of R is finitely generated.*

Proof. Let $\mathcal{S}$ be the set of all ideals of R which are not finitely generated. If $\mathcal{S}$ is nonempty, then use Zorn's Lemma to find a maximal element P of $\mathcal{S}$. P is prime by Proposition 2.4 and hence finitely generated by hypothesis.

This is a contradiction unless $\mathcal{S} = \emptyset$. Therefore, R is Noetherian by Theorem 1.9. $\square$

§2 Primary Decomposition

Throughout this section, A be a commutative ring, and that module [resp. homomorphism] are A -modules [resp. A -module homomorphisms] unless otherwise stated.

Definition 2.1. *Let M be an A -module.*

1. *A submodule Q of M is said to be **primary** if $Q \neq M$, and if principal homomorphism $a_{M/Q}$ is injective or nilpotent for each $a \in A$.*

*A ideal $\mathfrak{q}$ in A is said to be **primary** if it is a primary submodule of the regular module ${}_A A$.*

Remark. *Ideal $\mathfrak{q}$ is primary in A iff $xy \in \mathfrak{q}, x \notin \mathfrak{q} \Rightarrow y^n \in \mathfrak{q}$ for some $n > 0$*

2. *Let Q be a primary submodule of M . The **associated prime ideal** of Q is the prime ideal $\mathfrak{p} = \text{Rad}(\text{Ann}(M/Q))$. We shall say that Q is **primary for $\mathfrak{p}$** or that Q is **$\mathfrak{p}$ -primary**.*

Proposition 2.2. *If $Q_1, \dots, Q_n$ are $\mathfrak{p}$ -primary submodules of M , then so is their intersection $Q_1 \cap Q_2 \cap \dots \cap Q_n$.*

Theorem 2.3. *Let A be a commutative ring, $\mathfrak{q}$ and $\mathfrak{p}$ be ideals in A . Then $\mathfrak{q}$ is primary for $\mathfrak{p}$ if and only if*

$$(i) \quad \mathfrak{q} \subset \mathfrak{p} \subset \text{Rad}(\mathfrak{q})$$

$$(ii) \quad \text{if } ab \in \mathfrak{q} \text{ and } a \notin \mathfrak{q}, \text{ then } b \in \mathfrak{p}.$$

Proof. Suppose (i) and (ii) hold. If $ab \in \mathfrak{q}$ with $a \notin \mathfrak{q}$, then $b \in \mathfrak{p} \subset \text{Rad } \mathfrak{q}$, whence $b^n \in \mathfrak{q}$ for some $n > 0$. Therefore $\mathfrak{q}$ is primary.

To show that $\mathfrak{q}$ is primary for $\mathfrak{p}$ we need only show $\mathfrak{p} = \text{Rad } \mathfrak{q}$. By (i), $\mathfrak{p} \subset \text{Rad } \mathfrak{q}$. If $b \in \text{Rad } \mathfrak{q}$, let n be the least integer such that $b^n \in \mathfrak{q}$. If $n = 1$, $b \in \mathfrak{q} \subset \mathfrak{p}$. If $n > 1$, then $b^{n-1}b = b^n \in \mathfrak{q}$, with $b^{n-1} \notin \mathfrak{q}$ by the minimality of n . By (ii), $b \in \mathfrak{p}$. Thus $b \in \text{Rad } \mathfrak{q}$ implies $b \in \mathfrak{p}$, whence $\text{Rad } \mathfrak{q} \subset \mathfrak{p}$. $\square$

Proposition 2.4. *Clearly every prime ideal is primary. Also the contraction of a primary ideal is primary, for if $f : A \rightarrow B$ and if $\mathfrak{q}$ is a primary ideal in B , then $A/\mathfrak{q}^c$ is isomorphic to a subring of $B/\mathfrak{q}$.*

Proposition 2.5. *If $\text{Rad}(\mathfrak{a})$ is maximal, then $\mathfrak{a}$ is primary. In particular, the powers of a maximal ideal $\mathfrak{m}$ are $\mathfrak{m}$ -primary.*

Proof. Let $\text{Rad}(\mathfrak{a}) = \mathfrak{m}$. The image of $\mathfrak{m}$ in $A/\mathfrak{a}$ is the nilradical of $A/\mathfrak{a}$, hence $A/\mathfrak{a}$ has only one prime ideal $\pi(\mathfrak{m})$, by (1.8). Hence every element of $A/\mathfrak{a}$ is either a unit or nilpotent, and so every zero-divisor in $A/\mathfrak{a}$ is nilpotent. $\square$

Definition 2.6. *Let N be a submodule of an A -module M .*

1. *When N is written as a finite intersection of primary submodules of M ,*

$$N = Q_1 \cap Q_2 \cap \cdots \cap Q_n$$

*we shall say that N has a **primary decomposition** in M .*

2. *If each associated prime ideal $\mathfrak{p}_i$ of Q_i is distinct, then the primary decomposition is said to be **reduced**.*
3. *If $\mathfrak{p}_i$ does not contain any $\mathfrak{p}_j$ for $j \neq i$, then $\mathfrak{p}_i$ is called an **isolated prime** of the primary decomposition.*

Remark. *Using 2.3, we can always obtain from a given primary decomposition of N another reduced primary decomposition.*

Theorem 2.7 (Uniqueness of reduced primary decomposition). *Let N be a submodule of M with two reduced primary decompositions,*

$$Q_1 \cap Q_2 \cap \cdots \cap Q_k = N = Q'_1 \cap Q'_2 \cap \cdots \cap Q'_s$$

where Q_i is $\mathfrak{p}_i$ -primary and Q'_j is $\mathfrak{p}'_j$ -primary. Then $k = s$ and (after reordering if necessary) $\mathfrak{p}_i = \mathfrak{p}'_i$ for all $i = 1, 2, \dots, k$. Furthermore if Q_i and Q'_i both are $\mathfrak{p}_i$ -primary and $\mathfrak{p}_i$ is an isolated prime, then $Q_i = Q'_i$.

Theorem 2.8 (Existence of reduced primary decomposition). *Let M be a Noetherian A -module and N a submodule of M . Then N admits a reduced primary decomposition.*

Proof. Let $\mathcal{S}$ be the set of all submodules of M that do not have a primary decomposition. Clearly no primary submodule is in $\mathcal{S}$. We must show that $\mathcal{S}$ is actually empty. If $\mathcal{S}$ is nonempty, then $\mathcal{S}$ contains a maximal element C by Theorem 1.4.

Since C is not primary, there exist $r \in R$ and $b \in M - C$ such that $rb \in C$ but $r^n M \not\subset C$ for all $n > 0$. Let $M_n = (C : r^n) = \{x \in M \mid r^n x \in C\}$. Then each M_n is a submodule of M and $M_1 \subset M_2 \subset M_3 \subset \cdots$. By hypothesis there exists $k > 0$ such that $M_i = M_k$ for $i \geq k$. Let D be the submodule $r^k M + C = \{x \in M \mid x = r^k y + c \text{ for some } y \in M, c \in C\}$. Clearly $C \subset M_k \cap D$.

Conversely, if $x \in M_k \cap D$, then $x = r^k y + c$ and $r^k x \in C$, whence $r^{2k} v = r^k (r^k y) = r^k (x - c) = r^k x - r^k c \in C$. Therefore, $y \in M_{2k} = M_k$. Consequently, $r^k y \in C$ and hence $x = r^k y + c \in C$. Therefore $M_k \cap D \subset C$, whence $M_k \cap D = C$. Now $C \neq M_k \neq M$ and $C \neq D \neq M$ since $b \in M_k - C$ and $r^k M \not\subset C$. By the maximality of C in $\mathcal{S}$, M_k and D must have primary decompositions. Thus C has a primary decomposition, which is a contradiction. Therefore $\mathcal{S}$ is empty and every submodule has a primary decomposition. Consequently, every submodule has a reduced primary decomposition by ??.

□

Corollary 2.9. *If R is a commutative Noetherian ring with identity and M is a finitely generated unitary R -module. Then every submodule $N (\neq M)$ of M has a reduced primary decomposition.*

Proof. This is an immediate consequence of ?? ?? and 2.8

□

Chapter VI

Artinian

Contents

§1	Artinian Rings	17
§2	Commutative Artinian Rings	18

§1 Artinian Rings

Lemma 1.1. *If A is a left Artinian ring, then $J(A)$ is nilpotent.*

Proof. Consider the descending chain of ideals

$$J \supset J^2 \supset J^3 \supset \dots$$

there exists an integer n such that $J^n = J^{n+1} = \dots$. Denote J^n by I , then $I = JI$. Assume $I \neq 0$ and let nonempty set $I^2 = I \neq 0$

$$\mathcal{S} := \{\text{left ideal } L : IL \neq 0\}$$

Then there exists a minimal element L_0 in $\mathcal{S}$ by 1.3. Furthermore, $L_0 = Aa$ for some $a \in L_0$ and $IL_0 = L_0$ by minimality. Hence, there exists a nonzero $x \in I \subset J^n$ such that $xa = a$, that is, $1_A - x$ is a left zero divisor which contradicts the definition of J . Therefore, $J^n = I = 0$. $\square$

Lemma 1.2. *Let A is simple left Artinian ring, then the following condition are equivalent for a left A -module M :*

1. Noetherian
2. Artinian
3. has finite length.

Theorem 1.3 (Hopkins-Levitzki). *Let A be a left Artinian ring and a left A -modules M . If M is Artinian, then M is Noetherian.*

Proof. Consider the descending chain of submodules

$$M \supset JM \supset J^2M \supset \cdots \supset J^nM = 0$$

where $J = J(A)$. Then J^kM and $J^kM/J^{k+1}M$ are Artinian for all k by 1.5.

On the other hand, $J^kM/J^{k+1}M$ is a module over the semisimple Artinian ring A/J , and is also Artinian. Thus each $J^kM/J^{k+1}M$ is Noetherian and thus M is Noetherian by 1.5. $\square$

Example 1.4. Let ring $A = \begin{pmatrix} \mathbb{Q} & \mathbb{R} \\ 0 & \mathbb{R} \end{pmatrix}$, A is right Artinian but not left Artinian. infinite descending chain of left ideals

$$\begin{pmatrix} 0 & V_n \\ 0 & 0 \end{pmatrix} \supsetneq \begin{pmatrix} 0 & V_{n+1} \\ 0 & 0 \end{pmatrix} \supsetneq \cdots$$

where $\{V_n\}$ is a infinite descending chain of $\mathbb{Q}$ -subspace of $\mathbb{R}$.

In generally, the ring of upper triangular matrices

$$T := \begin{pmatrix} D_1 & M \\ 0 & D_2 \end{pmatrix}$$

where D_1, D_2 are rings and M is a (D_1, D_2) -bimodule. Then T is left Artinian if and only if D_1, D_2 are left Artinian and M_{D_2} has finite length.

§2 Commutative Artinian Rings

In this subsection we shall study commutative Artinian rings.

Theorem 2.1. Let A be a commutative ring with identity. Then the following conditions are equivalent:

1. A is Artinian
2. A is Noetherian and $\dim A = 0$.
3. $\text{Spec}(A)$ is finite and $\text{Krull dim } A = 0$
4. (Structure theorem) A is isomorphic to a finite direct product of Artin local rings.

Proof. 1. $\Rightarrow$ 2. By 1.3, A is Noetherian. If $\mathfrak{p}$ is a prime ideal in A , then $A/\mathfrak{p}$ is an integral domain and Artinian, whence a field. Thus $\dim A = 0$.

2. $\Rightarrow$ 3. Let $\mathcal{S} := \{\mathfrak{p}_1 \cap \mathfrak{p}_1 \cdots \cap \mathfrak{p}_r : \mathfrak{p}_i \text{ maximal ideal}\}$, then $\mathcal{S}$ has a minimal element $\mathfrak{a} = \mathfrak{p}_1 \cap \mathfrak{p}_1 \cdots \cap \mathfrak{p}_n$ by 1.3. Hence for any maximal ideal $\mathfrak{p}$, $\mathfrak{p} \supset \mathfrak{a}$ by minimality of $\mathfrak{a}$, whence $\mathfrak{m} = \mathfrak{m}_i$ for some i by . $\square$

Corollary 2.2. In a commutative Artin ring the

1. *nilradical is equal to the Jacobson radical.*
2. *A has finitely many maximal ideals*

Proposition 2.3. *Let A be an Artin local ring. Then the following are equivalent:*

1. *every ideal in A is principal;*
2. *the maximal ideal $\mathfrak{m}$ is principal;*
3. $\dim_k (\mathfrak{m}/\mathfrak{m}^2) \leq 1$.

Chapter VII

The Structure of Rings and Modules

Contents

§1	Simplicity	20
§1.1	Primitivity	21
§1.2	Jacobson Density Theorem	21
§1.3	Artinian simple ring	23
§2	Semisimplicity	24
§2.1	Structure of semisimple rings	24
§2.2	Structure of modules over semisimple rings	26
§3	Jacobson Radical	26
§3.1	Jacobson radical	26
§3.2	Nakayama's Lemma	28
§4	Characterizations of semisimple rings	28
§5	Algebra	28

§1 Simplicity

Definition 1.1. A ring A is said to be **simple** if A has no proper two-sided ideals.

Definition 1.2. A left module M over a ring A is said to be **simple** (or **irreducible**) if M has no proper submodules.

Remark. A left ideal $\mathfrak{a}$ of A is a simple left A -module if and only if $\mathfrak{a}$ is a minimal left ideal of A . In this case, we call $\mathfrak{a}$ the **simple left ideal** of A .

Proposition 1.3. Let A be a ring and M be a simple left A -module, then

1. $M = Am$ for every $0 \neq m \in M$.

2. If $0 \neq u \in M$, then $M \cong A/\text{Ann}(u)$, thus $\text{Ann}(u)$ is a left maximal ideal.

Conversely, if $\mathfrak{m}$ is left maximal in A , then $A/\mathfrak{m}$ is a simple A -module with $\text{Ann}(A/\mathfrak{m}) = \mathfrak{m}$.

3. If A is not a division ring, then M is a torsion module. $m \in M$.

Lemma 1.4 (Schur). Let M be a simple left module over a ring A and let N_i be any A -module.

1. Every nonzero A -module homomorphism $f : M \rightarrow N_1$ is a monomorphism;
2. Every nonzero A -module homomorphism $g : N_2 \rightarrow M$ is an epimorphism;
3. The endomorphism ring $D = \text{Hom}_A(M, M)$ is a division ring, then M is a vector space over $\text{Hom}_A(M, M)$ with $fa = f(a)$

§1.1 Primitivity

Definition 1.5. Let A be a ring.

1. The ring A is said to be **left primitive** if there exists a simple faithful left A -module.
2. An two-sided ideal $\mathfrak{a}$ of a ring A is said to be **left primitive** if the quotient ring $A/\mathfrak{a}$ is a left primitive ring.

Remark. If M is a simple left A -module, then $A/\text{Ann}(M)$ is left primitive with faithful simple left $A/\text{Ann}(M)$ -module M .

Proposition 1.6. Let A be a ring.

1. A simple ring A is left primitive with faithful simple left A -module $A/\mathfrak{m}$ for any left maximal ideal $\mathfrak{m}$ of A .
2. A commutative ring A is primitive if and only if A is a field.

§1.2 Jacobson Density Theorem

Definition 1.7. Let V be a vector space over a division ring D . A subring A of $\text{End}_D(V)$ is called a **dense ring of endomorphisms** of V if for every positive integer n , every linearly independent subset $\{u_1, \dots, u_n\}$ of V and every arbitrary subset $\{v_1, \dots, v_n\}$ of V , there exists $f \in A$ such that $f(u_i) = v_i, (i = 1, 2, \dots, n)$.

Lemma 1.8. Let M be a simple module over a ring A . Consider M as a vector space over the division ring $D = \text{End}_A(M)$. If V is a finite dimensional D -subspace of M and $a \in M - V$, then there exists $r \in A$ such that $ra \neq 0$ and $rV = 0$.

Remark. In other words, the element $r \in \text{Ann}_A(V)$ only annihilates D -subspace V .

Proof. The proof is by induction on $n = \dim_D V$. If $n = 0$, then $V = 0$ and $a \neq 0$. Since M is simple, $M = Ra$. Consequently, there exists $r \in R$ such that $ra = a \neq 0$ and $rV = r0 = 0$.

Suppose now $\dim_D V = n > 0$ and the theorem is true for dimensions less than n . Let $\{u_1, \dots, u_{n-1}, u\}$ be a D -basis of V and let $W = \text{span}\{u_1, \dots, u_{n-1}\}$ ($W = 0$ if $n = 1$). Then $V = W \oplus Du$ (vector space direct sum, W may not be an R -submodule of M) the left annihilator $I = \text{Ann}_R(W)$ is a left ideal of R .

Consequently, Iu is an R -submodule of M . Since $u \in M - W$, the induction hypothesis implies that there exists $r \in R$ such that $ru \neq 0$ and $rW = 0$. Consequently $r \in I$ and $0 \neq ru \in Iu$, whence $Iu \neq 0$. Therefore $M = Iu$ by simplicity.

We must find $r \in R$ such that $ra \neq 0$ and $rV = 0$. If no such r exists, $\text{Ann}(a) \subset \text{Ann}(V)$, then we can define a map $\theta : M \rightarrow M$ as follows. For $ru \in Iu = M$ let $\theta(ru) = ra \in M$. We claim that θ is well defined. If $r_1u = r_2u$ ($r_i \in I$), then $(r_1 - r_2)u = 0$, whence $(r_1 - r_2)V = (r_1 - r_2)(W \oplus Du) = 0$. Consequently by hypothesis $(r_1 - r_2)a = 0$. Therefore, $\theta(r_1u) = r_1a = r_2a = \theta(r_2u)$. Verify that $\theta \in \text{Hom}_R(M, M) = D$. Then for every $r \in I$,

$$0 = \theta(ru) - ra = r\theta(u) - ra = r(\theta(u) - a)$$

Therefore $\theta(u) - a \in W$ by induction hypothesis. Consequently

$$a = \theta u - (\theta u - a) \in Du + W = V,$$

which contradicts the fact that $a \notin V$. Therefore, there exists $r \in R$ such that $ra \neq 0$ and $rV = 0$. $\square$

Theorem 1.9 (Classic Jacobson Density Theorem). *Let A be a left primitive ring with a faithful simple left A -module M and division ring $D = \text{End}_A(M)$. Then A is a dense ring of endomorphisms of the D -vector space M (viewed $\alpha : A \hookrightarrow \text{End}_D(M)$ by $a \mapsto \alpha_a$ where left regular action $\alpha_a : m \mapsto am$ in M).*

Proof. It clear that $\alpha : A \rightarrow \text{End}_D(M)$ is a ring monomorphism since M is faithful. Let $\{u_1, u_2, \dots, u_n\}$ be a D -linearly independent subset and $\{v_1, v_2, \dots, v_n\}$ be an arbitrary subset. For each i let

$$V_i = \text{span}\{u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_n\}.$$

Since U is D -linearly independent, $u_i \notin V_i$. Consequently, by 1.8 there exists $r_i \in R$ such that

$$r_i u_i \neq 0 \text{ and } r_i V_i = 0$$

whence $Rr_i u_i = M$ by simplicity. Therefore exists $t_i \in R$ such that $t_i r_i u_i = v_i$. Let

$$r = t_1 r_1 + t_2 r_2 + \dots + t_n r_n \in R.$$

Consequently for each $i = 1, 2, \dots, n$

$$\alpha_r(u_i) = (t_1 r_1 + \dots + t_n r_n) u_i = v_i$$

Therefore $\text{Im } \alpha$ is a dense ring of endomorphisms of the D -vector space M . □

Corollary 1.10. *If M is a simple left A -module, then A acts densely on M . That is, for every positive integer n , every linearly independent subset $\{u_1, \dots, u_n\}$ of M and every arbitrary subset $\{v_1, \dots, v_n\}$ of M , there exists $a \in A$ such that $au_i = v_i$, ($i = 1, 2, \dots, n$).*

§1.3 Artinian simple ring

Lemma 1.11. *Let A be a dense ring of endomorphisms of a vector space V over a division ring D . Then A is left Artinian if and only if $\dim_D V$ is finite, in which case $A = \text{End}_D(V) \cong M_n(D)$.*

Proof. If A is Artinian and $\dim_D V$ is infinite, then there exists an infinite linearly independent subset $\{u_1, u_2, \dots\}$ of V . The V is a left simple $\text{End}_D(V)$ -module and hence a left simple A -module.

For each n , let $I_n = \text{Ann}_A \{u_1, \dots, u_n\}$. Then $I_1 \supsetneq I_2 \supsetneq \dots$ is a properly descending chain by lemma 1.8, which is a contradiction. Hence $\dim_D V$ is finite.

Conversely if $\dim_D V$ is finite, then V has a finite basis $\{v_1, \dots, v_m\}$. Then $A = \text{Hom}_D(V, V) \cong M_n(D)$ is Artinian. □

Corollary 1.12 (Artin-Wedderburn). *If A is a simple ring, then*

1. A is left Artinian
2. $A \cong M_n(D)$ for some positive integer n and some division ring $D \cong \text{End}_A(A/\mathfrak{m})$ for some maximal left ideal $\mathfrak{m}$ of A .
3. A is right Artinian

Lemma 1.13 (Schur's lemma). *Let V be a nonzero vector space over a division ring D . If $g : V \rightarrow V$ is a homomorphism of additive groups such that $gf = fg$ for all $f \in \text{Hom}_D(V, V)$, then there exists $\lambda \in D$ such that $g(x) = \lambda x$ for all $x \in V$.*

Remark. *That is, that is,*

$$g \in \text{End}_{\text{End}_D(V)}(V) \cong D$$

since V is a simple $\text{End}_D(V)$ -module.

Lemma 1.14. *The simple left $M_n(D)$ -modules for some division ring D is isomorphic to D^n with the usual left matrix multiplication.*

Proposition 1.15. *If $M_{n_1}(D_1) \cong M_{n_2}(D_2)$ for some positive integers n_1, n_2 and some division rings D_1, D_2 , then $n_1 = n_2$ and $D_1 \cong D_2$.*

Proof.

$$D_1 \cong \text{End}_{M_{n_1}(D_1)}(D_1^{n_1}) \cong \text{End}_{M_{n_2}(D_2)}(D_2^{n_2}) \cong D_2$$

□

§2 Semisimplicity

Theorem 2.1. *Let A be a ring and M a left A -module. The following conditions on M are equivalent:*

1. M is the sum of a family of simple submodules.
2. M is the direct sum of a family of simple submodules.
3. Every submodule N is a direct summand of M .

Proof. (1) $\Rightarrow$ (2) Let $\mathcal{S}$ be the set of all families $\mathcal{F}$ of simple submodules of M such that the sum of the members of $\mathcal{F}$ is direct. Since M is the sum of a family of simple submodules, $\mathcal{S}$ is nonempty. Partially order $\mathcal{S}$ by inclusion and let $\mathcal{C}$ be a chain in $\mathcal{S}$. Then $\mathcal{U} = \bigcup_{\mathcal{F} \in \mathcal{C}} \mathcal{F}$ is an upper bound for $\mathcal{C}$ in $\mathcal{S}$. By Zorn's lemma there exists a maximal element $\mathcal{F}_0$ in $\mathcal{S}$. We claim that $M = \bigoplus_{N \in \mathcal{F}_0} N$. If not, there exists a simple submodule K of M such that

$$K \cap \left(\bigoplus_{N \in \mathcal{F}_0} N \right) = 0.$$

Consequently, $\mathcal{F}_0 \cup \{K\} \in \mathcal{S}$, contradicting the maximality of $\mathcal{F}_0$.

(2) $\Rightarrow$ (3) Let $M = \bigoplus_{i \in I} N_i$, where each N_i is a simple submodule of M , and let N be a submodule of M . For each $i \in I$, either $N_i \subset N$ or $N_i \cap N = 0$ by simplicity. Let $J = \{i \in I \mid N_i \subset N\}$ and $K = I - J$. Then

$$M = N \oplus \left(\bigoplus_{i \in K} N_i \right).$$

(3) $\Rightarrow$ (1) Let N be the sum of all simple submodules of M . By hypothesis, $M = N \oplus P$ for some submodule P . □

Remark. A left A -module M satisfying the three conditions is said to be **semisimple**. Similarly one defines a right semisimple module.

Proposition 2.2. *Every submodule and every factor module of a left semisimple module is left semisimple.*

§2.1 Structure of semisimple rings

Definition 2.3. A ring A is called **semisimple** if $1_A \neq 0$, and if ${}_A A$ is a semisimple module.

Proposition 2.4. *If A is a semisimple ring, then A is left Artinian.*

Proof. Since ${}_A A$ is semisimple module, there exist simple left ideals L_j of A such that

$${}_A A \cong \bigoplus_{j \in J} L_j$$

with projection maps $\pi_j : {}_A A \rightarrow {}_A L_j$ for each $j \in J$. Since the sum is direct, $\{j_k : \pi_{j_k}(1_A) \neq 0\}$ is finite. Then $J = \{j_k : \pi_{j_k}(1_A) \neq 0\}$ is finite and

$${}_A A \cong L_1 \oplus L_2 \oplus \cdots \oplus L_n$$

Therefore ring A is left Artinian. □

Remark. *It is clear that any simple left ideal L of A is isomorphic to some L_j , thus A has finitely many minimal left ideals up to isomorphism.*

Lemma 2.5. *If L and L' are minimal left ideals in a ring R , then each of the following statements implies the one below it:*

1. $LL' \neq (0)$.
2. $\text{Hom}_R(L, L') \neq \{0\}$ and there exists $b' \in L'$ with $L' = Lb'$.
3. $L \cong L'$ as left R -modules.

If also $L^2 \neq (0)$, then (iii) implies (i) and the three statements are equivalent.

Theorem 2.6 (Wedderburn-Artin). *If A is a semisimple ring, then we have ring isomorphisms*

$$A \cong \prod_{i=1}^t A_i \cong \prod_{i=1}^t M_{n_i}(D_i)$$

where $A_i \cong M_{n_i}(D_i)$ $i \in I$.

Proof. Let $\{L_{j_i}\}_{i=1}^t$ be all distinct simple left ideals of A up to isomorphism and define A_i be the sum

$${}_A A_i := \bigoplus_{L_j \cong L_i} L_j$$

Thus, as left A -modules,

$${}_A A \cong \bigoplus_{i=1}^t {}_A A_i.$$

Then A_i is a two-sided ideal of A by lemma 2.5 and hence a ring with identity element $e_i = \pi_i(1_A)$ (π_i the projection from A to A_i). The projection from A to A_i induce a ring homomorphism

$$A \cong \prod_{i=1}^t A_i$$

since $1_A = e_1 + e_2 + \cdots + e_t$ and $e_i e_j = \delta_{ij} 1_A$.

Finally, each A_i is a simple ring since any nonzero two-sided ideal $\mathfrak{a}$ of A_i contains a minimal left ideal L of A_i and hence a minimal left ideal of A isomorphic to L_i . Then $Lb \subset L$ runs over all minimal left ideals of A isomorphic to L_i by lemma 2.5, whence $A_i \subset L \subset \mathfrak{a}$. $\square$

§2.2 Structure of modules over semisimple rings

Theorem 2.7. *Let semisimple ring*

$$A \cong \prod_{i=1}^t A_i$$

where each $A_i \cong M_{n_i}(D_i)$. Then every left A -module M is isomorphic to

$$\bigoplus_{i=1}^t e_i M \cong \bigoplus_{i=1}^t S_i^{(\kappa_i)}$$

where $e_i M \cong S_i \cong D_i^{n_i}$ and κ_i is a cardinal number.

And the decomposition is unique in the sense that if

$$M \cong \bigoplus_{i=1}^t S_i^{(\kappa_i)} \cong \bigoplus_{i=1}^t S_i^{(\kappa'_i)}$$

then $\kappa_i = \kappa'_i$ for each i

Remark. If M is finitely generated, then each κ_i is finite.

§3 Jacobson Radical

Definition 3.1. Let R be a ring. A element $x \in R$ is said to be **right quasi-regular** if there exists $y \in R$ such that $x + y - xy = 0$, y is called a **right quasi-inverse** of x .

Remark. That is, $1 - x$ has a right inverse $1 - y$.

Definition 3.2. A element $a \in R$ is said **right quasi-nilpotent element** if for every $r \in R$, ra is right quasi-regular.

Remark. That is, $1 - ra$ has a right inverse for every $r \in R$.

§3.1 Jacobson radical

Theorem 3.3. Let R be a ring, then the following sets are equal:

1. the intersection of all maximal left ideals of R
- 1' the intersection of all maximal right ideals of R

2. the intersection of all the annihilators of simple left R -modules;
- 2' the intersection of all the annihilators of simple right R -modules;
3. $\{x \in R : x \text{ is right quasi-nilpotent element}\}$
- 3' $\{x \in R : x \text{ is left quasi-nilpotent element}\}$
4. Nakayama's $\{x \in R : \text{for any finitely generated module } M, xM = M \Rightarrow M = 0\}$
5. the ideal J such that R/J is semisimple.

the two-sided ideal is called the **Jacobson radical** of R and is denoted by $J(R)$.

Corollary 3.4. 1. $J(\prod_{i \in I} R_i) = \prod_{i \in I} J(R_i)$.

2. $J(J(R)) = J(R)$.

3. $R/J(R)$ is semisimple.

Proposition 3.5. If R is a Artinian ring, then the radical $J(R)$ is a nilpotent ideal. Thus $J(R) = \text{Rad}(R)$

Proof. Since R is Artinian, the descending chain of ideals

$$J(R) \supseteq J(R)^2 \supseteq J(R)^3 \supseteq \dots$$

stabilizes, say $J(R)^n = J(R)^{n+1}$ for some positive integer n . On the other hand, ${}_R R$ is Noetherian thus $J(R)^n$ is finitely generated submodule of ${}_R R$ and $J(R)^n = 0$ by Nakayama's lemma. $\square$

Proposition 3.6. Let R be a ring. $J(M_n R) = M_n(J(R))$.

Proof. (a) If A is a left R -module, consider the elements of $A^n = A \oplus A \oplus \dots \oplus A$ (n summands) as column vectors; then A^n is a left $(\text{Mat}_n R)$ -module (under ordinary matrix multiplication).

(b) If A is a simple R -module, A^n is a simple $(\text{Mat}_n R)$ -module.

(c) $J(\text{Mat}_n R) \subset \text{Mat}_n J(R)$.

(d) $\text{Mat}_n J(R) \subset J(\text{Mat}_n R)$. [Hint: prove that $\text{Mat}_n J(R)$ is a left quasi-regular ideal of $\text{Mat}_n R$ as follows. For each $k = 1, 2, \dots, n$ let K_k consist of all matrices (a_{ij}) such that $a_{ij} \in J(R)$ and $a_{ij} = 0$ if $j \neq k$. Show that K_k is a left quasi-regular left ideal of $\text{Mat}_n R$ and observe that $K_1 + K_2 + \dots + K_n = \text{Mat}_n J(R)$.] $\square$

Corollary 3.7. Let R be a ring. If matrix $A \in M_n(J(R))$, then $I - A$ is invertible in $M_n(R)$.

§3.2 Nakayama's Lemma

Theorem 3.8 (Nakayama's lemma). *Let M be a finitely generated left R -module and a subset $X \subset J(R)$. Then $\mathfrak{a}M = M$ implies $M = 0$.*

Corollary 3.9. *Let M be a finitely generated left R -module, N a submodule of M , left ideal $\mathfrak{a} \subset J(R)$. Then $M = \mathfrak{a}M + N \Rightarrow M = N$.*

Corollary 3.10. *If R is a Noetherian local ring with maximal ideal $\mathfrak{m}$, then $\bigcap_{n=1}^{\infty} \mathfrak{m}^n = 0$.*

§4 Characterizations of semisimple rings

Theorem 4.1. *The following conditions on a ring R are equivalent:*

1. R is a semisimple ring.
2. Every left R -module is a semisimple module.
3. Every left R -module is injective.
4. Every left R -module is projective.
5. Every short exact sequence of left R -modules splits.
6. R is Artinian and $J(R) = 0$.

§5 Algebra

Definition 5.1. *Let A be an algebra over a commutative ring K with identity.*

1. A **left algebra A -module** is a left K -module M such that M is a left module over the ring A and $k(am) = (ka)m = a(km)$ for all $k \in K, a \in A, m \in M$. Indeed,

$$\begin{cases} (k_1a_1 + k_2a_2)(m_1 + m_2) = k_1a_1m_1 + k_1a_1m_2 + k_2a_2m_1 + k_2a_2m_2 \\ k(am) = (ka)m = a(km) \\ 1_K m = m, 1_K a = a \end{cases}$$

for all $k \in K, a \in A, m \in M$

2. A left algebra A -submodule of M is a subset of M which is itself an left algebra A -module.
3. A left algebra A -module M is **simple** (or **irreducible**) if M has no proper A -submodules.
4. A homomorphism $f : M \rightarrow N$ of algebra A -modules is a map that is both a K -module and an A -module homomorphism.

Remark.

Theorem 5.2. *Let A be a K -algebra. The Jacobson radical of the ring A coincides with the Jacobson radical of the algebra A . In particular A is a semisimple ring if and only if A is a semisimple algebra.*

Theorem 5.3. *Let A be a K -algebra.*

- (1) Every simple algebra A -module is a simple module over the ring A .*
- (2) Every simple module M over the ring A can be given a unique K -module structure in such a way that M is a simple algebra A -module.*

Chapter VIII

Integral

§1 Rings Extensions

Definition 1.1. Let B be a commutative ring with identity and A a subring of B containing 1_B .

1. Then B is said to be an **extension ring** of A .
2. An element $s \in B$ is said to be **integral** over A if s is a root of a monic polynomial in $A[x]$.
3. If every element of B is integral over A , B is said to be an **integral extension** of A .
4. The **integral closure** of A in B is the set of elements of B that are integral over A .
5. The ring A is said to be **integrally closed** in B if A is equal to its integral closure in B .

The integral closure of an integral domain R in its field of fractions is called the **normalization** of R . An integral domain is called integrally closed or normal if it is integrally closed in its field of fractions.

Remark. It follows from corollary 1.3 that the integral closure of R in S is a subring of S containing R .

Theorem 1.2. Let B be an extension ring of A and $s \in B$. Then the following conditions are equivalent.

1. s is integral over A
2. Subring $A[s]$ is a finitely generated A -module
3. There is a subring T that $A[s] \subset T \subset B$, which is finitely generated as an A -module;
4. There is a faithful $A[s]$ -submodule M which is finitely generated as an A -module.

Corollary 1.3. Let B be an extension ring of A . Then

1. If B is finitely generated as an A -module, then B is an integral extension of A .

2. If $s_1, \dots, s_t \in B$ are integral over A , then $A[s_1, \dots, s_t]$ is a finitely generated A -module and an integral extension ring of A .
3. If T is an integral extension ring of B and B is an integral extension ring of A , then T is an integral extension ring of A .

Proof. It immediately follows from 1.2 □

Proof. We have a tower of extension rings:

$$R \subset R[s_1] \subset R[s_1, s_2] \subset \cdots \subset R[s_1, \dots, s_t]$$

For each i , s_i is integral over R and hence integral over $R[s_1, \dots, s_{i-1}]$. Since $R[s_1, \dots, s_i] = R[s_1, \dots, s_{i-1}][s_i]$, $R[s_1, \dots, s_i]$ is a finitely generated module over $R[s_1, \dots, s_{i-1}]$ by 1.2. Thus $R[s_1, \dots, s_n]$ is a finitely generated R -module, then $R[s_1, \dots, s_n]$ is an integral extension ring of R by 1. □

Proof. T is obviously an extension ring of R . If $t \in T$, then t is integral over S and therefore the root of some monic polynomial $f \in S[x]$, say $f = \sum_{i=0}^n s_i x^i$. Since f is also a polynomial over the ring $R[s_0, s_1, \dots, s_{n-1}]$, t is integral over $R[s_0, \dots, s_{n-1}]$.

By 1.2 $R[s_0, \dots, s_{n-1}][t]$ is a finitely generated $R[s_0, \dots, s_{n-1}]$ -module. But since S is integral over R , $R[s_0, \dots, s_{n-1}]$ is a finitely generated R -module by 2. Then

$$R[s_0, \dots, s_{n-1}][t] = R[s_0, \dots, s_{n-1}, t]$$

is a finitely generated R -module. Since $R[t] \subset R[s_0, \dots, s_{n-1}, t]$, t is integral over R by 1.2. □

Proposition 1.4. 1. Every unique factorization domain is integrally closed.

2. In particular, the polynomial ring $F[x_1, \dots, x_n]$ (F a field) is integrally closed in its quotient field $F(x_1, \dots, x_n)$.

§1.1 Integral extension

Theorem 1.5. Let T be a multiplicative subset of an integral domain R such that $0 \notin T$. If R is integrally closed, then $T^{-1}R$ is an integrally closed integral domain.

Proof. $T^{-1}R$ is an integral domain and R may be identified with a subring of $T^{-1}R$ by 1.2. Extending this identification, the quotient field $Q(R)$ of R may be considered as a subfield of the quotient field $Q(T^{-1}R)$ of $T^{-1}R$. Verify that $Q(R) = Q(T^{-1}R)$.

Let $u \in Q(T^{-1}R)$ be integral over $T^{-1}R$; then for some $r_i \in R$ and $s_i \in T$,

$$u^n + (r_{n-1}/s_{n-1})u^{n-1} + \cdots + (r_1/s_1)u + (r_0/s_0) = 0.$$

Multiply through this equation by s^n , where $s = s_0 s_1 \cdots s_{n-1} \in T$, and conclude that su is integral over R . Since $su \in Q(T^{-1}R) = Q(R)$ and R is integrally closed, $su \in R$. Therefore, $u = su/s \in T^{-1}R$, whence $T^{-1}R$ is integrally closed. $\square$

Theorem 1.6. *Let S be an integral extension ring of R . Then the following statements hold.*

1. *Assume that S is an integral domain. Then R is a field if and only if S is a field.*
2. *Let $\mathfrak{p}$ be a prime ideal in R . Then there is a prime ideal $\mathfrak{q}$ in S with $\mathfrak{p} = \mathfrak{q} \cap R$.
Moreover, $\mathfrak{p}$ is maximal if and only if $\mathfrak{q}$ is maximal.*
3. *(The Going-up Theorem) Let $\mathfrak{p}_1 \subseteq \mathfrak{p}_2 \subseteq \cdots \subseteq \mathfrak{p}_n$ be a chain of prime ideals in R and suppose there are prime ideals $\mathfrak{q}_1 \subseteq \mathfrak{q}_2 \subseteq \cdots \subseteq \mathfrak{q}_m$ of S with $\mathfrak{p}_i = \mathfrak{q}_i \cap R$, $1 \leq i \leq m$ and $m < n$. Then the ascending chain of ideals can be completed: there are prime ideals $\mathfrak{q}_{m+1} \subseteq \cdots \subseteq \mathfrak{q}_n$ in S such that $\mathfrak{p}_i = \mathfrak{q}_i \cap R$ for all i .*

Theorem 1.7 (The Going-down Theorem). *Assume that S is an integral domain and R is integrally closed in S . Let $\mathfrak{p}_1 \supseteq \mathfrak{p}_2 \supseteq \cdots \supseteq \mathfrak{p}_n$ be a chain of prime ideals in R and suppose there are prime ideals $\mathfrak{q}_1 \supseteq \mathfrak{q}_2 \supseteq \cdots \supseteq \mathfrak{q}_m$ of S with $\mathfrak{p}_i = \mathfrak{q}_i \cap R$, $1 \leq i \leq m$ and $m < n$. Then the descending chain of ideals can be completed: there are prime ideals $\mathfrak{q}_{m+1} \supseteq \cdots \supseteq \mathfrak{q}_n$ in S such that $\mathfrak{p}_i = \mathfrak{q}_i \cap R$ for all i .*

Theorem 1.8. *Let S be an integral extension ring of R and let $\mathfrak{q}$ be a prime ideal in S which lies over a prime ideal $\mathfrak{p}$ in R . Then $\mathfrak{q}$ is maximal in S if and only if $\mathfrak{p}$ is maximal in R .*

Proof. Suppose $\mathfrak{q}$ is maximal in S . By ?? there is a maximal ideal $\mathfrak{m}$ of R that contains $\mathfrak{p}$ and $\mathfrak{m}$ is prime by ?. By ?? there is a prime ideal $\mathfrak{q}'$ in S such that $\mathfrak{q} \subset \mathfrak{q}'$ and $\mathfrak{q}'$ lies over $\mathfrak{m}$. Since $\mathfrak{q}'$ is prime, $\mathfrak{q}' \neq S$. The maximality of $\mathfrak{q}$ implies that $\mathfrak{q} = \mathfrak{q}'$, whence $\mathfrak{p} = \mathfrak{q} \cap R = \mathfrak{q}' \cap R = \mathfrak{m}$. Therefore, $\mathfrak{p}$ is maximal in R .

Conversely suppose $\mathfrak{p}$ is maximal in R . Since $\mathfrak{q}$ is prime in S , $\mathfrak{q} \neq S$ and there is a maximal ideal N of S containing $\mathfrak{q}$ and N is prime, whence $1_R = 1_S \notin N$. Since $\mathfrak{p} = R \cap \mathfrak{q} \subset R \cap N \subset R$, we must have $\mathfrak{p} = R \cap N$ by maximality. Thus $\mathfrak{q}$ and N both lie over $\mathfrak{p}$ and $\mathfrak{q} \subset N$. Therefore, $\mathfrak{q} = N$ by 1.8. $\square$

§2 Discrete Valuation Ring

Definition 2.1. *The following conditions on a principal ideal domain are equivalent:*

1. *A has exactly one nonzero prime ideal;*
2. *up to associates, A has exactly one prime element;*
3. *A is local and is not a field.*

*A ring satisfying these conditions is called a **discrete valuation ring**.*

Theorem 2.2. *An integral domain A is a discrete valuation ring if and only if*

- (i) A is noetherian,
- (ii) A is integrally closed, and
- (iii) A has exactly one nonzero prime ideal.

§3 Dedekind Domain

Definition 3.1. *A **Dedekind domain** is an integral domain R satisfying the following equivalent conditions:*

1. R is Noetherian, integrally closed and has Krull dimension one (Every nonzero prime ideal of R is maximal).
2. Every nonzero ideal of R is invertible
3. Every finitely generated torsion-free R -module is free.
4. the localization $R_{\mathfrak{p}}$ at each prime ideal $\mathfrak{p}$ of R is a discrete valuation ring.
5. Every nonzero proper ideal of R can be written as a product of prime ideals of R , and this factorization is unique up to the order of the factors.

Proposition 3.2. *Let A be an integral domain, and let S be a multiplicative subset of A .*

1. *If A is noetherian, then so also is $S^{-1}A$.*
2. *If A is integrally closed, then so also is $S^{-1}A$.*
3. *If A has Krull dimension one, then so also does $S^{-1}A$.*
4. *If A is a Dedekind domain, then so also is $S^{-1}A$.*

Proposition 3.3. *A noetherian integral domain A is a Dedekind domain if and only if, for every nonzero prime ideal $\mathfrak{p}$ in A , the localization $A_{\mathfrak{p}}$ is a discrete valuation ring.*

§3.1 Unique factorization of ideals

Theorem 3.4. *Let A be a Dedekind domain. Every proper nonzero ideal $\mathfrak{a}$ of A can be written in the form*

$$\mathfrak{a} = \mathfrak{p}_1^{r_1} \cdots \mathfrak{p}_n^{r_n}$$

with the $\mathfrak{p}_i$ distinct prime ideals and the $r_i > 0$; the $\mathfrak{p}_i$ and the r_i are uniquely determined.

§3.2 The ideal class group

Definition 3.5. Let R be an integral domain with quotient field K . A **fractional ideal** of R is

- (i) a nonzero R -submodule I of K
- (ii) there exists a nonzero $d \in R$ such that $dI \subset R$ i.e., $(R : I) \cap R \neq \emptyset$

Definition 3.6. If R is an integral domain with quotient field K , then the set of all fractional ideals of R forms a commutative monoid, with identity R and multiplication given by

$$IJ = \left\{ \sum_{i=1}^n a_i b_i \mid a_i \in I; b_i \in J; n \in \mathbb{Z}_{\geq 1} \right\}$$

A fractional ideal I of an integral domain R is said to be **invertible** if $IJ = R$ for some fractional ideal J of R .

Theorem 3.7. Let A be a Dedekind domain. The set $\text{Id}(A)$ of fractional ideals is a group; in fact, it is the free abelian group on the set of nonzero prime ideals.

Definition 3.8. We define the **ideal class group** $\text{Cl}(A)$ of A to be the quotient $\text{Cl}(A) = \text{Id}(A)/\text{P}(A)$ of $\text{Id}(A)$ by the subgroup of principal ideals. The **class number** of A is the order of $\text{Cl}(A)$ (when finite).

In the case that A is the ring of integers $\mathcal{O}_K$ in a number field K , we often refer to $\text{Cl}(\mathcal{O}_K)$ as the ideal class group of K , and its order as the class number of K .

Proposition 3.9. Let R be an integral domain with quotient field K .

1. Indeed for any fractional ideal I the set $I^{-1} = \{a \in K \mid aI \subset R\}$ is easily seen to be a fractional ideal such that $I^{-1}I = II^{-1} \subset R$.
2. The inverse of an invertible fractional ideal I is unique and is $I^{-1} = \{a \in K \mid aI \subset R\}$. If I is invertible and $IJ = JI = R$, then clearly $J \subset I^{-1}$. Conversely, since I^{-1} and J are R -submodules of K , $I^{-1} = RI^{-1} = (JI)I^{-1} = J(II^{-1}) \subset JR = RJ \subset J$, whence $J = I^{-1}$.
3. If I, A, B are fractional ideals of R such that $IA = IB$ and I is invertible, then $A = RA = (I^{-1}I)A = I^{-1}(IB) = RB = B$.
4. If I is an ordinary ideal in R , then $R \subset I^{-1}$.

Lemma 3.10. Let $I, I_1, I_2, \dots, I_n$ be ideals in an integral domain R .

1. The ideal $I_1 I_2 \cdots I_n$ is invertible if and only if each I_j is invertible.
2. If $\mathfrak{p}_1 \cdots \mathfrak{p}_m = I = \mathfrak{q}_1 \cdots \mathfrak{q}_n$, where the $\mathfrak{p}_i$ and $\mathfrak{q}_j$ are prime ideals in R and every $\mathfrak{p}_i$ is invertible, then $m = n$ and (after reindexing) $\mathfrak{p}_i = \mathfrak{q}_i$ for each $i = 1, \dots, m$.

Lemma 3.11. *If I is a fractional ideal of an integral domain R with quotient field K and $f \in \text{Hom}_R(I, R)$, then for all $a, b \in I : af(b) = bf(a)$.*

Lemma 3.12. *Every invertible fractional ideal of an integral domain R with quotient field K is a finitely generated R -module.*

Theorem 3.13. *Let R be an integral domain and I a fractional ideal of R . Then I is invertible if and only if I is a projective R -module.*

§4 Discrete valuations

Definition 4.1. *Let K be a field. A **discrete valuation** on K is a nonzero homomorphism $v : K^\times \rightarrow \mathbb{Z}$ such that $v(a + b) \geq \min(v(a), v(b))$.*

*As v is not the zero homomorphism, its image is a nonzero subgroup of $\mathbb{Z}$, and is therefore of the form $m\mathbb{Z}$ for some $m \in \mathbb{Z}$. If $m = 1$, then $v : K^\times \rightarrow \mathbb{Z}$ is surjective, and v is said to be **normalized**; otherwise, $x \mapsto m^{-1} \cdot v(x)$ will be a normalized discrete valuation.*

We extend v to a map $K \rightarrow \mathbb{Z} \cup \{\infty\}$ by setting $v(0) = \infty$, where ∞ is a symbol $\geq n$ for all $n \in \mathbb{Z}$.

Remark. *We have*

1. $v(\zeta) = 0$ for some $\zeta \in K^\times$
2. $v(-a) = v(a)$ for all $a \in K$;
3. $v(a + b) = \max\{v(a), v(b)\}$ if $v(a) \neq v(b)$.

We often use "ord" rather than " v " to denote a discrete valuation.

Proposition 4.2. *Let v be a discrete valuation on K , then*

$$A := \{a \in K \mid v(a) \geq 0\}$$

is a principal ideal domain with maximal ideal

$$\mathfrak{m} = \{a \in K \mid v(a) > 0\}$$

If $v(K^\times) = m\mathbb{Z}$, then the ideal $\mathfrak{m}$ is generated by every element of $v^{-1}(m)$.

Definition 4.3. *Let A be a Dedekind domain and let $\mathfrak{p}$ be a prime ideal in A . For any $c \in K^\times$, let $v(c)$ be the exponent of $\mathfrak{p}$ in the factorization of (c) . Then v is a normalized discrete valuation on K , called the **discrete valuation associated to $\mathfrak{p}$** , denoted by $\text{ord}_{\mathfrak{p}}$.*

Proposition 4.4. *Let $x_1, \dots, x_m$ be elements of a Dedekind domain A , and let $\mathfrak{p}_1, \dots, \mathfrak{p}_m$ be distinct prime ideals of A . For every integer n , there is an $x \in A$ such that*

$$\text{ord}_{\mathfrak{p}_i}(x - x_i) > n, \quad i = 1, 2, \dots, m.$$

§5 Integral closures of Dedekind domains

Theorem 5.1. *Let A be a Dedekind domain with field of fractions K and L/K be a finite separable extension, then the integral closure of A in L is Dedekind domain.*

Definition 5.2. *Let A be a Dedekind domain with field of fractions K , and let B be the integral closure of A in a finite separable extension L of K . A prime ideal $\mathfrak{p}$ of A will factor in B ,*

$$\mathfrak{p}B = \mathfrak{P}_1^{e_1} \cdots \mathfrak{P}_g^{e_g}$$

where $\mathfrak{P}$ are distinct prime ideals in B and $e_i \geq 1$,

1. If any of the numbers is > 1 , then we say that $\mathfrak{p}$ is **ramified** in B (or L). The number e_i is called the **ramification index**.

2. We say $\mathfrak{P}$ divides $\mathfrak{p}$, written $\mathfrak{P} \mid \mathfrak{p}$, if $\mathfrak{P}$ occurs in the factorization of $\mathfrak{p}$ in B .

We then write $e(\mathfrak{P}/\mathfrak{p})$ for the ramification index and $f(\mathfrak{P}/\mathfrak{p})$ for the degree of the field extension $[B/\mathfrak{P} : A/\mathfrak{p}]$ (called the **residue class degree**).

3. $\mathfrak{p}$ is said to **split** (or split completely) in L if $e_i = f_i = 1$ for all i

4. $\mathfrak{p}$ is said to be **inert** in L if $\mathfrak{p}B$ is a prime ideal (so $g = 1 = e$).

Theorem 5.3. *Let m be the degree of L over K , and let $\mathfrak{P}_1, \dots, \mathfrak{P}_g$ be the prime ideals dividing $\mathfrak{p}$; then*

$$\sum_{i=1}^g e_i f_i = m$$

where $e_i = e(\mathfrak{P}_i/\mathfrak{p})$ and $f_i = f(\mathfrak{P}_i/\mathfrak{p})$. If L is Galois over K , then all the ramification numbers are equal, and all the residue class degrees are equal, and so

$$efg = m.$$

Chapter IX

Completions

Contents

§1		37
§2	Graded rings and graded modules	37
§2.1	Graded ring	37
§2.2	Graded modules	38
§2.3	Graded algebra	39
§3	Filtrations	39
§4	Associated graded ring	40

§1

§2 Graded rings and graded modules

§2.1 Graded ring

Definition 2.1. A *graded ring* is a ring A together with a family $\{A_n\}_{n \geq 0}$ of subgroups of A , such that

(i) $A = \bigoplus_{n=0}^{\infty} A_n$ in $\mathbf{Ab}$

(ii) and $A_m A_n \subseteq A_{m+n}$ for all $m, n \geq 0$.

Thus A_0 must be a subring of A containing 1_A , and A is a A_0 -algebra. A is said to be **standard graded ring** if

(iii) $A = A_0[A_1]$ as a A_0 -algebra.

Proposition 2.2. Let $A = \bigoplus_{n \geq 0} A_n$ be a graded ring. Then the following statements are equivalent:

1. A is a Noetherian ring.
2. A_0 is Noetherian and A is finitely generated as A_0 -algebra (A_0 -ring)

Definition 2.3. Let $A = \bigoplus A_n$ be a graded ring.

1. A ideal (left, right, two-sided) I of A is called **graded ideal** if $I = \bigoplus_{n \geq 0} (A_n \cap I)$ in **Ab**.
2. If I is a two-sided graded ideal of A , then quotient graded ring $A/I = \bigoplus_{i \geq 0} A_i / (A_i \cap I)$ is well-defined, called the **quotient graded ring** of A by I .

Remark. It clear that a ideal I is graded if and only if it is generated by homogeneous elements.

Definition 2.4. Let A and B be graded rings.

1. A **graded ring homomorphism** from A to B is a ring homomorphism $f : A \rightarrow B$ such that $f(A_n) \subseteq B_n$ for all $n \geq 0$.
2. the **kernel** of a graded ring homomorphism is a two-sided graded ideal.

§2.2 Graded modules

Definition 2.5. Let $A = \bigoplus_{n \geq 0} A_n$ be a graded ring. A **graded left A -module** is a left A -module M together with a family $\{M_n\}_{n \geq 0}$ of subgroups of M such that

- (i) $M = \bigoplus_{n=0}^{\infty} M_n$ as an abelian group
- (ii) $A_m M_n \subseteq M_{m+n}$ for all $m, n \geq 0$.

Elements of M_n are called **homogeneous elements of degree n** .

Remark. Thus each M_n is an A_0 -module.

Definition 2.6. Let A be a graded ring and M, N graded A -modules.

1. A **graded submodule** of M is a submodule N of M such that $N = \bigoplus_{n \geq 0} (N \cap M_n)$ in **Ab**.
2. then the $M/N = \bigoplus_{n \geq 0} M_n / (N \cap M_n)$ is well-defined, called the **quotient graded module** of M by N .
3. A **graded A -homomorphism** from M to N is an A -module homomorphism $f : M \rightarrow N$ such that $f(M_n) \subseteq N_n$ for all $n \geq 0$.

§2.3 Graded algebra

Definition 2.7. Let k be a commutative ring (usually a field). A **graded k -algebra** is k -algebra A such that

- (i) $A = \bigoplus_{n \geq 0} A_n$ is a graded ring.
- (ii) the structure morphism $\varphi : k \rightarrow A$ with $\text{Im}(\varphi) \subset A_0$ (scalar action preserves degrees).

Remark. In this case, A_0 is naturally a k -algebra, and A may be viewed as a graded A_0 -algebra with base ring A_0 instead of k . In many situations one further assumes that the base ring $k = A_0$.

We give a graded ring of particular importance in commutative algebra.

Definition 2.8. Let k be commutative ring (usually a field). A **standard graded k -algebra** is a graded k -algebra $A = \bigoplus_{n \geq 0} A_n$ such that A is generated by A_1 as a k -algebra.

§3 Filtrations

Definition 3.1. Let A be a ring. A descending **filtration** of A one means a sequence of subgroups

$$A = A_0 \supset A_1 \supset A_2 \supset \cdots \supset A_n \supset \cdots$$

such that $A_m A_n \subseteq A_{m+n}$ for all $m, n \geq 0$. (thus each A_n is a two-sided ideal of A).

If $\mathfrak{a}$ is an ideal of A , then the $\{\mathfrak{a}^n\}$ is called the **$\mathfrak{a}$ -adic filtration**.

Definition 3.2. Let A be a filtered ring with descending filtration $\mathcal{F} = \{A_i\}$. A left A -module M is called a **filtered left A -module** if it has a sequence of subgroups (hence submodules) $\{M_i\}$ such that

- (i) $M = M_0 \supset M_1 \supset M_2 \supset \cdots \supset M_n \supset \cdots$
- (ii) $A_i \cdot M_j \subseteq M_{i+j}$ for all i, j .

The $\{M_i\}$ is called a **filtration** of M with respect to $\mathcal{F}$.

Similarly, $\mathfrak{a}$ -adic filtration of M is the filtration $\{\mathfrak{a}^n M\}$.

Definition 3.3. Let A be a ring, $\mathfrak{a}$ an ideal and M an A -module with a filtration $\{M_n\}$.

1. We say that it is an **$\mathfrak{a}$ -filtration** if $\mathfrak{a}M_n \subseteq M_{n+1}$ for all n .
2. We say that an $\mathfrak{a}$ -filtration is **$\mathfrak{a}$ -stable** if we have $\mathfrak{a}M_n = M_{n+1}$ for all n sufficiently large.

Lemma 3.4. If $\{M_n\}, \{M'_n\}$ are stable $\mathfrak{a}$ -filtrations of M , then they have bounded difference: that is, there exists an integer n_0 such that $M_{n+n_0} \subseteq M'_n$ and $M'_{n+n_0} \subseteq M_n$ for all $n \geq 0$.

Proof. Enough to take $M'_n = \mathfrak{a}^n M$. Since $\mathfrak{a}M_n \subseteq M_{n+1}$ for all n , we have $\mathfrak{a}^n M \subseteq M_n$; also $\mathfrak{a}M_n = M_{n+1}$ for all $n \geq n_0$ say, hence $M_{n+n_0} = \mathfrak{a}^n M_{n_0} \subseteq \mathfrak{a}^n M$. $\square$

Lemma 3.5. *Let A be a Noetherian ring, M a finitely-generated A -module, (M_n) an $\mathfrak{a}$ -filtration of M . Then the following are equivalent:*

1. $M^* = \bigoplus_{n \geq 0} M_n$ is a finitely-generated A^* -module;
2. The filtration (M_n) is stable.

Proof. Each M_n is finitely generated, hence so is each $Q_n = \bigoplus_{r=0}^n M_r$: this is a subgroup of M^* but not (in general) an A^* -submodule. However, it generates one in M^* , namely

$$M_n^* = M_0 \oplus \cdots \oplus M_n \oplus \mathfrak{a}M_n \oplus \mathfrak{a}^2M_n \oplus \cdots \oplus \mathfrak{a}^rM_n \oplus \cdots$$

Since Q_n is finitely generated as an A -module, M_n^* is finitely generated as an A^* -module. The M_n^* form an ascending chain, whose union is M^* . Since A^* is Noetherian, M^* is finitely generated as an A^* -module $\Leftrightarrow$ the chain stops, i.e., $M^* = M_{n_0}^*$ for some $n_0 \Leftrightarrow M_{n_0+r} = \mathfrak{a}^r M_{n_0}$ for all $r \geq 0 \Leftrightarrow$ the filtration is stable. $\square$

Proposition 3.6 (Artin-Rees). *Let A be a Noetherian ring, $\mathfrak{a}$ an ideal, M a finitely generated A -module and $\{M_n\}$ a stable $\mathfrak{a}$ -filtration. If M' is a submodule of M , then $\{M' \cap M_n\}$ is a stable $\mathfrak{a}$ -filtration of M' .*

Proof. We have $\mathfrak{a}(M' \cap M_n) \subseteq \mathfrak{a}M' \cap \mathfrak{a}M_n \subseteq M' \cap M_{n+1}$, hence $(M' \cap M_n)$ is an $\mathfrak{a}$ -filtration. Hence it defines a graded A^* -module which is a submodule of M^* and therefore finitely generated. $\square$

Corollary 3.7. *There exists an integer s such that for all integers $n \geq s$ we have*

$$\mathfrak{a}^n M \cap M' = \mathfrak{a}^{n-s} (\mathfrak{a}^s M \cap M')$$

Corollary 3.8. *Let A be a Noetherian ring, $\mathfrak{a}$ an ideal, M a finitely-generated A -module and M' a submodule of M . Then the $\mathfrak{a}$ -topology of M' coincides with the topology induced by the $\mathfrak{a}$ -topology of M .*

§4 Associated graded ring

Definition 4.1. *Let A be a ring with filtration $\mathcal{F}$, and M an filtered A -module with respect to $\mathcal{F}$.*

1. We define **associated graded ring** of A with respect to $\mathcal{F}$

$$\mathrm{gr}_{\mathcal{F}}(A) = \bigoplus_{n=0}^{\infty} A_n / A_{n+1}$$

2. We define **associated graded module** of A with respect to $\mathcal{F}$

$$\mathrm{gr}(M) = \bigoplus_{n=0}^{\infty} M_n / M_{n+1}$$

then $\mathrm{gr}(M)$ is a graded $\mathrm{gr}_{\mathcal{F}}(A)$ -module.

Proposition 4.2. *Let A be a Noetherian ring and $\mathfrak{a}$ an ideal. Then*

1. $\mathrm{gr}_{\mathfrak{a}}(A)$ is Noetherian.
2. *If M is a finitely generated A -module with a stable $\mathfrak{a}$ -filtration, then $\mathrm{gr}(M)$ is a finitely generated graded $\mathrm{gr}_{\mathfrak{a}}(A)$ -module.*

Chapter X

Dimension

§1 Hilbert Functions

Definition 1.1. Let $\mathcal{A}$ be an abelian category. An **additive function** $\lambda : \mathcal{A} \rightarrow \mathbf{Ab}$ is a covariant functor such that for every short exact sequence

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

in $\mathcal{A}$, we have

$$\lambda(M) = \lambda(M') + \lambda(M'')$$

In the following, $\lambda(N)$ is a finitely generated abelian group and we always view $\lambda(N)$ as the rank of $\lambda(N)$.

Definition 1.2. Let $A = \bigoplus_{n \geq 0} A_n$ be a commutative Noetherian graded ring and $M = \bigoplus_{n \geq 0} M_n$ a finitely generated graded A -module (It follows that each M_n is a finitely generated A_0 -module with an additive function λ).

1. The **Hilbert function** of M with respect to λ is the function defined by

$$h_M^\lambda(n) := \lambda(M_n)$$

2. the **Hilbert-Poincaré series** of M

$$P_M^\lambda(t) := \sum_{n=0}^{\infty} h_M^\lambda(n) t^n$$

Remark. If A_0 is an Artinian ring, $\lambda(N) = \ell_{A_0}(N)$. If A_0 is a field, $\lambda(N) = \dim_{A_0}(N)$.

Theorem 1.3 (Hilbert-Serre). The Poincaré series $P_M^\lambda(t)$ is a rational function of the form

$$\frac{f(t)}{\prod_{i=1}^d (1 - t^{k_i})}$$

where $f(t) \in \mathbb{Z}[t]$ and $k_1, \dots, k_d$ are the degrees of the homogeneous generators of A over A_0 . The order of the pole of $P_M^\lambda(t)$ at $t = 1$ is called the **dimension** of M with respect to λ , denoted by $\dim M$.

Proof. By induction on s , the number of generators of R over R_0 . Start with $s = 0$; this means that $R_n = 0$ for all $n > 0$, so that $R = R_0$ and M is a finitely-generated R_0 module, hence $M_n = 0$ for all large n . Thus $P_M^\lambda(t)$ is a polynomial in this case.

Now suppose $s > 0$ and the theorem true for $s - 1$. Multiplication by x_s is an R -module homomorphism of M_n into M_{n+k_s} , hence it gives an exact sequence, say

$$0 \rightarrow K_n \rightarrow M_n \xrightarrow{x_s} M_{n+k_s} \rightarrow L_{n+k_s} \rightarrow 0.$$

Let $K = \bigoplus_n K_n, L = \bigoplus_n L_n$; these are both finitely-generated A -modules (because K is a submodule and L a quotient module of M), and both are annihilated by x_s , hence they are $A_0[x_1, \dots, x_{s-1}]$ -modules. Applying λ to (1) we have, by (2.11)

$$\lambda(K_n) - \lambda(M_n) + \lambda(M_{n+k_s}) - \lambda(L_{n+k_s}) = 0;$$

multiplying by t^{n+k_s} and summing with respect to n we get

$$(1 - t^{k_s}) P(M, t) = P(L, t) - t^{k_s} P(K, t) + g(t)$$

where $g(t)$ is a polynomial. Applying the inductive hypothesis the result now follows. $\square$

Corollary 1.4 (Existence of the Hilbert polynomial). *Then there exists a polynomial $H(t) \in \mathbb{Q}[t]$ of degree $\dim M$ such that $H(n) = h_M^\lambda(n)$ for all sufficiently large integers n .*

*This polynomial is called the **Hilbert polynomial** of M with respect to λ .*

Corollary 1.5. *There exists a polynomial $\chi(t, A, M) \in \mathbb{Q}[t]$ such that*

$$\sum_{i=0}^n \lambda(M_i) = \chi(n)$$

for all sufficiently large integers n .

Definition 1.6. *The **multiplicity** of M with respect to λ is defined to be*

$$e(M, \lambda) := d! \cdot a_{d(M)}$$

where $a_{d(M)}$ is the leading coefficient of the Hilbert polynomial $\chi(t)$.

§2 Krull Dimension

In this section, all rings A are commutative Noetherian ring with identity and modules M are finitely generated A -module. Although some definitions and results hold in greater generality, the

most of useful properties stay in this case.

Definition 2.1. *Let A -module M .*

1. The **Krull dimension** of A is defined to be the supremum of the lengths n of all chains of prime ideals

$$\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n$$

say $\dim(A) = n$ if the supremum is attained, and $\dim(A) = \infty$ otherwise.

2. The **Krull dimension** of A -module M is defined to be the dimension of the quotient ring $A/\text{Ann}(M)$, that is,

$$\dim(M) := \dim(A/\text{Ann}(M))$$

Proposition 2.2. .

1. $\dim A = \dim \text{Spec } A$
2. If A is a local ring with maximal ideal $\mathfrak{m}$, then $\dim A = \sup \{n : \mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n = \mathfrak{m}\}$
3. if S is a multiplicatively closed subset of A , then $\dim S^{-1}A \leq \dim A$

Definition 2.3. *Let A be a commutative ring and $\mathfrak{p}$ a prime ideal of A . The **height** of $\mathfrak{p}$ is defined to be*

$$\text{ht}(\mathfrak{p}) := \sup \{n : \mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_n = \mathfrak{p}\}$$

Thus $\dim A = \sup_{\mathfrak{p} \in \text{Spec } A} \text{ht}(\mathfrak{p})$.

Theorem 2.4 (Krull Principal Ideal Theorem). *Let A be a Noetherian commutative ring and a ideal $\mathfrak{a} = (f_1, \dots, f_r)$. If $\mathfrak{p}$ is a minimal element of $\{\mathfrak{q} \in \text{Spec } A : \mathfrak{a} \subset \mathfrak{q}\}$. Then $\text{ht}(\mathfrak{p}) \leq r$.*

Chapter XI

§1

Definition 1.1. The space X is called **Noetherian** if it satisfies the descending chain condition for closed subsets, i.e., every descending chain of closed subsets eventually stabilizes.

Definition 1.2. The space X is called **irreducible** if it cannot be expressed as the union of two proper closed subsets. The subset Y of X is called **irreducible** if it is irreducible as a subspace of X , and empty set is irreducible.

Remark. It is easy to see that X is irreducible if and only if

1. Any two non-empty open subsets of X intersect. Any two non-empty closed subsets of X intersect.
2. Every non-empty open subset of X is dense in X . Every non-empty closed subset of X has empty interior.

§2 Spectrum of a Ring

Definition 2.1. Let A be a commutative ring with identity. The **spectrum** of A , denoted by

$$\operatorname{Spec} A := \{\mathfrak{p} : \mathfrak{p} \text{ is a prime ideal of } A\}$$

, is the set of all prime ideals of A . If $\mathfrak{a}$ is an ideal of A , then the **vanishing (zeros) set** of $\mathfrak{a}$ is defined to be

$$V(\mathfrak{a}) := \{\mathfrak{p} \in \operatorname{Spec} A : \mathfrak{a} \subset \mathfrak{p}\} \subset \operatorname{Spec} A$$

The **Zariski topology** on $\operatorname{Spec} A$ is defined by taking the collection of all vanishing sets $V(\mathfrak{a})$ as the closed sets.

Definition 2.2. Defining a sheaf $\mathcal{O}$ on $X = \operatorname{Spec} A$, for each open set $U \subseteq X$, we define $\mathcal{O}(U)$ be the set of functions

$$s : U \rightarrow \prod_{\mathfrak{p} \in U} A_{\mathfrak{p}}, \quad \text{or simplified } (s_{\mathfrak{p}})_{\mathfrak{p} \in U}$$

such that $s(\mathfrak{p}) \in A_{\mathfrak{p}}$ for all $\mathfrak{p} \in U$, and such that s is locally a quotient of elements of A , i.e., for each $\mathfrak{p} \in U$, there exists an open neighborhood $V \subseteq U$ and elements $a, f \in A$ such that $f \notin \mathfrak{q}$ for all $\mathfrak{q} \in V$ and $s|_V = a/f$.

Definition 2.3. Let A be a commutative ring with identity. The **spectrum** of A is the pair consisting of the topological space $\text{Spec } A$ and the sheaf of rings $\mathcal{O}$ defined above.

Definition 2.4. Let $f \in A$, the **distinguished open set** of f is defined to be

$$D(f) := \{\mathfrak{p} \in \text{Spec } A : f \notin \mathfrak{p}\} = \text{Spec } A - V(f)$$