

- Q.27 What is a Carnot cycle? List the four stages of the Carnot cycle.
- Q.28 Differentiate between single and multiple reactions with one example of each.
- Q.29 What is a second-order reaction? Give an example involving two reactants.
- Q.30 State the formula for the efficiency of a Carnot engine. Explain what each term represents.
- Q.31 What is activation energy? Describe its role in a chemical reaction.
- Q.32 What are the main differences between a batch reactor and a continuous reactor?
- Q.33 Draw a simple graph comparing plus flow, mixed flow and batch reactor.
- Q.34 State and explain Dalton's law and Henry's law.
- Q.35 Discuss any one of the following
(i) Vander Waal's Equation (ii) Amagat's Law

Section-D

Note: Long answer questions. Attempt any two question out of three Questions. (2x10=20)

- Q.36 Explain the basic construction and working principle of a Plug Flow Reactor with its neat sketch.
- Q.37 Calculate the efficiency of a Carnot engine operating between a hot reservoir at 500 K and a cold reservoir at 300K.
- Q.38 Define and differentiate between a closed system, an open system, and an isolated system. Provide one practical example for each type of system.

No. of Printed Pages : 4
Roll No.

180642/120642/030642

4th Sem.

Branch: CHEMICAL ENGINEERING (P&P)

Sub : CTRE

Time : 3 Hrs.

M.M. : 100

SECTION-A

Note: Multiple type Questions. All Questions are compulsory. (10x1=10)

- Q.1 Which of the following is an extensive property?
a) Temperature b) Pressure
c) Volume d) Density
- Q.2 Which of the following is NOT a state function?
a) Internal Energy b) Enthalpy
c) Heat d) Entropy
- Q.3 The zeroth law of thermodynamics is the basis for the measurement of:
a) Temperature b) Pressure
c) Volume d) Energy
- Q.4 What is the unit of entropy?
a) Joules b) Joules per Kelvin
c) Joules per mole d) Kelvin
- Q.5 Henry's Law is used to describe:
a) The pressure of an ideal gas
b) The solubility of gases in liquids
c) The behavior of real gases
d) The phase diagram of a substance

- Q.6 The equilibrium constant of a reaction is affected by:
- Temperature only
 - Pressure only
 - Both temperature & pressure
 - Neither temperature nor pressure
- Q.7 Activation energy of a reaction is :
- The minimum energy required for a reaction to occur
 - The energy released during a reaction
 - The total energy of reactants
 - The energy difference between reactants and products
- Q.8 A batch reactor is characterized by :
- Continuous input and output of reactants and products
 - No flow in or out during the reaction
 - Perfect mixing of reactants and products
 - Steady-state operation
- Q.9 Molecularity of a reaction refers to :
- The order of the reaction
 - The number of molecules involved in the reaction step
 - The rate of the reaction
 - The activation energy of the reaction
- Q.10 What defines an elementary reaction?
- A reaction involving multiple steps
 - A reaction that occurs in a single step
 - A reaction that reaches equilibrium
 - A reaction that does not involve any intermediates

Section-B

Note: Objective type questions. All questions are compulsory. (10x1=10)

- Q.11 Expand MFR.
- Q.12 Define entropy.
- Q.13 What is isobaric process.
- Q.14 Define molecularity of reaction.
- Q.15 Give one example of open system.
- Q.16 Write ideal gas equation.
- Q.17 What is state function.
- Q.18 What is equilibrium.
- Q.19 Write Raoult's law.
- Q.20 Write any two scales of temperature measurement.

Section-C

Note: Short answer type Question. Attempt any twelve questions out of fifteen Questions. (12x5=60)

- Q.21 Explain the difference between intensive and extensive properties. Provide one example of each.
- Q.22 State the Zeroth Law of Thermodynamics and explain its significance in temperature measurement.
- Q.23 Define a thermodynamic system and classify it into open, closed, and isolated systems with examples.
- Q.24 Explain the First Law of Thermodynamics for a closed system. With one example.
- Q.25 Derive the ideal gas law from the basic principles and state its assumptions.
- Q.26 Calculate the value of universal gas constant (R) from ideal gas law in $\text{m}^3 \text{ atm/k. mol. K}$.